

Extending the Harper Identity to Iterated Belief Change

Richard Booth
Cardiff University
Cardiff, UK
boothr2@cardiff.ac.uk

Jake Chandler
La Trobe University
Melbourne, Australia
jacob.chandler@latrobe.edu.au

Abstract The field of iterated belief change has focused mainly on *revision*, with the other main operator of AGM belief change theory, i.e., *contraction* receiving relatively little attention. In this paper we extend the Harper Identity from single-step change to define iterated contraction in terms of iterated revision. Specifically, just as the Harper Identity provides a recipe for defining the *belief set* resulting from contracting A in terms of (i) the initial belief set and (ii) the belief set resulting from revision by $\neg A$, we look at ways to define the *plausibility ordering* over worlds resulting from contracting A in terms of (iii) the initial plausibility ordering, and (iv) the plausibility ordering resulting from revision by $\neg A$. After noting that the most straightforward such extension leads to a trivialisation of the space of permissible orderings, we provide a family of operators for combining plausibility orderings that avoid such a result. These operators are characterised in our domain of interest by a pair of intuitively compelling properties, which turn out to enable the derivation of a number of iterated contraction postulates from postulates for iterated revision. We finish by observing that a salient member of this family allows for the derivation of counterparts for contraction of some well known iterated revision operators, as well as for defining new iterated contraction operators.

1 Introduction

Since the publication of Darwiche and Pearl’s seminal paper on the topic in the mid 90’s [Darwiche and Pearl, 1997], a substantial body of research has now accumulated on the problem of iterated belief revision—the problem of how to adjust one’s corpus of beliefs in response to a temporal sequence of successive additions to its members [Booth and Meyer, 2006; 2011; Boutilier, 1996; Jin and Thielscher, 2007; Nayak *et al.*, 2003; Peppas, 2014].

In contrast, work on the parallel problem of iterated *contraction*—the problem of how to adjust one’s corpus in response to a sequence of successive retractions—was only initiated far more recently and remains comparatively underdeveloped

[Chopra *et al.*, 2008; Hansson, 2012; Hild and Spohn, 2008; Nayak *et al.*, 2006; 2007; Ramachandran *et al.*, 2012; Rott, 2009].

One obvious way to level out this discrepancy would be to introduce a principle that enables us to derive, from constraints on iterated revision, corresponding constraints on iterated contraction. But while there exists a well known and widely accepted postulate connecting *single-shot* revision and contraction, the ‘Harper Identity’ [Harper, 1976], there has been no discussion to date of how to extend this principle to the iterated case.¹ One idea, which we pursue in this paper, is that whereas the Harper Identity says the *belief set* resulting from contracting sentence A should be formed by combining (i) the initial belief set and (ii) the belief set resulting from revision by $\neg A$, we look for ways to define the *plausibility ordering* over worlds resulting from contracting A in terms of (iii) the initial plausibility ordering, and (iv) the plausibility ordering resulting from revision by $\neg A$.

In the present paper, we first of all show that the simplest extension of the Harper Identity to iterated belief change is too strong a principle, being inconsistent with basic principles of belief dynamics on pairs of triviality (Section 3). This leads us to consider a set of collectively weaker principles, which we show to characterise, in our domain of interest, a family of binary combination operators for total preorders that we call *TeamQueue* combinators (Section 4). After recapitulating a number of existing postulates from both iterated revision and contraction, we show how these two lists of postulates can be linked via the use of any *TeamQueue* combinator (Section 5). Then we prove some more specific results of this type using a particular *TeamQueue* combinator that we call *Synchronous TeamQueue* (Section 6). Finally we conclude and mention some ideas for future work. Proofs of the various propositions and theorems have been relegated to the appendix.

2 Preliminaries

We represent the beliefs of an agent by a so-called belief state Ψ , which we treat as a primitive. Ψ determines a belief set $[\Psi]$, a deductively closed set of sentences, drawn from a finitely generated propositional, truth-functional language L . The set of classical logical consequences of a sentence $A \in L$ is denoted by $\text{Cn}(A)$. The set of propositional worlds is denoted by W , and the set of models of a given sentence A is denoted by $\llbracket A \rrbracket$.

The dynamics of belief states are modelled by two operations—contraction and revision, which respectively return the posterior belief states $\Psi * A$ and $\Psi \div A$ resulting from an adjustment of the prior belief state Ψ to accommodate, respectively, the inclusion and exclusion of A .

¹It should be noted that [Nayak *et al.*, 2006] and Ramachandran *et al* [Ramachandran *et al.*, 2012] do propose a principle that they call the ‘New Harper Identity’. But while this may be suggestive of an attempted extension of the Harper Identity to the iterated case, the New Harper Identity simply appears to be a representation, in terms of plausibility orderings, of a particular set of postulates for iterated contraction.

We assume that these operations satisfy the so-called AGM postulates [Alchourrón *et al.*, 1985], which enforce a principle of ‘minimal mutilation’ of the initial belief set in meeting the relevant exclusion or inclusion constraint. Regarding revision, we have:

- (AGM*1) $\text{Cn}([\Psi * A]) \subseteq [\Psi * A]$
- (AGM*2) $A \in [\Psi * A]$
- (AGM*3) $[\Psi * A] \subseteq \text{Cn}([\Psi] \cup \{A\})$
- (AGM*4) If $\neg A \notin [\Psi]$, then $\text{Cn}([\Psi] \cup \{A\}) \subseteq [\Psi * A]$
- (AGM*5) If A is consistent, then so too is $[\Psi * A]$
- (AGM*6) If $\text{Cn}(A) = \text{Cn}(B)$, then $[\Psi * A] = [\Psi * B]$
- (AGM*7) $[\Psi * (A \wedge B)] \subseteq \text{Cn}([\Psi * A] \cup \{B\})$
- (AGM*8) If $\neg B \notin [\Psi * A]$, then $\text{Cn}([\Psi * A] \cup \{B\}) \subseteq [\Psi * (A \wedge B)]$

Regarding contraction:

- (AGM÷1) $\text{Cn}([\Psi \div A]) \subseteq [\Psi \div A]$
- (AGM÷2) $[\Psi \div A] \subseteq [\Psi]$
- (AGM÷3) If $A \notin [\Psi]$, then $[\Psi \div A] = [\Psi]$
- (AGM÷4) If $A \notin \text{Cn}(\emptyset)$, then $A \notin [\Psi \div A]$
- (AGM÷5) If $A \in [\Psi]$, then $[\Psi] \subseteq \text{Cn}([\Psi \div A] \cup \{A\})$
- (AGM÷6) If $\text{Cn}(A) = \text{Cn}(B)$, then $[\Psi \div A] = [\Psi \div B]$
- (AGM÷7) $[\Psi \div A] \cap [\Psi \div B] \subseteq [\Psi \div A \wedge B]$
- (AGM÷8) If $A \notin [\Psi \div A \wedge B]$, then $[\Psi \div A \wedge B] \subseteq [\Psi \div A]$

We also assume that they are linked in the one-step case by the Harper Identity (HI):

$$(HI) \quad [\Psi \div A] = [\Psi] \cap [\Psi * \neg A]$$

We follow a number of authors in making use of a ‘semantic’ representation of the ‘syntactic’ one-step revision and contraction dispositions associated with a particular belief state Ψ in terms of a total preorder (*tpo*) \preceq_Ψ over the set W of possible worlds. Intuitively \preceq_Ψ orders the worlds according to *plausibility* (with more plausible worlds lower down the ordering). Then the set $\min(\preceq_\Psi, \llbracket A \rrbracket) := \{x \in \llbracket A \rrbracket \mid \forall y \in \llbracket A \rrbracket, x \preceq_\Psi y\}$ of minimal A -worlds corresponds to the set of worlds in which all and only the sentences in $[\Psi * A]$ are true, with $\llbracket [\Psi] \rrbracket = \min(\preceq_\Psi, W)$ for any Ψ (see, for instance, the representation results in [Grove, 1988; Katsuno and Mendelzon, 1991]). Viewed in this way, the question of iterated belief change becomes a question about the dynamics of \preceq_Ψ under contraction and revision, with HI translating into the constraint $\min(\preceq_{\Psi \div A}, W) = \min(\preceq_\Psi, W) \cup \min(\preceq_{\Psi * \neg A}, W)$. We will denote the set of all *tpo*s over W by $T(W)$. The strict part of \preceq_Ψ will be denoted by \prec_Ψ and its symmetric part by \sim_Ψ .

A tpo \preceq_Ψ can also be represented by an ordered partition $\langle S_1, S_2, \dots, S_m \rangle$ of W , with $x \preceq_\Psi y$ iff $r(x, \preceq_\Psi) \leq r(y, \preceq_\Psi)$, where $r(x, \preceq_\Psi)$ denotes the ‘rank’ of x with respect to \preceq_Ψ and is defined by taking $S_{r(x, \preceq_\Psi)}$ to be the cell in the partition that contains x .

3 A triviality result

What should an agent believe after performing a contraction followed by a revision? We would like to extend the Harper Identity to cover this case.

In syntactic terms, the most straightforward suggestion would be to simply extend HI to cover not just one’s beliefs, but also one’s commitments to retain or lose various beliefs upon subsequent revisions:

$$(EHI) \quad [(\Psi \div A) * B] = [\Psi * B] \cap [(\Psi * \neg A) * B]$$

If $B \equiv \top$ then we obtain HI as a special case. Note that under weak assumptions, EHI can equivalently be restated in terms of contraction only:

Proposition 1 *EHI entails*

$$(EHIC) \quad [(\Psi \div A) \div B] = [\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B]$$

*and is equivalent to it in the presence of AGM*3 and the Levi Identity:*

$$(LI) \quad [\Psi * A] = Cn([\Psi \div \neg A] \cup \{A\}).$$

However, as Gärdenfors’ classic triviality result and its subsequent refinements [Gärdenfors, 1986; Rott, 1989; Etlin, 2009] have taught us, the unqualified extension of principles of belief dynamics to cover conditional beliefs is a risky business. And as it turns out, the above proposal is too strong: it can be shown that, under mild constraints on single shot revision and contraction, it places unacceptable restrictions on the space of permissible belief sets resulting from single revisions:

Proposition 2 *In the presence of AGM*5, AGM*6 and AGM \div 3, EHI (and more specifically, HI, alongside its left-to-right half $[(\Psi \div A) * B] \subseteq [\Psi * B] \cap [(\Psi * \neg A) * B]$) entails that there does not exist a belief state Ψ such that: (i) $[\Psi] = Cn(p \wedge q)$, (ii) $[\Psi * \neg p] = Cn(\neg p \wedge q)$ and (iii) $[\Psi * p \leftrightarrow \neg q] = Cn(p \leftrightarrow \neg q)$, where p and q are propositional atoms.²*

The above strategy and its shortcomings can equivalently be recast in semantic terms. Let us call a function \oplus that takes pairs of tpos as inputs and yields a tpo as an output a tpo combination operator, or a ‘combinator’. For convenience, we denote $\preceq_1 \oplus \preceq_2$ by ‘ $\preceq_{1 \oplus 2}$ ’.

In extending the Harper Identity to the iterated case, we are essentially looking for an appropriate combinator \oplus such that:

²The problem that we have just noted for EHI is closely related to the observation that an intersection of two sets of ‘rational doxastic conditionals’ need not itself be rational, which is familiar from the literature on default reasoning [Lehmann and Magidor, 1992].

$$\text{(COMBI)} \quad \preceq_{\Psi \div A} = \preceq_{\Psi} \oplus \preceq_{\Psi * \neg A}$$

Now, just as HI corresponds, given COMBI, to the following semantic principle:

$$\text{(\oplus HI)} \quad \min(\preceq_{1 \oplus 2}, W) = \min(\preceq_1, W) \cup \min(\preceq_2, W)$$

EHI amounts to

$$\text{(\oplus EHI)} \quad \text{For all } S \subseteq W, \min(\preceq_{1 \oplus 2}, S) = \min(\preceq_1, S) \cup \min(\preceq_2, S)$$

What our result above effectively demonstrates is that no combinator \oplus satisfies \oplus EHI unless we place undesirable restrictions on its domain: \oplus EHI is too much to ask for.

We will continue approaching our issue of interest from a predominantly semantic perspective for the remainder of the paper. In the following section, we retreat from \oplus EHI to offer an altogether weaker set of minimal postulates for \oplus , before taking a look at a concrete family of ‘Team Queuing’ combinators that satisfy them. We first establish a general characterisation of this family before showing that our set of minimal postulates suffices to characterise it in our restricted domain of interest.

4 Combinators: the bottom line

Since we are in the business of extending the Harper Identity, we will begin by requiring satisfaction of \oplus HI. We call combinators that satisfy this property ‘*basic*’ combinators.

In addition, even though EHI is too strong, certain weakenings of it do seem to be compelling. Specifically, it seems appropriate to require that our combination method leads to the following weak lower and upper bound principles:

$$\text{(LB)} \quad [\Psi * B] \cap [(\Psi * \neg A) * B] \subseteq [(\Psi \div A) * B]$$

$$\text{(UB)} \quad [(\Psi \div A) * B] \subseteq [\Psi * B] \cup [(\Psi * \neg A) * B]$$

We note that the former corresponds to the half of EHI that was *not* implicated in our earlier triviality result. Given COMBI, these will be ensured by requiring, respectively, the following upper and lower bounds on $\min(\preceq_{1 \oplus 2}, S)$ for any $S \subseteq W$ (note an upper, resp. lower bound on world-sets yields a lower, resp. upper bound on belief sets):

$$\text{(\oplus UB)} \quad \min(\preceq_{1 \oplus 2}, S) \subseteq \min(\preceq_1, S) \cup \min(\preceq_2, S)$$

$$\text{(\oplus LB)} \quad \text{Either } \min(\preceq_1, S) \subseteq \min(\preceq_{1 \oplus 2}, S) \text{ or } \min(\preceq_2, S) \subseteq \min(\preceq_{1 \oplus 2}, S)$$

\oplus UB and \oplus LB can be repackaged using only binary comparisons:

Proposition 3 \oplus UB and \oplus LB are respectively equivalent to the following:

$$\text{(\oplus SPU+)} \quad \text{If } x \prec_1 y \text{ and } z \prec_2 y \text{ then either } x \prec_{1 \oplus 2} y \text{ or } z \prec_{1 \oplus 2} y$$

$$\text{(\oplus WPU+)} \quad \text{If } x \preceq_1 y \text{ and } z \preceq_2 y \text{ then either } x \preceq_{1 \oplus 2} y \text{ or } z \preceq_{1 \oplus 2} y$$

$\oplus\text{SPU}+$ and $\oplus\text{WPU}+$ owe their names to their being respective strengthenings of the following principles of strict and weak preference unanimity, which are analogues of the ‘weak Pareto’ and ‘Pareto weak preference’ principles found in the social choice literature:

($\oplus\text{SPU}$) If $x \prec_1 y$ and $x \prec_2 y$ then $x \prec_{1\oplus 2} y$

($\oplus\text{WPU}$) If $x \preceq_1 y$ and $x \preceq_2 y$ then $x \preceq_{1\oplus 2} y$

We now consider a concrete family of basic combinators that satisfy both $\oplus\text{SPU}+$ and $\oplus\text{WPU}+$, and, indeed, can be shown to be characterised by precisely these principles in our domain of interest. We call these ‘TeamQueue’ combinators.

The basic idea behind this family—and motivation behind the name given to it—can be grasped by means of the following analogy: A number of couples go shopping for groceries. The supermarket that they frequent is equipped with two tills. For each till, we find a sequence of various groups of people queueing to pay for their items. In order to minimise the time spent in the store, each couple operates by “team queueing”: each member of the pair joins a group in a different queue and leaves their place to join their partner’s group in case this group arrives at the till first. After synchronously processing their first group of customers, the tills may or may not then operate at different and variable speeds. We consider the temporal sequence of sets of couples leaving the store. In our setting, the queues are the two tpos (with lower elements towards the head of the queue) and the couples are pairs of copies of each world.

More formally, we assume, for each ordered pair $\langle \preceq_1, \preceq_2 \rangle$ of tpos, a sequence $\langle a_{\preceq_1, \preceq_2}(i) \rangle_{i \in \mathbb{N}}$ such that:

(a1) $\emptyset \neq a_{\preceq_1, \preceq_2}(i) \subseteq \{1, 2\}$ for each i ,

(a2) $a_{\preceq_1, \preceq_2}(1) = \{1, 2\}$

$a_{\preceq_1, \preceq_2}(i)$ specifies which queue is to be processed at each step. Then (a1) ensures either one or both are processed, and (a2) says both are processed at the initial stage (which will ensure $\oplus\text{HI}$ holds for the resulting combinators). Then we construct the ordered partition $\langle T_1, T_2, \dots, T_m \rangle$ corresponding to $\preceq_{1\oplus 2}$ inductively as follows:

$$T_i = \bigcup_{j \in a_{\preceq_1, \preceq_2}(i)} \min\left(\bigcap_{k < i} T_k^c, \preceq_j\right)$$

(where ‘ T^c ’ denotes the complement of set T) and m is minimal such that $\bigcup_{i \leq m} T_i = W$. With this in hand, we can now offer:

Definition 1 \oplus is a TeamQueue combinator iff, for each ordered pair $\langle \preceq_1, \preceq_2 \rangle$ of tpos there exists a sequence $\langle a_{\preceq_1, \preceq_2}(i) \rangle_{i \in \mathbb{N}}$ satisfying (a1) and (a2) such that $\preceq_{1\oplus 2}$ is obtained as above.

It is easily verified that TeamQueue combinators are indeed basic combinators. The following example provides an elementary illustration of the combinator at work:

Example 1 Suppose that $W = \{w, x, y, z\}$, that \preceq_1 is the tpo represented by the ordered partition $\langle \{z\}, \{w\}, \{x, y\} \rangle$, and that \preceq_2 is represented by $\langle \{x, z\}, \{y\}, \{w\} \rangle$. Let \oplus be a TeamQueue combinator such that $\langle a_{\preceq_1, \preceq_2}(i) \rangle_{i \in \mathbb{N}} = \langle \{1, 2\}, \{2\}, \{1\}, \dots \rangle$. Then the ordered partition corresponding to $\preceq_{1 \oplus 2}$ is $\langle T_1, T_2, T_3 \rangle = \langle \{x, z\}, \{y\}, \{w\} \rangle$, since

$$\begin{aligned} T_1 &= \bigcup_{j \in \{1, 2\}} \min(W, \preceq_j) = \{x, z\} \\ T_2 &= \min(T_1^c, \preceq_2) = \{y\} \\ T_3 &= \min(T_1^c \cap T_2^c, \preceq_1) = \{w\} \end{aligned}$$

As noted above, TeamQueue combinators satisfy both $\oplus\text{SPU+}$ and $\oplus\text{WPU+}$. In fact, one can show that this family can actually be characterised by these two conditions, in the presence of a third:

Theorem 1 \oplus is a TeamQueue combinator iff it is a basic combinator that satisfies $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and the following ‘no overtaking’ property;

($\oplus\text{NO}$) For $i \neq j$, if $x \prec_i y$ and $z \preceq_j y$, then either $x \prec_{1 \oplus 2} y$ or $z \preceq_{1 \oplus 2} y$

Taken together, the three postulates $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and $\oplus\text{NO}$ say that in $\preceq_{1 \oplus 2}$, no world x is allowed to improve its position w.r.t. *both* input orderings \preceq_1 and \preceq_2 . Indeed each postulate blocks one of the three possible ways in which this ‘no double improvement’ condition could be violated. We note that this condition can be cashed out in terms of the following remarkable property:

Proposition 4 \oplus is a TeamQueue combinator iff it is a basic combinator that satisfies the following ‘trifurcation’ property, for all $S \subseteq W$:

($\oplus\text{TRI}$) $\min(\preceq_{1 \oplus 2}, S)$ is equal to either $\min(\preceq_1, S)$, $\min(\preceq_2, S)$ or $\min(\preceq_1, S) \cup \min(\preceq_2, S)$

Given COMBI, $\oplus\text{TRI}$ yields the claim that $[(\Psi \div A) * B]$ is equal to either $[\Psi * B]$, $[(\Psi * \neg A) * B]$ or $[\Psi * B] \cap [(\Psi * \neg A) * B]$.

To wrap up this section, it should be noted that the results so far have been perfectly domain-general, in the sense that they hold for combinators whose domain corresponded to the entire space of pairs of tpos defined over W . Our problem of interest is somewhat narrower in scope, however, since we are interested in the special case in which one of the tpos is obtained from the other by means of a revision. In particular, we assume the first two semantic postulates of [Darwiche and Pearl, 1997].

(CR*1) If $x, y \in \llbracket A \rrbracket$ then $x \preceq_{\Psi * A} y$ iff $x \preceq y$

(CR*2) If $x, y \in \llbracket \neg A \rrbracket$ then $x \preceq_{\Psi * A} y$ iff $x \preceq y$

In other words, \preceq_1 and \preceq_2 will always be $\llbracket A \rrbracket$ -variants for some sentence A , in the following sense:

Definition 2 Given $\preceq_1, \preceq_2 \in T(W)$ and $S \subseteq W$, we say \preceq_1 and \preceq_2 are S -variants iff $[x \preceq_1 y \text{ iff } x \preceq_2 y]$ holds for all $x, y \in (S \times S) \cup (S^c \times S^c)$. We let $V(W)$ denote the set of all $\langle \preceq_1, \preceq_2 \rangle \in T(W) \times T(W)$ such that \preceq_1, \preceq_2 are S -variants for some $S \subseteq W$.

Example 2 Suppose that $W = \{w, x, y, z\}$, that \preceq_1 is the tpo represented by the ordered partition $\langle \{w\}, \{x\}, \{y\}, \{z\} \rangle$, and that \preceq_2 is represented by $\langle \{w\}, \{x, y\}, \{z\} \rangle$. Then \preceq_1 and \preceq_2 are $\{y, z\}$ -variants, since (i) $w \prec_1 x$ and $w \prec_2 x$, as well as (ii) $y \prec_1 z$ and $y \prec_2 z$. They are not, however, $\{x, y\}$ -variants, since $x \prec_1 y$ but $y \not\prec_2 x$.

This leads to the following domain restriction on \oplus :

$$(\oplus\text{DOM}) \quad \text{Domain}(\oplus) \subseteq V(W)$$

As it turns out, this constraint allows for a noteworthy simplification of the characterisation of TeamQueue combinators:

Proposition 5 Given $\oplus\text{DOM}$, \oplus is a TeamQueue combinator iff it is a basic combinator that satisfies $\oplus\text{SPU+}$ and $\oplus\text{WPU+}$.

We also note, in passing, that

Proposition 6 Given $\oplus\text{DOM}$, \oplus satisfies $\oplus\text{SPU+}$ and $\oplus\text{WPU+}$ iff it satisfies $\oplus\text{SPU}$ and $\oplus\text{WPU}$, respectively.

Given Proposition 4, the potentially surprising upshot of Proposition 5 is that, in our domain of interest, satisfaction of $\oplus\text{LB}$ and $\oplus\text{UB}$ entails satisfaction of $\oplus\text{TRI}$.

5 Iterated Contraction via TeamQueue Combination

A central result of AGM theory says that, under assumption of HI, if $*$ satisfies the AGM revision postulates, then \div automatically satisfies the AGM contraction postulates. In this section we look at some of the postulates for both iterated revision and contraction that have been proposed in the literature. We show that, if $\preceq_{\Psi \div A}$ is defined from \preceq and $\preceq_{\Psi * \neg A}$ using COMBI via a TeamQueue combinator, then satisfaction of some well known sets of postulates for iterated revision leads to satisfaction of other well known sets of postulates for iterated contraction.

The most widely cited postulates for iterated revision are the four DP postulates of [Darwiche and Pearl, 1997]. These, like most of the postulates for iterated belief change, come in two flavours: a *semantic* one in terms of requirements on the tpo $\preceq_{\Psi * A}$ associated to the revised state $\Psi * A$, and a *syntactic* one in terms of requirements on the belief set $[(\Psi * A) * B]$ following a sequence of two revisions. Turning first to the semantic versions, we've already encountered the first two of these postulates—CR*1 and CR*2—in the previous section. The other two are

$$(\text{CR}^*3) \quad \text{If } x \in \llbracket A \rrbracket, y \in \llbracket \neg A \rrbracket \text{ and } x \prec y \text{ then } x \prec_{\Psi * A} y$$

$$(\text{CR}^*4) \quad \text{If } x \in \llbracket A \rrbracket, y \in \llbracket \neg A \rrbracket \text{ and } x \preceq y \text{ then } x \preceq_{\Psi * A} y$$

Each of these has an equivalent corresponding syntactic version as follows:

- (C*1) If $A \in \text{Cn}(B)$ then $[(\Psi * A) * B] = [\Psi * B]$
- (C*2) If $\neg A \in \text{Cn}(B)$ then $[(\Psi * A) * B] = [\Psi * B]$
- (C*3) If $A \in [\Psi * B]$ then $A \in [(\Psi * A) * B]$
- (C*4) If $\neg A \notin [\Psi * B]$ then $\neg A \notin [(\Psi * A) * B]$

Chopra et al [2008] proposed a list of ‘counterparts’ to the DP postulates for the case of $\Psi \div A$. The semantic versions of these were:

- (CR \div 1) If $x, y \in \llbracket \neg A \rrbracket$ then $x \preceq_{\Psi \div A} y$ iff $x \preceq y$
- (CR \div 2) If $x, y \in \llbracket A \rrbracket$ then $x \preceq_{\Psi \div A} y$ iff $x \preceq y$
- (CR \div 3) If $x \in \llbracket \neg A \rrbracket, y \in \llbracket A \rrbracket$ and $x \prec y$ then $x \prec_{\Psi \div A} y$
- (CR \div 4) If $x \in \llbracket \neg A \rrbracket, y \in \llbracket A \rrbracket$ and $x \preceq y$ then $x \preceq_{\Psi \div A} y$

Chopra et al [2008] showed (their Theorem 2) that, in the presence of the AGM postulates (reformulated as in our setting to apply to belief *states* rather than just belief *sets*) each of these postulates has an equivalent syntactic version as follows:

- (C \div 1) If $\neg A \in \text{Cn}(B)$ then $[(\Psi \div A) * B] = [\Psi * B]$
- (C \div 2) If $A \in \text{Cn}(B)$ then $[(\Psi \div A) * B] = [\Psi * B]$
- (C \div 3) If $\neg A \in [\Psi * B]$ then $\neg A \in [(\Psi \div A) * B]$
- (C \div 4) $A \notin [\Psi * B]$ then $A \notin [(\Psi \div A) * B]$

As it turns out, the definition of $\preceq_{\Psi \div A}$ from \preceq and $\preceq_{\Psi * \neg A}$ using COMBI via a TeamQueue combinator allows us to show the precise sense in which Chopra *et al*’s postulates are ‘Darwiche-Pearl-like’, as they put it:

Proposition 7 *Let \oplus be a TeamQueue combinator, let $*$ be an AGM revision operator and let \div be such that $\preceq_{\Psi \div A}$ is defined from $*$ via COMBI using \oplus . Then, for each $i = 1, 2, 3, 4$, if $*$ satisfies CR**i* then \div satisfies CR \div *i*.*

As a corollary, given the AGM postulates, we recover the same result for the syntactic versions as well.

Finally, Nayak *et al* [2007] have endorsed the following principle of ‘Principled Factored Intersection’, which they show to be satisfied by a number of proposals for iterated contraction:

- (PFI) Given $B \in [\Psi \div A]$
 - (a) If $\neg B \rightarrow \neg A \in [(\Psi \div A) \div B]$, then $[(\Psi \div A) \div B] = [\Psi \div A] \cap [\Psi \div \neg A \rightarrow B]$
 - (b) If $\neg B \rightarrow \neg A, \neg B \rightarrow A \notin [(\Psi \div A) \div B]$, then $[(\Psi \div A) \div B] = [\Psi \div A] \cap [\Psi \div \neg A \rightarrow B] \cap [\Psi \div A \rightarrow B]$
 - (c) If $\neg B \rightarrow A \in [(\Psi \div A) \div B]$, then $[(\Psi \div A) \div B] = [\Psi \div A] \cap$

$$[\Psi \div A \rightarrow B]$$

The rationale for PFI remains rather unclear to date. Indeed, the only justifications provided appear to be (a) that PFI avoids a particular difficulty faced by another constraint that has been proposed in the literature—namely Rott’s ‘Qualified Intersection’ principle [Rott, 2001]—and which can be reformulated in a manner that is superficially rather similar to PFI and (b) that PFI entails a pair of prima facie appealing principles. Neither of these considerations strike us as being particularly compelling. For one, Rott’s Qualified Intersection principle remains itself unclearly motivated. Secondly, plenty of ill-advised principles can be shown to have certain plausible consequences.

The TeamQueue approach, however, allows us to rest the principle on a far firmer foundation. Indeed:

Proposition 8 *Let \oplus be a TeamQueue combinator, let $*$ be an AGM revision operator and let \div be such that $\preceq_{\Psi \div A}$ is defined from $*$ via COMBI using \oplus . If $*$ satisfies CR*1 and CR*2 then \div satisfies PFI.*

6 The Synchronous TeamQueue Combinator

A special case of TeamQueue combinators takes $a_{\preceq_1, \preceq_2}(i) = \{1, 2\}$ for all ordered pairs $\langle \preceq_1, \preceq_2 \rangle$ and all i . This represents a particularly *fair* way of combining tpos. In terms of our supermarket analogy, it corresponds to the situation in which the tills process groups of customers at the same speed.

Definition 3 *The Synchronous TeamQueue (STQ) combinator is the TeamQueue combinator for which $a_{\preceq_1, \preceq_2}(i) = \{1, 2\}$ for all ordered pairs $\langle \preceq_1, \preceq_2 \rangle$ and all i . We will denote the STQ combinator by \oplus_{STQ} .*

Example 3 *Suppose $W = \{x, y, z, w\}$, that \preceq_1 is the tpo represented by the ordered partition $\langle \{z\}, \{w\}, \{x, y\} \rangle$ and \preceq_2 is represented by $\langle \{x, z\}, \{y\}, \{w\} \rangle$. Then the ordered partition corresponding to $\preceq_{1 \oplus_{\text{STQ}} 2}$ is $\langle T_1, T_2 \rangle = \langle \{x, z\}, \{w, y\} \rangle$.*

Roughly, $\preceq_{1 \oplus_{\text{STQ}} 2}$ tries to make each world as low in the ordering as possible, while trying to preserve the information contained in \preceq_1 and \preceq_2 . (The idea is similar to that of the *rational closure* construction in default reasoning [Lehmann and Magidor, 1992].) We remark that \oplus_{STQ} is commutative, i.e., $\preceq_{1 \oplus 2} = \preceq_{2 \oplus 1}$. It can be characterised semantically, in the absence of domain restrictions, as follows:

Theorem 2 *\oplus_{STQ} is the only basic combinator that satisfies both $\oplus\text{SPU+}$ and the following ‘Parity’ constraint:*

$$(\oplus\text{PAR}) \quad \text{If } x \prec_{1 \oplus 2} y \text{ then for each } i \in \{1, 2\} \text{ there exists } z \text{ s.t. } x \sim_{1 \oplus 2} z \text{ and } z \prec_i y$$

Note that $\oplus\text{WPU+}$ is not listed among the characteristic principles: it is entailed by the conjunction of $\oplus\text{SPU+}$ and $\oplus\text{PAR}$.

Whilst $\oplus\text{PAR}$ may not be immediately easy to grasp, it can be given a nice formulation in our setting in terms of the notion of *strong belief* [Battigalli and Siniscalchi, 2002; Stalnaker, 1996]. A sentence $A \in [\Psi]$ is strongly believed in Ψ in case the only way it can be dislodged by the next revision input B is if B is logically *inconsistent* with A . That is, A is strongly believed in Ψ iff (i) $A \in [\Psi]$, and (ii) $A \in [\Psi * B]$ for all sentences B such that $A \wedge B$ is consistent. Semantically, a consistent sentence A is strongly believed in Ψ iff every A -world is strictly more plausible than every $\neg A$ -world, i.e., $x \prec_{\Psi} y$ for every $x \in \llbracket A \rrbracket$, $y \in \llbracket \neg A \rrbracket$. With this in hand, one can show:

Proposition 9 $\oplus\text{PAR}$ is equivalent to:

$$(\oplus\text{SB}) \quad \text{If } x \prec_{1 \oplus 2} y \text{ for every } x \in S^c, y \in S, \text{ then } \min(\preceq_1, S) \cup \min(\preceq_2, S) \subseteq \min(\preceq_{1 \oplus 2}, S)$$

Given COMBI, $\oplus\text{SB}$ yields: If $\neg B$ is strongly believed in $\Psi \div A$ then $[(\Psi \div A) * B] \subseteq [\Psi * B] \cap [(\Psi * \neg A) * B]$. Thus, although we cannot have EHI for all A, B , the STQ combinator *does* guarantee it to hold for a certain restricted class of pairs of sentences, namely those A, B such that $\neg B$ is strongly believed after removing A .

To finish this section, we turn to further behaviour for iterated contraction that can be captured thanks to the further principles satisfied by $\oplus\text{STQ}$.

Three popular approaches to supplementing the AGM postulates for revision and the DP postulates can be found in the literature: the ‘natural’ [Boutilier, 1996], ‘restrained’ [Booth and Meyer, 2006], and ‘lexicographic’ [Nayak, 1994] approaches. All of these have the semantic consequence that the prior tpo \preceq_{Ψ} determines the posterior tpo $\preceq_{\Psi * A}$. All three promote the lowest A -worlds in \preceq_{Ψ} to become the lowest overall in $\preceq_{\Psi * A}$, but differ on what to do with the rest of the ordering. Natural revision leaves everything else unchanged, restrained revision preserves the strict ordering \prec_{Ψ} while additionally making every A -world x strictly lower than every $\neg A$ -world y for which $x \preceq_{\Psi} y$, and lexicographic revision just makes every A -world lower than every $\neg A$ -world, while preserving the ordering within each of $\llbracket A \rrbracket$ and $\llbracket \neg A \rrbracket$.

This raises an obvious question, namely: Which principles of iterated contraction does one recover from the natural, restrained and lexicographic revision operators, respectively, if one defines \div from $*$ using $\oplus\text{STQ}$? As it turns out, both the natural and the restrained revision operator yield the very same iterated contraction operator, which has been discussed in the literature under the name of ‘natural contraction’ [Nayak *et al.*, 2007], and which sets $\min(\preceq_{\Psi}, \llbracket \neg A \rrbracket) \cup \min(\preceq_{\Psi}, W)$ to be the lowest rank in $\preceq_{\Psi \div A}$ while leaving $\preceq_{\Psi \div A}$ otherwise unchanged from \preceq_{Ψ} .

Proposition 10 Let $*$ be any revision operator—such as the natural or restrained revision operator—satisfying the following property:

$$\text{If } x, y \notin \min(\preceq_{\Psi}, \llbracket A \rrbracket) \text{ and } x \prec_{\Psi} y, \text{ then } x \prec_{\Psi * A} y$$

Let \div be the contraction operator defined from $*$ via COMBI using \oplus_{STQ} . Then \div is the natural contraction operator.

We do not have a characterisation of the operator that is recovered from lexicographic revision in this manner, which we call the *STQ-lex contraction operator*. That is, STQ-lex contraction sets $\preceq_{\Psi \div A} = \preceq_{\Psi} \oplus_{\text{STQ}} \preceq_{\Psi *_{\text{L}} \neg A}$, where $*_{\text{L}}$ is lexicographic revision. We can report, however, that it is distinct from both lexicographic and priority contraction, the other two iterated contraction operators discussed in the literature alongside natural contraction [Nayak *et al.*, 2007]. Roughly, lexicographic contraction works by setting the i^{th} rank S_i of $\preceq_{\Psi \div A}$ to be the union of the i^{th} -lowest A -worlds with the i^{th} -lowest $\neg A$ -worlds.

Example 4 Suppose $W = \{x, y, z, w\}$ and \preceq_{Ψ} is the tpo represented by $\langle \{x\}, \{y\}, \{z\}, \{w\} \rangle$. Let $\llbracket A \rrbracket = \{x, w\}$, so that $\preceq_{\Psi *_{\text{L}} \neg A} = \langle \{y\}, \{z\}, \{x\}, \{w\} \rangle$. Then lexicographic contraction yields $\preceq_{\Psi \div A} = \langle \{x, y\}, \{z, w\} \rangle$ while STQ-lex contraction yields $\preceq_{\Psi \div A} = \langle \{x, y\}, \{z\}, \{w\} \rangle$.

Both lexicographic and priority contraction can, however, still be recovered via the TeamQueue approach. Lexicographic contraction can be recovered from lexicographic revision by combining, not \preceq_{Ψ} and $\preceq_{\Psi *_{\text{L}} \neg A}$, but rather $\preceq_{\Psi *_{\text{L}} A}$ and $\preceq_{\Psi *_{\text{L}} \neg A}$ using \oplus_{STQ} . Priority contraction can be recovered from lexicographic revision by combining \preceq_{Ψ} and $\preceq_{\Psi *_{\text{L}} \neg A}$ using a TeamQueue combinator. However, the combinator involved is not \oplus_{STQ} but rather the TeamQueue combinator that is most ‘biased’ towards \preceq_2 : the combinator for which, for all ordered pairs $\langle \preceq_1, \preceq_2 \rangle$, $a_{\preceq_1, \preceq_2}(1) = \{1, 2\}$, then $a_{\preceq_1, \preceq_2}(j) = \{2\}$ for all $j > 1$.

7 Conclusions

We have shown that the issue of extending the Harper identity to iterated belief change (a) is not a straightforward affair but (b) can be fruitfully approached by combining a pair of total preorders by means of TeamQueue combinator. We have also noted that one particular such combinator, the Synchronic TeamQueue combinator \oplus_{STQ} can be put to work to derive various counterparts for contraction of the three best known iterated revision operators.

Whilst the normative appeal of the characteristic syntactic properties \oplus_{LB} and \oplus_{UB} of the TeamQueue family of combinators is clear enough, we do not, at this stage, have a clear enough grasp of the normative appeal of the further syntactic requirement \oplus_{SB} that characterises \oplus_{STQ} . We plan to investigate this issue further in future work.

A second issue that we would like to explore is the question of whether or not it is possible to show that the Darwiche-Pearl postulates are *equivalent* to the ones proposed by Chopra *et al*, given a suitable further bridge principle taking us from iterated *contraction* to iterated *revision*. Such a task would first involve providing a compelling generalisation of the Levi Identity mentioned in Proposition 1 above.

Appendix

Proposition 1 *EHI entails*

$$(EHIC) \quad [(\Psi \div A) \div B] = [\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B]$$

and is equivalent to it in the presence of AGM*3 and the Levi Identity

$$(LI) \quad [\Psi * A] = Cn([\Psi \div \neg A] \cup \{A\})$$

Proof: From EHI to EHIC: By HI, which EHI entails, $[(\Psi \div A) \div B] = [\Psi \div A] \cap [(\Psi \div A) * \neg B] = [\Psi] \cap [\Psi * \neg A] \cap [(\Psi \div A) * \neg B]$. By EHI, we have $[(\Psi \div A) * \neg B] = [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B]$ and hence $[(\Psi \div A) \div B] = [\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B]$ as required.

From EHIC to EHI: By LI, we have $[(\Psi \div A) * \neg B] = Cn([\Psi \div A] \div B] \cup \{\neg B\})$. By EHIC, we have $[(\Psi \div A) \div B] = [\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B]$. So to recover EHI, we need to show that $Cn([\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B] \cup \{\neg B\}) = [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B]$. The left-to-right direction, i.e. $Cn([\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B] \cup \{\neg B\}) \subseteq [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B]$, is immediate. Regarding the right-to-left, assume, for some arbitrary C , that $C \in [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B]$. Firstly, it follows by AGM*3 and the deduction theorem that $\neg B \rightarrow C \in [\Psi]$ and $\neg B \rightarrow C \in [\Psi * \neg A]$. Secondly, it follows by deductive closure of belief sets that $\neg B \rightarrow C \in [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B]$. Therefore $\neg B \rightarrow C \in [\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B]$ and hence $C \in Cn([\Psi] \cap [\Psi * \neg B] \cap [\Psi * \neg A] \cap [(\Psi * \neg A) * \neg B] \cup \{\neg B\})$, as required.

Proposition 2 *In the presence of AGM*5, AGM*6 and AGM \div 3, EHI (and more specifically, HI, alongside its left-to-right half $[(\Psi \div A) * B] \subseteq [\Psi * B] \cap [(\Psi * \neg A) * B]$) entails that there does not exist a belief state Ψ such that: (i) $[\Psi] = Cn(p \wedge q)$, (ii) $[\Psi * \neg p] = Cn(\neg p \wedge q)$ and (iii) $[\Psi * p \leftrightarrow \neg q] = Cn(p \leftrightarrow \neg q)$, where p and q are propositional atoms.*

Proof: We first show that HI and the left-to-right half of EHI jointly entail that $[(\Psi \div A) \div B] \subseteq [\Psi * \neg B]$. Indeed, by HI, $[(\Psi \div A) \div B] = [\Psi \div A] \cap [(\Psi \div A) * \neg B] \subseteq [(\Psi \div A) * \neg B]$. By the left-to-right half of EHI, we then have $[(\Psi \div A) \div B] \subseteq [\Psi * \neg B] \cap [(\Psi * \neg A) * \neg B] \subseteq [\Psi * \neg B]$ as required.

We now establish that, in the presence of AGM*5, AGM*6 and AGM \div 3, HI and the left-to-right half of EHI jointly entail the following “vacuity” principle:

$$(VAC) \quad \text{If } A \text{ is consistent and } B \in [\Psi * A], \text{ then } [\Psi] \cap [\Psi * A] \subseteq [\Psi * B]$$

Indeed, assume that A is consistent and that $B \in [\Psi * A]$. Since A is consistent, so too is $[\Psi * A]$, by AGM*5, and hence $\neg B \notin [\Psi * A]$. Since, by HI, we have $[\Psi \div \neg A] = [\Psi] \cap [\Psi * A]$ (with help from AGM*6), it follows that $\neg B \notin [\Psi \div \neg A]$.

Given AGM \div 3, we then have $[(\Psi \div \neg A) \div \neg B] = [\Psi \div \neg A]$, and, by HI, $[(\Psi \div \neg A) \div \neg B] = [\Psi] \cap [\Psi * A]$. By the inclusion $[(\Psi \div \neg A) \div \neg B] \subseteq [\Psi * B]$, which we have shown above to be derivable from HI and the left-to-right half of EHI (plus AGM*6), it then follows that $[\Psi] \cap [\Psi * A] \subseteq [\Psi * B]$, as required.

With this in place, assume VAC and, for reductio, that there exists a belief set satisfying (i) to (iii). It follows from (ii) that $p \leftrightarrow \neg q \in [\Psi * \neg p]$. Given the latter, it then follows from VAC that $[\Psi] \cap [\Psi * \neg p] \subseteq [\Psi * p \leftrightarrow \neg q]$. But by (i) and (ii), $[\Psi] \cap [\Psi * \neg p] = \text{Cn}(p \wedge q) \cap \text{Cn}(\neg p \wedge q) = \text{Cn}(q)$. Hence, by $[\Psi] \cap [\Psi * \neg p] \subseteq [\Psi * p \leftrightarrow \neg q]$, we have $q \in [\Psi * p \leftrightarrow \neg q]$. But (iii) tells us that $[\Psi * p \leftrightarrow \neg q] = \text{Cn}(p \leftrightarrow \neg q)$. Contradiction.

Proposition 3 $\oplus\text{UB}$ and $\oplus\text{LB}$ are respectively equivalent to

$(\oplus\text{SPU}+)$ If $x \prec_1 y$ and $z \prec_2 y$ then $x \prec_{1\oplus 2} y$ or $z \prec_{1\oplus 2} y$

and

$(\oplus\text{WPU}+)$ If $x \preceq_1 y$ and $z \preceq_2 y$ then either $x \preceq_{1\oplus 2} y$ or $z \preceq_{1\oplus 2} y$

Proof: From $\oplus\text{UB}$ to $\oplus\text{SPU}+$: Suppose that $x \prec_1 y$ and $z \prec_2 y$. From the former, we know that $\min(\preceq_1, \{x, y, z\}) \subseteq \{x, z\}$ and from the latter we know that $\min(\preceq_2, \{x, y, z\}) \subseteq \{x, z\}$. Thus, by $\oplus\text{UB}$, $\min(\preceq_{1\oplus 2}, \{x, y, z\}) \subseteq \{x, z\}$. From this, it must be the case that $y \notin \min(\preceq_{1\oplus 2}, \{x, y, z\})$, so either $x \prec_{1\oplus 2} y$ or $z \prec_{1\oplus 2} y$, as required.

From $\oplus\text{SPU}+$ to $\oplus\text{UB}$: Assume for contradiction that there exists an x , such that $x \in \min(\preceq_{1\oplus 2}, S)$ but $x \notin \min(\preceq_1, S) \cup \min(\preceq_2, S)$. From the latter, there exist $y, z \in S$, such that $y \prec_1 x$ and $z \prec_2 x$. By $\oplus\text{SPU}+$, it then follows that either $y \prec_{1\oplus 2} x$ or $z \prec_{1\oplus 2} x$, contradicting $x \in \min(\preceq_{1\oplus 2}, S)$. Thus, $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_1, S) \cup \min(\preceq_2, S)$, as required.

From $\oplus\text{LB}$ to $\oplus\text{WPU}+$: We derive the contrapositive of $\oplus\text{WPU}+$, namely:

If $y \prec_{1\oplus 2} x$ and $y \prec_{1\oplus 2} z$, then $y \prec_1 x$ or $y \prec_2 z$

Assume then that $y \prec_{1\oplus 2} x$ and $y \prec_{1\oplus 2} z$. It follows from this that $\min(\preceq_{1\oplus 2}, \{x, y, z\}) \subseteq \{y\}$. By $\oplus\text{LB}$, we then recover either (i) $\min(\preceq_1, \{x, y, z\}) \subseteq \{y\}$ or (ii) $\min(\preceq_2, \{x, y, z\}) \subseteq \{y\}$. Assume (i). It follows that $y \prec_1 x$. Assume (ii). It follows that $y \prec_2 z$. Hence, either $y \prec_1 x$ or $y \prec_2 z$, as required.

From $\oplus\text{WPU}+$ to $\oplus\text{LB}$: Assume for reductio that $\oplus\text{LB}$ fails, so that there exist an x and a y such that $y \in \min(\preceq_1, S)$ and $z \in \min(\preceq_2, S)$, but $y, z \notin \min(\preceq_{1\oplus 2}, S)$. From the latter, there exist an x_1 and x_2 such that $x_1, x_2 \in S$, $x_1 \prec_{1\oplus 2} y$ and $x_2 \prec_{1\oplus 2} z$. Since $\preceq_{1\oplus 2}$ is a total preorder, we may assume that there exists an x such that $x \in S$, $x \prec_{1\oplus 2} y$ and $x \prec_{1\oplus 2} z$. By $\oplus\text{WPU}+$, we then have either $x \prec_1 y$ or $x \prec_2 z$, contradicting our assumption that $y \in \min(\preceq_1, S)$ and $z \in \min(\preceq_2, S)$.

Theorem 1 \oplus is a TeamQueue combinator iff it is a basic combinator that satisfies $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and the following ‘no overtaking’ property;

($\oplus\text{NO}$) If either (i) $x \preceq_1 y$ and $z \prec_2 y$ or (ii) $x \preceq_2 y$ and $z \prec_1 y$, then either $x \preceq_{1\oplus 2} y$ or $z \prec_{1\oplus 2} y$

Proof: We prove that \oplus satisfies $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and $\oplus\text{NO}$ iff it satisfies

($\oplus\text{TRI}$) $\min(\preceq_{1\oplus 2}, S)$ is equal to either $\min(\preceq_1, S)$, $\min(\preceq_2, S)$ or $\min(\preceq_1, S) \cup \min(\preceq_2, S)$

The desired result then follows from Proposition 4 below.

We first show that $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and $\oplus\text{NO}$ entail $\oplus\text{TRI}$.

We know that $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_1, S) \cup \min(\preceq_2, S)$ from $\oplus\text{SPU+}$. Indeed, assume that $y \in \min(\preceq_{1\oplus 2}, S)$ but, for reductio, that $y \notin \min(\preceq_1, S) \cup \min(\preceq_2, S)$. Then $\exists x, z \in S$ such that $x \prec_1 y$ and $z \prec_2 y$. Then, by $\oplus\text{SPU+}$, either $x \prec_{1\oplus 2} y$ or $z \prec_{1\oplus 2} y$. Either way, we get $y \notin \min(\preceq_{1\oplus 2}, S)$. Contradiction. Hence, $y \in \min(\preceq_1, S) \cup \min(\preceq_2, S)$, as required.

Now if the converse holds, i.e. $\min(\preceq_1, S) \cup \min(\preceq_2, S) \subseteq \min(\preceq_{1\oplus 2}, S)$, then we have $\min(\preceq_{1\oplus 2}, S) = \min(\preceq_1, S) \cup \min(\preceq_2, S)$ and we are done. So assume $\min(\preceq_1, S) \cup \min(\preceq_2, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$. Then either $\min(\preceq_1, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$ or $\min(\preceq_2, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$. Let’s assume $\min(\preceq_1, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$. We will show that this implies $\min(\preceq_{1\oplus 2}, S) = \min(\preceq_2, S)$, which will suffice. (If instead we assume $\min(\preceq_2, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$, then the same reasoning will show $\min(\preceq_{1\oplus 2}, S) = \min(\preceq_1, S)$, which also suffices.) Since $\min(\preceq_1, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$, let $x \in \min(\preceq_1, S)$ but $x \notin \min(\preceq_{1\oplus 2}, S)$.

We first derive $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_2, S)$. Let $y \in \min(\preceq_{1\oplus 2}, S)$ and assume for reductio that $y \notin \min(\preceq_2, S)$. Then $\exists z \in S$ such that $z \prec_2 y$. From $y \in \min(\preceq_{1\oplus 2}, S)$, we know that $y \preceq_{1\oplus 2} z$. From $x \in \min(\preceq_1, S)$, we also know that $x \preceq_1 y$. From $z \prec_2 y$, $y \preceq_{1\oplus 2} z$ and $x \preceq_1 y$, we can deduce by $\oplus\text{NO}$ that $x \preceq_{1\oplus 2} y$, in contradiction with $x \notin \min(\preceq_{1\oplus 2}, S)$. Hence, $y \in \min(\preceq_2, S)$, as required.

We now derive $\min(\preceq_2, S) \subseteq \min(\preceq_{1\oplus 2}, S)$. Let $y \in \min(\preceq_2, S)$ and assume, for reductio, that $y \notin \min(\preceq_{1\oplus 2}, S)$. From $x, y \notin \min(\preceq_{1\oplus 2}, S)$, $\exists z \in S$, such that $z \prec_{1\oplus 2} x$ and $z \prec_{1\oplus 2} y$. Then, from $\oplus\text{WPU+}$, we have either $z \prec_1 x$ or $z \prec_2 y$. If $z \prec_1 x$, then we contradict $x \in \min(\preceq_1, S)$. If $z \prec_2 y$, then we contradict $y \in \min(\preceq_2, S)$. Either way, we get a contradiction, so $y \in \min(\preceq_{1\oplus 2}, S)$, as required.

Finally, we show that $\oplus\text{TRI}$ entails $\oplus\text{SPU+}$, $\oplus\text{WPU+}$ and $\oplus\text{NO}$.

Regarding $\oplus\text{SPU+}$: From $\oplus\text{TRI}$, we know that, $\forall S$, $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_1, S) \cup \min(\preceq_2, S)$. Now suppose that $x \prec_1 y$ and $z \prec_2 y$. Then $y \notin \min(\preceq_1, \{x, y, z\}) \cup \min(\preceq_2, \{x, y, z\})$. Hence $y \notin \min(\preceq_{1\oplus 2}, \{x, y, z\})$, so $x \prec_{1\oplus 2} y$ or $z \prec_{1\oplus 2} y$, as required.

Regarding $\oplus\text{WPU}+$: From $\oplus\text{TRI}$, we know that, $\forall S$, either $\min(\preceq_1, S) \subseteq \min(\preceq_{1\oplus 2}, S)$ or $\min(\preceq_2, S) \subseteq \min(\preceq_{1\oplus 2}, S)$. This is the property $\oplus\text{LB}$ and we already proved in Proposition 3 that it entails $\oplus\text{WPU}+$.

Regarding $\oplus\text{NO}$: From $\oplus\text{TRI}$, we know that, $\forall S, i \neq j$, either $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_i, S)$ or $\min(\preceq_j, S) \subseteq \min(\preceq_{1\oplus 2}, S)$. Now assume $x \prec_i y, y \preceq_{1\oplus 2} x, z \preceq_j y$ and, for reductio, $y \prec_{1\oplus 2} z$. From $y \preceq_{1\oplus 2} x$ and $y \prec_{1\oplus 2} z$, we get $y \in \min(\preceq_{1\oplus 2}, \{x, y, z\})$ but from $x \prec_i y$, we get $y \notin \min(\preceq_i, \{x, y, z\})$. Hence $\min(\preceq_{1\oplus 2}, \{x, y, z\}) \not\subseteq \min(\preceq_i, \{x, y, z\})$. From this and the property cited at the beginning of this paragraph, we get $\min(\preceq_j, \{x, y, z\}) \subseteq \min(\preceq_{1\oplus 2}, \{x, y, z\})$. We also know from $\oplus\text{TRI}$ that $\min(\preceq_{1\oplus 2}, \{x, y, z\}) \subseteq \min(\preceq_1, \{x, y, z\}) \cup \min(\preceq_2, \{x, y, z\})$. Hence, since $y \in \min(\preceq_{1\oplus 2}, \{x, y, z\})$ and $y \notin \min(\preceq_i, \{x, y, z\})$, we get $y \in \min(\preceq_j, \{x, y, z\})$. Hence, since $z \preceq_j y, z \in \min(\preceq_j, \{x, y, z\})$ and so, from $\min(\preceq_j, \{x, y, z\}) \subseteq \min(\preceq_{1\oplus 2}, \{x, y, z\})$, $z \in \min(\preceq_{1\oplus 2}, \{x, y, z\})$, contradicting $y \prec_{1\oplus 2} z$. Hence $z \preceq_{1\oplus 2} y$, as required.

Proposition 4 \oplus is a TeamQueue combinator iff it is a basic combinator that satisfies the following ‘trifurcation’ property:

$$(\oplus\text{TRI}) \quad \min(\preceq_{1\oplus 2}, S) \text{ is equal to either } \min(\preceq_1, S), \min(\preceq_2, S) \text{ or } \min(\preceq_1, S) \cup \min(\preceq_2, S)$$

Proof: Right-to-left direction: Let \oplus be any combinator that satisfies those properties. We must specify a sequence a_{\preceq_1, \preceq_2} for each ordered pair $\langle \preceq_1, \preceq_2 \rangle$ such that (i) \oplus_a satisfies properties (a1) and (a2) and (ii) $\oplus_a = \oplus$.

Assume that $\langle S_1, S_2, \dots, S_n \rangle$ represents $\preceq_{1\oplus 2}$. Then we specify a_{\preceq_1, \preceq_2} by setting, for all i ,

$$j \in a_{\preceq_1, \preceq_2}(i) \text{ iff } \min\left(\bigcap_{k < i} S_k^c, \preceq_j\right) \subseteq S_i (= \min\left(\bigcap_{k < i} S_k^c, \preceq_{1\oplus 2}\right))$$

Regarding (i), \oplus_a satisfies (a1) since \oplus satisfies $\oplus\text{TRI}$ and (a2) since \oplus satisfies $\oplus\text{HI}$

Regarding (ii), let $\langle T_1, T_2, \dots, T_m \rangle$ represent $\preceq_{1\oplus a 2}$. We prove by induction that $T_i = S_i$. Regarding $i = 1$: The result follows from $\oplus\text{HI}$. Regarding the inductive step: Assume $T_j = S_j, \forall j < i$. We want to show $T_i = S_i$. By construction, $T_i = \bigcup_{j \in a(i)} \min(\preceq_j, \bigcap_{k < i} S_k^c)$. So we need to show $\min(\bigcap_{k < i} S_k^c, \preceq_{1\oplus 2}) = \bigcup_{j \in a(i)} \min(\preceq_j, \bigcap_{k < i} S_k^c)$. This follows from $\oplus\text{TRI}$.

Left-to-right direction: We show that \oplus_a satisfies each of $\oplus\text{SPU}+, \oplus\text{WPU}+$ and $\oplus\text{NO}$.

- Regarding $\oplus\text{SPU}+$: We prove the contrapositive. Suppose $y \preceq_{1\oplus 2} x$ and $y \preceq_{1\oplus 2} z$. Assume $y \in S_i = \bigcup_{j \in a(i)} \min(\preceq_j, \bigcap_{k < i} S_k^c) \subseteq \min(\preceq_1, \bigcap_{k < i} S_k^c) \cup \min(\preceq_2, \bigcap_{k < i} S_k^c)$. Assume $y \in \min(\preceq_1, \bigcap_{k < i} S_k^c)$. Since

$y \preceq_{1\oplus 2} x$, we know that $x \in \bigcap_{k < i} S_k^c$, hence $y \preceq_1 x$, as required. Similarly, if $y \in \min(\preceq_2, \bigcap_{k < i} S_k^c)$, then $y \preceq_2 z$.

- Regarding $\oplus\text{WPU}+$: We prove the contrapositive. Suppose $y \prec_{1\oplus 2} x$ and $y \prec_{1\oplus 2} z$. Assume $y \in S_i$. Since $y \prec_{1\oplus 2} x$ and $y \prec_{1\oplus 2} z$, we know that $x, z \in \bigcap_{k < i} S_k^c \cap S_i^c$. Now, we know that S_i equals one of $\min(\preceq_1, \bigcap_{k < i} S_k^c)$, $\min(\preceq_2, \bigcap_{k < i} S_k^c)$ or $\min(\preceq_1, \bigcap_{k < i} S_k^c) \cup \min(\preceq_2, \bigcap_{k < i} S_k^c)$. We consider each case in turn:

- (1) $S_i = \min(\preceq_1, \bigcap_{k < i} S_k^c)$: From $y \in S_i$ and $x \in \bigcap_{k < i} S_k^c \cap S_i^c$, we have $y \prec_1 x$, as required.
- (2) $S_i = \min(\preceq_2, \bigcap_{k < i} S_k^c)$: From $y \in S_i$ and $z \in \bigcap_{k < i} S_k^c \cap S_i^c$, we have $y \prec_2 z$, as required.
- (3) $S_i = \min(\preceq_1, \bigcap_{k < i} S_k^c) \cup \min(\preceq_2, \bigcap_{k < i} S_k^c)$: Either $y \in \min(\preceq_1, \bigcap_{k < i} S_k^c)$, in which case $y \prec_1 x$, or $y \in \min(\preceq_2, \bigcap_{k < i} S_k^c)$, in which case $y \prec_2 z$.

- Regarding $\oplus\text{NO}$: We show: If $x \prec_i y$, $y \preceq_{1\oplus 2} x$ and $z \preceq_j y$, then $z \preceq_{1\oplus 2} y$, $i \neq j$, $i, j \in \{1, 2\}$. Suppose that $x \prec_i y$, $y \preceq_{1\oplus 2} x$ and $z \preceq_j y$. We must show that $z \preceq_{1\oplus 2} y$. Assume $y \in S_t$. Then, from $y \preceq_{1\oplus 2} x$ and $z \preceq_j y$, we have $x, z \in \bigcap_{k < t} S_k^c$ and furthermore $z \in S_t^c$. We know that S_t equals one of $\min(\preceq_1, \bigcap_{k < t} S_k^c)$, $\min(\preceq_2, \bigcap_{k < t} S_k^c)$ or $\min(\preceq_1, \bigcap_{k < t} S_k^c) \cup \min(\preceq_2, \bigcap_{k < t} S_k^c)$. From $x \prec_i y$, we know that $y \notin \min(\preceq_i, \bigcap_{k < t} S_k^c)$, hence we must have $y \in \min(\preceq_j, \bigcap_{k < t} S_k^c)$. Furthermore, we are left with either $S_t = \min(\preceq_j, \bigcap_{k < t} S_k^c)$ or $S_t = \min(\preceq_1, \bigcap_{k < t} S_k^c) \cup \min(\preceq_2, \bigcap_{k < t} S_k^c)$. In either case, we must have $y \prec_j z$, as required.

Proposition 5 *Given $\oplus\text{DOM}$, \oplus is a TeamQueue combinator iff it satisfies $\oplus\text{SPU}+$ and $\oplus\text{WPU}+$.*

Proof: We show that, given $\oplus\text{DOM}$, if \oplus satisfies $\oplus\text{SPU}+$ and $\oplus\text{WPU}+$, then it satisfies $\oplus\text{NO}$ and hence, by Propositions 1 and 6, is a TeamQueue combinator.

Suppose $x \prec_i y$, $y \preceq_{1\oplus 2} x$ and $z \preceq_j y$, with $i \neq j$. We must show $z \preceq_{1\oplus 2} y$. If we can show $z \preceq_i y$, then we can conclude $z \preceq_{1\oplus 2} y$ from $\oplus\text{WPU}$. So suppose for reductio that $y \prec_i z$. From $\oplus\text{DOM}$, $\exists S$, such that, $\forall u, v \in S$, $u \preceq_1 v$ iff $u \preceq_2 v$ and $\forall u, v \in S^c$, $u \preceq_1 v$ iff $u \preceq_2 v$. From $z \preceq_j y$ and $y \prec_i z$, it must be the case that $y \in S$ and $z \in S^c$. If $x \in S$, then from $x \prec_i y$, we get $x \prec_j y$ and so $x \prec_{1\oplus 2} y$ from $\oplus\text{SPU}$, contradicting $y \preceq_{1\oplus 2} x$. If $x \in S^c$, then, since $x \prec_i y \prec_i z$ and $z \in S^c$, $x \prec_j z$. So from this and $z \preceq_j y$, we get $x \prec_j y$ and so again $x \prec_{1\oplus 2} y$ from $\oplus\text{SPU}$, contradicting $y \preceq_{1\oplus 2} x$. Hence, it must be that $z \preceq_i y$, as required.

Proposition 6 *Given $\oplus\text{DOM}$, \oplus satisfies $\oplus\text{SPU}+$ and $\oplus\text{WPU}+$ iff it satisfies $\oplus\text{SPU}$ and $\oplus\text{WPU}$, respectively.*

Proof: We prove this by demonstrating the equivalence, given $\oplus\text{DOM}$, of $\oplus\text{SPU}$ and $\oplus\text{WPU}$ with $\oplus\text{UB}$ and $\oplus\text{LB}$, respectively, which we have shown (see Proposition 3) to be equivalent to $\oplus\text{SPU}+$ and $\oplus\text{WPU}+$, respectively.

Regarding $\oplus\text{SPU}$ and $\oplus\text{UB}$, our proof is direct. Regarding $\oplus\text{WPU}$ and $\oplus\text{LB}$, we first show that $\oplus\text{WPU}$ is equivalent to the following weakening $\oplus\text{WLB}$ of $\oplus\text{LB}$:

$$(\oplus\text{WLB}) \quad \min(\preceq_1, S) \cap \min(\preceq_2, S) \subseteq \min(\preceq_{1\oplus 2}, S)$$

before showing that $\oplus\text{WLB}$ is equivalent to $\oplus\text{LB}$ under the domain restriction $\oplus\text{DOM}$.

From $\oplus\text{UB}$ to $\oplus\text{SPU}$: The result follows from the fact that $x \preceq y$ iff $\min(\preceq, \{x, y\}) \subseteq \{x\}$.

From $\oplus\text{SPU}$ to $\oplus\text{UB}$: It suffices to show that $\min(\preceq_{1\oplus 2}, S) \subseteq \min(\preceq_1, S) \cup \min(\preceq_2, S)$. Assume $\oplus\text{DOM}$, $\oplus\text{SPU}$ and that there exists an x , such that $x \in \min(\preceq_{1\oplus 2}, S)$ but, for contradiction, that $x \notin \min(\preceq_1, S) \cup \min(\preceq_2, S)$. From the latter, there exist $y_1, y_2 \in S$, such that (i) $y_1 \prec_1 x$ and (ii) $y_2 \prec_2 x$. From the former, (iii) $x \preceq_{1\oplus 2} y_1$ and (iv) $x \preceq_{1\oplus 2} y_2$. From (i) and (iii) on the one hand and (ii) and (iv) on the other, by $\oplus\text{SPU}$, we recover (v) $x \preceq_2 y_1$ and (vi) $x \preceq_1 y_2$, respectively. The conjunctions of (i) and (vi), i.e. $y_1 \prec_1 x \preceq_1 y_2$, and of (ii) and (v), i.e. $y_2 \prec_2 x \preceq_2 y_1$, however, jointly contradict $\oplus\text{DOM}$, since the latter entails that there exist no x, y_1, y_2 such that $y_1 \prec_1 x \preceq_1 y_2$ but $y_2 \prec_2 x \preceq_2 y_1$. Hence $x \in \min(\preceq_1, S) \cup \min(\preceq_2, S)$, as required.

From $\oplus\text{WPU}$ to $\oplus\text{WLB}$: Let $x \in \min(\preceq_1, S) \cap \min(\preceq_2, S)$ and assume for reductio that $x \notin \min(\preceq_{1\oplus 2}, S)$. Then there exists $y \in S$ such that $y \prec_{1\oplus 2} x$. By $\oplus\text{WPU}$, either $y \prec_1 x$ or $y \prec_2 x$. Assume $y \prec_1 x$ (the other case is analogous). Then $x \notin \min(\preceq_1, S)$ and hence $x \notin \min(\preceq_1, S) \cap \min(\preceq_2, S)$. Contradiction. Hence, $x \in \min(\preceq_{1\oplus 2}, S)$, as required.

From $\oplus\text{WLB}$ to $\oplus\text{WPU}$: Suppose $x \preceq_1 y$ and $x \preceq_2 y$. Then $x \in \min(\preceq_1, \{x, y\}) \cap \min(\preceq_2, \{x, y\})$. Assume for reductio that $y \prec_{1\oplus 2} x$. Then $x \notin \min(\preceq_{1\oplus 2}, \{x, y\})$, so, from $\oplus\text{WLB}$, $x \notin \min(\preceq_1, \{x, y\}) \cap \min(\preceq_2, \{x, y\})$. Contradiction. Hence $x \preceq_{1\oplus 2} y$, as required.

From $\oplus\text{LB}$ to $\oplus\text{WLB}$: Obvious.

From $\oplus\text{WLB}$ to $\oplus\text{LB}$: Assume that $\oplus\text{LB}$ doesn't hold. Then there exists an S such that $\min(\preceq_1, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$ and $\min(\preceq_2, S) \not\subseteq \min(\preceq_{1\oplus 2}, S)$. So there exist $x, y \in S$ such that $x \in \min(\preceq_1, S)$, $y \in \min(\preceq_2, S)$ and $x, y \notin \min(\preceq_{1\oplus 2}, S)$. Hence there exists $z \in S$ such that $z \prec_{1\oplus 2} x$ and $z \prec_{1\oplus 2} y$. By $\oplus\text{WLB}$, we know from $z \prec_{1\oplus 2} x$ that either $z \prec_1 x$ or $z \prec_2 x$. From this and the fact that $x \in \min(\preceq_1, S)$, we recover the result that $z \prec_2 x$. Similarly, we also recover $z \prec_1 y$. So we obtain the following pattern: $x \preceq_1 z \prec_1 y$ and $y \preceq_2 z \prec_2 x$. But this is not possible given $\oplus\text{DOM}$. Hence $\oplus\text{LB}$ holds, as required.

Proposition 7 Let \oplus be a TeamQueue combinator, let $*$ be an AGM revision operator and let \div be such that $\preceq_{\Psi \div A}$ is defined from $*$ using \oplus . Then, for each $i = 1, 2, 3, 4$, if $*$ satisfies (CR $*i$) then \div satisfies (CR $\div i$).

Proof: From CR $*1$ to CR $\div 1$: Let $x, y \in \llbracket \neg A \rrbracket$. We must show that $x \preceq_{\Psi \div A} y$ iff $x \preceq_{\Psi} y$. Note that from CR $*1$, we have (1) $x \preceq_{\Psi * \neg A} y$ iff $x \preceq_{\Psi} y$. Regarding the left-to-right direction of the equivalence: Assume (2) $y \prec x$. From (1) and (2), we recover (3) $y \prec_{\Psi * \neg A} x$. From (2) and (3), by \oplus SPU, it follows that $y \prec_{\Psi \div A} x$, as required. Regarding the right-to-left-direction: Assume (4) $x \preceq_{\Psi} y$. From (1) and (4), we recover (5) $x \preceq_{\Psi * \neg A} y$. From (4) and (5), by \oplus WPU, it follows that $x \preceq_{\Psi \div A} y$, as required.

From CR $*2$ to CR $\div 2$: Similar proof as the one given for the derivation of CR $\div 1$ from CR $*1$.

From CR $*3$ to CR $\div 3$: Let $x \in \llbracket \neg A \rrbracket, y \in \llbracket A \rrbracket$ and (1) $x \prec_{\Psi} y$. We must show that $x \prec_{\Psi \div A} y$. From CR $*3$, we recover (2) $x \prec_{\Psi * \neg A} y$. From (1) and (2), by \oplus SPU, we then obtain $x \prec_{\Psi \div A} y$, as required.

From CR $*4$ to CR $\div 4$: Let $x \in \llbracket \neg A \rrbracket, y \in \llbracket A \rrbracket$ and (1) $x \preceq_{\Psi} y$. We must show that $x \preceq_{\Psi \div A} y$. From CR $*4$, we recover (2) $x \preceq_{\Psi * \neg A} y$. From (1) and (2), by \oplus WPU, we then obtain $x \preceq_{\Psi \div A} y$, as required.

Proposition 8 Let $*$ be any revision operator satisfying C $*1$ and C $*2$ and \div be the contraction operator defined from $*$ using any tpo aggregation function satisfying \oplus WPU, \oplus SPU and \oplus HI. Then \div satisfies PFI.

Proof: Assume that $*$ satisfies CR $*1$ and CR $*2$ and let \div be the contraction operator defined from $*$ using some tpo aggregation function satisfying \oplus WPU, \oplus SPU and \oplus HI. We saw above, in Proposition 7 that \div will also satisfy CR $\div 1$ and CR $\div 2$. The desired result then immediately follows from the theorem established by Ramachandran *et al* (2011, Theorem 1), according to which every contraction function \div obtained from a revision function $*$, such that \div and $*$ satisfy HI, satisfies PFI if it also satisfies CR $\div 1$ and CR $\div 2$.

Theorem 2 \oplus_{STQ} is the only basic combinator that satisfies both \oplus SPU+ and the following ‘Parity’ constraint:

(\oplus PAR) If $x \prec_{1 \oplus 2} y$ then for each $i \in \{1, 2\}$ there exists z s.t. $x \sim_{1 \oplus 2} z$ and $z \prec_i y$

Proof: We need to show that if \oplus satisfies \oplus SPU+ and \oplus PAR, for any \preceq_1, \preceq_2 , we have $\preceq_{1 \oplus 2} = \preceq_{1 \oplus_{\text{STQ}} 2}$. Assume that $\preceq_{1 \oplus 2} = \{S_1, S_2, \dots, S_m\}$ and $\preceq_{1 \oplus_{\text{STQ}} 2} = \{T_1, T_2, \dots, T_n\}$, where S_i, T_i are the ranks of the relevant tpos, with lower ranks being the most preferred.

We will prove, by induction on i , that $S_i = T_i, \forall i$. Assume $S_j = T_j, \forall j < i$. We must show $S_i = T_i$.

Regarding $S_i \subseteq T_i$: Let $x \in S_i$, so that $x \preceq_{1 \oplus 2} y, \forall y \in \bigcap_{j < i} S_j^c$. Assume for reductio that $x \notin T_i$. Since $x \in S_i$, we know that $x \in \bigcap_{j < i} S_j^c = \bigcap_{j < i} T_j^c$. Hence, since $x \notin T_i$ and, by construction of $\preceq_{1 \oplus \text{STQ}2}$, there exists $y_1 \in \bigcap_{j < i} T_j^c$ such that $y_1 \prec_1 x$ and there exists $y_2 \in \bigcap_{j < i} T_j^c$ such that $y_2 \prec_2 x$. Then, by $\oplus \text{SPU}+$, either $y_1 \prec_{1 \oplus \text{STQ}2} x$ or $y_2 \prec_{1 \oplus \text{STQ}2} x$, in both cases contradicting $x \preceq_{1 \oplus 2} y, \forall y \in \bigcap_{j < i} S_j^c$. Hence $x \in T_i$, as required.

Regarding $T_i \subseteq S_i$: Let $x \in T_i$. Then, by construction of $\preceq_{1 \oplus \text{STQ}2}$, we have $x \in \min(\preceq_1, \bigcap_{j < i} T_j^c) \cup \min(\preceq_2, \bigcap_{j < i} T_j^c)$. Assume for reductio that $x \notin S_i$. We know that $x \in \bigcap_{j < i} T_j^c$, so by the inductive hypothesis, $x \in \bigcap_{j < i} S_j^c$. From this and $x \notin S_i$ we know that there exists a $y \in S_i$, such that $y \prec_{1 \oplus 2} x$. Then from $\oplus \text{PAR}$, there exist a $z_1 \in S_i$ such that $z_1 \prec_1 x$ and a $z_2 \in S_i$ such that $z_2 \prec_2 x$. But this contradicts $x \in \min(\preceq_1, \bigcap_{j < i} T_j^c) \cup \min(\preceq_2, \bigcap_{j < i} T_j^c)$. Hence $x \in S_i$, as required.

Proposition 9 $\oplus \text{PAR}$ is equivalent to:

($\oplus \text{SB}$) If $x \prec_{1 \oplus 2} y$ for every $x \in S^c, y \in S$, then $\min(\preceq_1, S) \cup \min(\preceq_2, S) \subseteq \min(\preceq_{1 \oplus 2}, S)$

Proof: From $\oplus \text{PAR}$ to $\oplus \text{SB}$: Assume that $x \prec_{1 \oplus 2} y$ for every $x \in S^c, y \in S$. It suffices to show that $\min(\preceq_1, S) \cup \min(\preceq_2, S) \subseteq \min(\preceq_{1 \oplus 2}, S)$. So assume $x \in \min(\preceq_1, S) \cup \min(\preceq_2, S)$ but, for contradiction, $x \notin \min(\preceq_{1 \oplus 2}, S)$. Then $y \prec_{1 \oplus 2} x$ for some $y \in S$. From the latter, by $\oplus \text{PAR}$, we know that $z_1 \prec_1 x$ for some z_1 such that $y \preceq_{1 \oplus 2} z_1$ and $z_2 \prec_2 x$ for some z_2 such that $y \preceq_{1 \oplus 2} z_2$. Given our initial assumption, we can deduce from $y \preceq_{1 \oplus 2} z_1, y \preceq_{1 \oplus 2} z_2$ and $y \in S$ that $z_1, z_2 \in S$. But this, together with $z_1 \prec_1 x$ and $z_2 \prec_2 x$ contradicts $x \in \min(\preceq_1, S) \cup \min(\preceq_2, S)$. Hence $x \in \min(\preceq_{1 \oplus 2}, S)$, as required.

From $\oplus \text{SB}$ to $\oplus \text{PAR}$: Suppose $\oplus \text{PAR}$ does not hold, i.e. $\exists x, y$, such that $x \prec_{1 \oplus 2} y$ and for no z do we have $x \sim_{1 \oplus 2} z$ and $z \prec_1 y$ (similar reasoning will apply if we replace \prec_1 by \prec_2 here). We will show that $\oplus \text{SB}$ fails, i.e. that $\exists S \subseteq W$, such that $x \prec_{\Psi_{1 \oplus 2}} y$ for every $x \in S^c, y \in S$ and $\min(\preceq_1, S) \cup \min(\preceq_2, S) \not\subseteq \min(\preceq_{1 \oplus 2}, S)$.

Let $S = \{w \mid x \preceq_{1 \oplus 2} w\}$ (so that $S^c = \{w \mid w \prec_{1 \oplus 2} x\}$). Clearly $x \in S$ and, from $x \prec_{1 \oplus 2} y$, we know that $y \in S$ but $y \notin \min(\preceq_{1 \oplus 2}, S)$. Hence, to show $\min(\preceq_1, S) \cup \min(\preceq_2, S) \not\subseteq \min(\preceq_{1 \oplus 2}, S)$ and therefore that $\oplus \text{SB}$ fails, it suffices to show $y \in \min(\preceq_1, S)$. But if $y \notin \min(\preceq_1, S)$, then $z \prec_1 y$ for some $z \in S$, i.e. some z , such that $x \preceq_{1 \oplus 2} z$. Since $\preceq_{1 \oplus 2}$ is a tpo we may assume $x \sim_{1 \oplus 2} z$. This contradicts our initial assumption that for no z do we have $x \sim_{1 \oplus 2} z$ and $z \prec_1 y$. Hence $y \in \min(\preceq_1, S)$, as required.

Proposition 10 Let $*$ be any revision operator—such as the natural or restrained revision operator—satisfying the following property:

If $x, y \notin \min(\preceq_\Psi, \llbracket A \rrbracket)$ and $x \prec_\Psi y$, then $x \prec_{\Psi * A} y$

Let \div be the contraction operator defined from $*$ using \oplus_{STQ} . Then \div is the natural contraction operator.

Proof: Recall the definition of natural contraction:

- (\div -NAT) $x \preceq_{\Psi \div A} y$ iff
- (a) $x \in \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$, or
 - (b) $x, y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$ and $x \preceq_\Psi y$

We must show that for any $x, y \in W$ and $A \in L$, $x \preceq_{\Psi \div A} y$ iff $x \preceq_{\Psi \div NA} y$. We split into two cases.

Case 1: $x \in \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$. Then, by the definitions of \div_N and \div , we have both $x \preceq_{\Psi \div A} y$ and $x \preceq_{\Psi \div NA} y$, so the desired result holds.

Case 2: $x \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$. Then by definition of \div_N , $x \preceq_{\Psi \div NA} y$ iff both $y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$ and $x \preceq_\Psi y$. We now consider each direction of the equivalence to be demonstrated separately.

- From $x \preceq_{\Psi \div NA} y$ to $x \preceq_{\Psi \div A} y$: Suppose $x \preceq_{\Psi \div NA} y$, and hence that both $y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$ and $x \preceq_\Psi y$. Assume for reductio that $y \prec_{\Psi \div A} x$. By $\oplus\text{PAR}$: if $y \prec_{\Psi \div A} x$, then there exists z such that $z \sim_{\Psi \div A} y$ and $z \prec_\Psi x$. Hence there exists z such that $z \sim_{\Psi \div A} y$ and $z \prec_\Psi x$. Since $x \preceq_\Psi y$, we therefore also have $z \prec_\Psi y$. If $z \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket)$, then from the postulate mentioned in the proposition, we get $z \prec_{\Psi * \neg A} y$ and then $z \prec_{\Psi \div A} y$. Contradiction. Hence we can assume $z \in \min(\preceq_\Psi, \llbracket \neg A \rrbracket)$. From $x \preceq_\Psi y$, $y \prec_{\Psi \div A} x$ and $\oplus\text{WPU}$, we know that $y \prec_{\Psi * \neg A} x$. From this, $\text{CR} * 2$, $\text{CR} * 4$ and $x \preceq_\Psi y$, we get $y \in \llbracket \neg A \rrbracket$. Hence, from $z \prec_\Psi y$ and $\text{CR} * 1$, we recover $z \prec_{\Psi * \neg A} y$ and then $z \prec_{\Psi \div A} y$ by $\oplus\text{SPU}$. Contradiction again. Hence $x \preceq_{\Psi \div A} y$, as required.
- From $x \preceq_{\Psi \div A} y$ to $x \preceq_{\Psi \div NA} y$: Assume that $x \preceq_{\Psi \div A} y$ and, for reductio, that either $y \prec_\Psi x$ or $y \in \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$. If the latter holds, then we know that $y \in \min(\preceq_{\Psi \div A}, W)$, by definition of \div . Hence, from this and $x \preceq_{\Psi \div A} y$, we also deduce that $x \in \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$, contradicting the assumption that $x \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$. So assume that $y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$ and $y \prec_\Psi x$. From the latter and our assumption that $x \preceq_{\Psi \div A} y$, it follows by $\oplus\text{SPU}$ that $x \preceq_{\Psi \neg A} y$. But it also follows from $y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket) \cup \min(\preceq_\Psi, W)$ and $y \prec_\Psi x$ that $x, y \notin \min(\preceq_\Psi, \llbracket \neg A \rrbracket)$. We then recover, from the property mentioned in the proposition, the result that $x \preceq_\Psi y$, contradicting our assumption that $y \prec_\Psi x$. Hence, $x \preceq_{\Psi \div NA} y$, as required.

References

- [Alchourrón *et al.*, 1985] Carlos E Alchourrón, Peter Gärdenfors, and David Makinson. On the logic of theory change: Partial meet contraction and revision functions. *Journal of Symbolic Logic*, 50(02):510–530, 1985.
- [Battigalli and Siniscalchi, 2002] Pierpaolo Battigalli and Marciano Siniscalchi. Strong belief and forward induction reasoning. *Journal of Economic Theory*, 106(2):356–391, 2002.
- [Booth and Meyer, 2006] Richard Booth and Thomas Meyer. Admissible and restrained revision. *Journal of Artificial Intelligence Research*, 26(1):127–151, 2006.
- [Booth and Meyer, 2011] Richard Booth and Thomas Meyer. How to revise a total preorder. *Journal of Philosophical Logic*, pages 1–46, 2011.
- [Boutilier, 1996] Craig Boutilier. Iterated revision and minimal change of conditional beliefs. *Journal of Philosophical Logic*, 25(3):263–305, 1996.
- [Chopra *et al.*, 2008] Samir Chopra, Aditya Ghose, Thomas Meyer, and Ka-Shu Wong. Iterated belief change and the recovery axiom. *Journal of Philosophical Logic*, 37(5):501–520, 2008.
- [Darwiche and Pearl, 1997] Adnan Darwiche and Judea Pearl. On the logic of iterated belief revision. *Artificial Intelligence*, 89(1):1–29, 1997.
- [Etlin, 2009] David Etlin. The problem of noncounterfactual conditionals. *Philosophy of Science*, 76(5):676–688, 2009.
- [Gärdenfors, 1986] Peter Gärdenfors. Belief revisions and the ramsey test for conditionals. *The Philosophical Review*, pages 81–93, 1986.
- [Grove, 1988] Adam Grove. Two modellings for theory change. *Journal of Philosophical Logic*, 17(2):157–170, 1988.
- [Hansson, 2012] Sven Ove Hansson. Global and iterated contraction and revision: An exploration of uniform and semi-uniform approaches. *Journal of philosophical logic*, 41(1):143–172, 2012.
- [Harper, 1976] William L Harper. Rational conceptual change. In *PSA: Proceedings of the Biennial Meeting of the Philosophy of Science Association*, pages 462–494. JSTOR, 1976.
- [Hild and Spohn, 2008] Matthias Hild and Wolfgang Spohn. The measurement of ranks and the laws of iterated contraction. *Artificial Intelligence*, 172(10):1195–1218, 2008.
- [Jin and Thielscher, 2007] Yi Jin and Michael Thielscher. Iterated belief revision, revised. *Artificial Intelligence*, 171(1):1–18, 2007.

- [Katsuno and Mendelzon, 1991] Hirofumi Katsuno and Alberto O Mendelzon. Propositional knowledge base revision and minimal change. *Artificial Intelligence*, 52(3):263–294, 1991.
- [Lehmann and Magidor, 1992] Daniel Lehmann and Menachem Magidor. What does a conditional knowledge base entail? *Artificial intelligence*, 55(1):1–60, 1992.
- [Nayak *et al.*, 2003] Abhaya C Nayak, Maurice Pagnucco, and Pavlos Peppas. Dynamic belief revision operators. *Artificial Intelligence*, 146(2):193–228, 2003.
- [Nayak *et al.*, 2006] Abhaya Nayak, Randy Goebel, Mehmet Orgun, and Tam Pham. Taking levi identity seriously: A plea for iterated belief contraction. In *Knowledge Science, Engineering and Management*, pages 305–317. Springer, 2006.
- [Nayak *et al.*, 2007] Abhaya C Nayak, Randy Goebel, and Mehmet A Orgun. Iterated belief contraction from first principles. In *IJCAI*, pages 2568–2573, 2007.
- [Nayak, 1994] Abhaya C Nayak. Iterated belief change based on epistemic entrenchment. *Erkenntnis*, 41(3):353–390, 1994.
- [Peppas, 2014] Pavlos Peppas. A panorama of iterated revision. In *David Makinson on Classical Methods for Non-Classical Problems*, pages 71–94. Springer, 2014.
- [Ramachandran *et al.*, 2012] Raghav Ramachandran, Abhaya C Nayak, and Mehmet A Orgun. Three approaches to iterated belief contraction. *Journal of Philosophical Logic*, 41(1):115–142, 2012.
- [Rott, 1989] Hans Rott. Conditionals and theory change: Revisions, expansions, and additions. *Synthese*, 81(1):91–113, 1989.
- [Rott, 2001] Hans Rott. *Change, Choice and Inference: A Study of Belief Revision and Nonmonotonic Reasoning*. Oxford University Press, 2001.
- [Rott, 2009] Hans Rott. Shifting priorities: Simple representations for twenty-seven iterated theory change operators. In *Towards Mathematical Philosophy*, pages 269–296. Springer, 2009.
- [Stalnaker, 1996] Robert Stalnaker. Knowledge, belief and counterfactual reasoning in games. *Economics and philosophy*, 12(02):133–163, 1996.