

Article

Fluctuation, Dissipation and the Arrow of Time

Michele Campisi* and Peter Hänggi

Institute of Physics, University of Augsburg, Universitätsstr. 1, D-86135 Augsburg, Germany

* Author to whom correspondence should be addressed; michele.campisi@physik.uni-augsburg.de

Version November 8, 2011 submitted to *Entropy*. Typeset by *TEX* using class file *mdpi.cls*

Abstract: The recent development of the theory of fluctuation relations has led to new insights into the ever-lasting question of how irreversible behavior emerges from time-reversal symmetric microscopic dynamics. We provide an introduction to fluctuation relations, examine their relation to dissipation and discuss their impact on the arrow of time question.

Keywords: work, entropy, second law, minus first law

Classification: PACS 1.70.+w 05.40.-a 05.70.Ln

1. Introduction

Irreversibility enters the laws of thermodynamics in two distinct ways:


Equilibrium Principle An isolated, macroscopic system which is placed in an arbitrary initial state within a finite fixed volume will attain a unique state of equilibrium.

Second Law (Clausius) For a non-quasi-static process occurring in a thermally isolated system, the entropy change between two equilibrium states is non-negative.

The first of these two principles is the *Equilibrium Principle* [1], whereas the second is the *Second Law of Thermodynamics* in the formulation given by Clausius [3]. Very often the Equilibrium Principle is loosely referred to as the Second Law of Thermodynamics, thus creating a great confusion in the literature. So much that proposing to raise the Equilibrium Principle to the rank of one of the fundamental laws of thermodynamic became necessary [1]. Indeed it was argued that this Law of Thermodynamics,

18 defining the very concept of state of equilibrium, is the most fundamental of all the Laws of Thermody-
 19 namics (which in fact are formulated in terms of equilibrium states) and for this reason the nomenclature
 20 *Minus-First Law of Thermodynamics* was proposed for it.

Figure 1. Autonomous vs. nonautonomous dynamics. Top: Autonomous evolution of a gas from a non-equilibrium state to an equilibrium state (Minus-First Law). Bottom: Nonautonomous evolution of a thermally isolated gas between two equilibrium states. The piston moves according to a pre-determined protocol specifying its position λ_t in time. The entropy change is non-negative (Second Law).



21 The Minus-First Law of Thermodynamics and the Second Law of Thermodynamics consider two very
 22 different situations, see Fig. 1. The Minus-First Law deals with a completely isolated system that begins
 23 in non-equilibrium and ends in equilibrium, following its spontaneous and *autonomous* evolution. In the
 24 Second Law one considers a *thermally* (but not mechanically) isolated system that begins in equilibrium.
 25 A mechanical perturbation drives the system out of equilibrium, the perturbation is then turned off and
 26 a final equilibrium will be reached, corresponding to higher entropy.¹ At variance with the Minus-First
 27 Law, here the system does not evolve autonomously, but rather in response to a driving: we speak in this
 28 case of *nonautonomous* evolution.

29 Both the Minus-First Law and the Second Law have to do with irreversibility and the arrow of time.
 30 While since the seminal works of Boltzmann, the main efforts of those working in the foundations of
 31 statistical mechanics were directed to reconcile the Minus-First Law with the time-reversal symmetric
 32 microscopic dynamics, recent developments in the theory of fluctuation relations, have brought new and
 33 deep insights into the microscopic foundations of the Second Law. As we shall see below, fluctuation
 34 theorems highlight in a most clear way the fascinating fact that the Second Law is deeply rooted in the
 35 time-reversal symmetric nature of the microscopic laws of microscopic dynamics [4,5].

¹That such final equilibrium state exists is dictated by the Minus-First Law. Here we see clearly the reason for assigning a higher rank to the Equilibrium Principle

36 This connection is best seen if one considers the Second Law in the formulation given by Kelvin,
37 which is equivalent to Clausius formulation [6]:

38 **Second Law (Kelvin)** No work can be extracted from a closed equilibrium system during a cyclic vari-
39 ation of a parameter by an external source.

40 The field of fluctuation theorems has recently gained much attention. Many fluctuation theorems have
41 been reported in the literature, referring to different scenarios. Fluctuation theorems exist for classical
42 dynamics, stochastic dynamics, and for quantum dynamics; for transiently driven systems, as well as
43 for non equilibrium steady states; for systems prepared in canonical, micro-canonical, grand-canonical
44 ensembles, and even for systems initially in contact with “finite heat baths” [7]; they can refer to dif-
45 ferent quantities like work (different kinds), entropy production, exchanged heat, exchanged charge,
46 and even information, depending on different set-ups. All these developments including discussions of
47 the experimental applications of fluctuation theorems, have been summarized in a number of reviews
48 [4,5,8,9].

49 In Sec. 2 we will give a brief introduction to the classical work Fluctuation Theorem of Bochkov
50 and Kuzovlev [10], which is the first fluctuation theorem reported in the literature. The discussion of
51 this theorem suffices for our purpose of highlighting the impact of fluctuation theory on dissipation (Sec.
52 3) and on the arrow of time issue (Sec. 4). Remarks of the origin of time’s arrow in this context are
53 collected in Sec. (5)

54 2. The fluctuation theorem

55 2.1. Autonomous dynamics

56 Consider a completely isolated mechanical system composed of f degrees of freedom. Its dynamics
57 are dictated by some Hamiltonian $H(\mathbf{q}, \mathbf{p})$, which we assume to be time reversal symmetric; i.e.,

$$H(\mathbf{q}, \mathbf{p}) = H(\mathbf{q}, -\mathbf{p}) \quad (1)$$

58 Here $(\mathbf{q}, \mathbf{p}) = (q_1 \dots q_f, p_1 \dots p_f)$ denotes the conjugate pairs of coordinates and momenta describing
59 the microscopic state of the system.

60 The assumption of time-reversal symmetry implies that if $[\mathbf{q}(t), \mathbf{p}(t)]$ is a solution of Hamilton equa-
61 tions of motion, then, for any τ , $[\mathbf{q}(\tau - t), -\mathbf{p}(\tau - t)]$ is also a solution of Hamilton equations of motion.
62 This is the well known principle of *microreversibility* for *autonomous* systems [11].

63 We assume that the system is at equilibrium described by the Gibbs ensemble:

$$\varrho(\mathbf{q}, \mathbf{p}) = e^{-\beta H(\mathbf{q}, \mathbf{p})} / Z(\beta) \quad (2)$$

64 where $Z(\beta) = \int d\mathbf{p}d\mathbf{q} e^{-\beta H(\mathbf{q}, \mathbf{p})}$ is the canonical partition function, and $\beta^{-1} = k_B T$, with k_B being the
65 Boltzmann constant and T denotes the temperature.

66 We next imagine to be able to observe the time evolution of all coordinates and momenta in some time
67 span $t \in [0, \tau]$. Fluctuation theorems are concerned with the probability² $P[\Gamma]$ that the trajectory Γ is

²To be more precise, the probability density functional (PDFL)

68 observed. We will reserve the symbol Γ to denote the whole trajectory (that is, mathematically speaking
 69 to denote a map from the interval $[0, \tau]$ to the $2f$ dimensional phase space), whereas the symbol Γ_t will
 70 be used to denote the specific point in phase space visited by the trajectory Γ at time t . The central
 71 question is how the probability $P[\Gamma]$ compares with the probability $P[\tilde{\Gamma}]$ to observe $\tilde{\Gamma}$, the time-reversal
 72 companion of Γ : $\tilde{\Gamma}_t = \varepsilon\Gamma_{\tau-t}$ where $\varepsilon(\mathbf{q}, \mathbf{p}) = (\mathbf{q}, -\mathbf{p})$ denotes the time reversal operator. The answer is
 73 given by the microreversibility principle which implies:

$$P[\Gamma] = P[\tilde{\Gamma}]. \quad (3)$$

74 To see this consider Hamiltonian dynamics but for the case that the trajectory Γ is *not* a solution of
 75 Hamilton equations, then $\tilde{\Gamma}$ is also not a solution, and both the probabilities $P[\Gamma]$ and $P[\tilde{\Gamma}]$ are trivially
 76 zero. Now consider the case when Γ is solution of Hamilton equations, then also $\tilde{\Gamma}$ is a solution. ut
 77 differently, because the dynamics are Hamiltonian, there is one and only one solution passing through
 78 the point Γ_0 at time $t = 0$, then the probability $P[\Gamma]$ is given by the probability to observe the system at
 79 Γ_0 at $t = 0$. By our equilibrium assumption this is given by $\varrho(\Gamma_0)$ ³. Likewise the $P[\tilde{\Gamma}]$ is given by $\varrho(\tilde{\Gamma}_0)$.
 80 Due to time-reversal symmetry and energy conservation we have $H(\tilde{\Gamma}_0) = H(\varepsilon\Gamma_\tau) = H(\Gamma_\tau) = H(\Gamma_0)$
 81 implying $\varrho(\tilde{\Gamma}_0) = \varrho(\Gamma_0)$, hence Eq. (3).

82 To summarize, the micro reversibility principle for autonomous systems in conjunction with the hy-
 83 pothesis of Gibbsian equilibrium implies that the probability to observe a trajectory and its time-reversal
 84 companion are equal. There is no way to distinguish between past and future in an autonomous system
 85 at equilibrium. Obviously, this is no longer so when the system is prepared out of equilibrium, as in Fig
 86 1, top.

87 2.2. Nonautonomous dynamics

88 Imagine now the nonautonomous case of a thermally insulated system driven through the variation
 89 of a parameter λ_t . Thermal insulation guarantees that the dynamics are still Hamiltonian. At variance
 90 with the autonomous case though, now the Hamiltonian is time dependent. Without loss of generality we
 91 assume that the varying parameter, denoted by λ_t couples linearly to some system observable $Q(\mathbf{q}, \mathbf{p})$,
 92 so that the Hamiltonian reads:

$$H(\mathbf{q}, \mathbf{p}; \lambda_t) = H_0(\mathbf{q}, \mathbf{p}) - \lambda_t Q(\mathbf{q}, \mathbf{p}) \quad (4)$$

93 This is the traditional form employed in the study of the fluctuation-dissipation theorem [12].⁴ In the
 94 following we shall reserve the symbol λ (without subscript) to denote the whole parameter variation
 95 protocol, and use the symbol λ_t , to denote the specific value taken by the parameter at time t . The
 96 succession of parameter values is assumed to be pre-specified (the system evolution does not affect the
 97 parameter evolution).

98 We assume that at time $t = 0$ the system is in the equilibrium Gibbs state

$$\varrho_0(\mathbf{q}, \mathbf{p}) = e^{-\beta H_0(\mathbf{q}, \mathbf{p})} / Z_0(\beta), \quad (5)$$


³To be more precise $P[\Gamma]\mathcal{D}\Gamma = \varrho(\Gamma_0)d\Gamma_0$ where $\mathcal{D}\Gamma$ is the measure on the Γ -trajectory space, and $d\Gamma_0$ is the measure in phase space

⁴For the sake of clarity we remark that the Hamiltonian describing the expansion of a gas, as depicted in Fig. 1, bottom, is not of this form. Our arguments however can be generalized to nonlinear couplings [10].

99 where $Z_0(\beta) = \int d\mathbf{q}d\mathbf{p}e^{-\beta H_0(\mathbf{q},\mathbf{p})}$, and that at any fixed value of the parameter the Hamiltonian is time
100 reversal symmetric:

$$H(\mathbf{q}, \mathbf{p}; \lambda_t) = H(\mathbf{q}, -\mathbf{p}; \lambda_t) \quad (6)$$

Figure 2. Microreversibility for nonautonomous classical (Hamiltonian) systems. The initial condition Γ_0 evolves to Γ_τ under the protocol λ , following the path Γ . The time-reversed final condition $\varepsilon\Gamma_\tau$ evolves to the time-reversed initial condition $\varepsilon\Gamma_0$ under the protocol $\tilde{\lambda}$, following the path $\tilde{\Gamma}$.



101 Note the prominent fact that energy is not conserved in the nonautonomous case because the Hamil-
102 tonian is time-dependent in this case. Microreversibility, as we have described it above, also does not
103 hold: Given a protocol λ , if Γ is a solution of the Hamilton equations of motion, in general $\tilde{\Gamma}$ is not.
104 However $\tilde{\Gamma}$ is a solution of the equations of motion generated by the time-reversed protocol $\tilde{\lambda}$, where
105 $\tilde{\lambda}_t = \lambda_{\tau-t}$. This is the *microreversibility principle for nonautonomous systems* [4]. It is illustrated in Fig.
106 2. Despite its importance we are not aware of any text-books in classical (or quantum) mechanics that
107 discusses it. A classical proof appears in [13, Sec. 1.2.3]. Corresponding quantum proofs were given in
108 Refs. [14] and [4, See appendix B].

109 As with the autonomous case we can ask how the probability distribution $P[\Gamma, \lambda]$ that the trajectory
110 Γ is realized under the protocol λ , compares with the probability distribution $P[\tilde{\Gamma}, \tilde{\lambda}]$ that the reversed
111 trajectory $\tilde{\Gamma}$ is realized under the reversed protocol $\tilde{\lambda}$. The answer to this was first given by Bochkov and
112 Kuzovlev [10], who showed that

$$P[\Gamma, \lambda] = P[\tilde{\Gamma}, \tilde{\lambda}]e^{\beta W_0} \quad (7)$$

where

$$W_0 = \int_0^\tau dt \lambda_t \dot{Q}_t. \quad (8)$$

113 Here, $Q_t = Q(\Gamma_t)$ denotes the evolution of the quantity Q along the trajectory Γ and W_0 is the so called
114 “exclusive work”. As discussed in [4,15–17] yet another definition of work is possible, the so called

115 “inclusive work” $W = - \int dt \dot{\lambda}_t Q_t$, leading to a different and equally important fluctuation theorem
 116 involving free energy differences [4,18,19]. Without entering the question about the physical meaning of
 117 the two quantities W and W_0 , it suffices for the present propose to notice that for a cyclic transformation
 118 $W_0 = W$.⁵ In the remaining of this section we will restrict our analysis to cyclic transformations
 119 ($\lambda_0 = \lambda_\tau$) in order to make contact with Kelvin postulate and to avoid any ambiguity regarding the
 120 usage of the word “work”.

121 Just like Eq. (3) is a direct expression of the principle of microreversibility for autonomous systems,
 122 so is Eq. (7) a direct expression of the more general principle of microreversibility for nonautonomous
 123 systems. Remarkably it expresses the second law in a most clear and refined way.

124 In order to see this it is important to realize that the work W_0 is odd under time-reversal. This is so
 125 because W_0 is linear in a quantity \dot{Q}_t , which is the time derivative of an even observable Q . The theorem
 126 says that the probability to observe a trajectory corresponding to some work $W_0 > 0$ under the driving
 127 λ is exponentially larger than the probability to observe the reversed trajectory (corresponding to $-W_0$)
 128 under the driving $\tilde{\lambda}$. This provides a statistical formulation of the second law

129 **Second Law (Fluctuation Theorem)** Injecting some amount of energy W_0 into a thermally insulated
 130 system at equilibrium at temperature T by the cyclic variation of a parameter, is exponentially (i.e.
 131 by a factor $e^{W_0/(k_B T)}$) more probable than withdrawing the same amount of energy from it by the
 132 reversed parameter variation.

133 Multiplying Eq. (7) by $e^{-\beta W_0}$ and integrating over all Γ -trajectories, leads to the relation [10]:

$$\langle e^{-\beta W_0} \rangle_\lambda = 1. \quad (9)$$

134 The subscript λ in Eq. (9) is there to recall that the average is taken over the trajectories generated by the
 135 protocol λ . In particular, the notation $\langle \cdot \rangle_\lambda$ denotes a nonequilibrium average. Combining Eq. (9) with
 136 Jensen’s inequality, $\langle \exp(x) \rangle \geq \exp(\langle x \rangle)$, leads to

$$\langle W_0 \rangle_\lambda \geq 0, \quad (10)$$

137 which now expresses Kelvin’s postulate as a nonequilibrium inequality. The quantum generalization of
 138 this fluctuation theorem and yet further relations have been given recently in Ref. [17].

139 3. Dissipation: Kubo’s formula

140 Before we continue with the implications of the fluctuation theorem for the arrow of time question, it
 141 is instructive to see in which way the fluctuation theorem relates to dissipation.

142 Given the distribution $P[\Gamma, \lambda]$, the distribution $p[Q, \lambda]$ that a trajectory Q of the observable $Q(\mathbf{q}, \mathbf{p})$
 143 occurs in the time span $[0, \tau]$, can be formally expressed as:

$$p[Q, \lambda] = \int \mathcal{D}\Gamma P[\Gamma, \lambda] \delta(Q - Q[\Gamma]) \quad (11)$$

⁵For a detailed discussion on the differences between the two work expressions we refer the readers to Sect. III. A in the colloquium [4].

144 where δ denotes Dirac's delta in the Q -trajectory space, the integration is a functional integration over
 145 all Γ -trajectories, and $Q[\Gamma]$ is defined as $Q[\Gamma]_t \doteq Q[\Gamma_t]$.

146 Multiplying Eq. (3) by $e^{-\beta \int \lambda_s \dot{Q}_s ds} \delta(Q - Q[\Gamma])$ and integrating over all Γ -trajectories, one finds:

$$p[Q, \lambda] e^{-\beta \int \lambda_s \dot{Q}_s ds} = p[\tilde{Q}, \tilde{\lambda}] , \quad (12)$$

147 where \tilde{Q} is the time reversal companion of Q : $\tilde{Q}_t = Q_{\tau-t}$. Now multiplying both sides of Eq. (12) by
 148 Q_τ and integrating over all Q -trajectories, one obtains:

$$\langle Q_\tau e^{-\beta \int \lambda_s \dot{Q}_s ds} \rangle_\lambda = \langle \tilde{Q}_\tau \rangle_{\tilde{\lambda}} \quad (13)$$

149 Note that $\langle \tilde{Q}_\tau \rangle_{\tilde{\lambda}} = \langle Q_0 \rangle_{\tilde{\lambda}}$ and that, due to causality, the value taken by the observable $Q(\mathbf{q}, \mathbf{p})$ at time
 150 $t = 0$ cannot be influenced by the subsequent evolution of the protocol $\tilde{\lambda}$. Therefore, the average presents
 151 a manifest equilibrium average; that is to say that it is an average over the initial canonical equilibrium
 152 $\rho_0(\mathbf{q}, \mathbf{p})$. We denote this equilibrium average by the symbol $\langle \cdot \rangle$ (with no subscript). Thus, Eq. (13) reads

$$\langle Q_\tau e^{-\beta \int \lambda_s \dot{Q}_s ds} \rangle_\lambda = \langle Q_0 \rangle \quad (14)$$

154 By expanding the exponential in Eq. (14) to first order in λ , one obtains:

$$\langle Q_\tau \rangle_\lambda - \langle Q_0 \rangle = \beta \left\langle Q_\tau \int_0^\tau \lambda_s \dot{Q}_s ds \right\rangle_\lambda + O(\lambda^2) . \quad (15)$$

155 Since the bracketed expression on the rhs is already $O(\lambda)$ we can replace the non-equilibrium average
 156 $\langle \cdot \rangle_\lambda$ with the equilibrium average $\langle \cdot \rangle$ on the rhs. Further, since averaging commutes with time integration
 157 one arrives, up to order $O(\lambda^2)$, at:

$$\langle Q_\tau \rangle_\lambda - \langle Q_0 \rangle = \beta \int_0^\tau \langle Q_\tau \dot{Q}_s \rangle \lambda_s ds , \quad (16)$$

$$= -\beta \int_0^\tau \langle \dot{Q}_{\tau-s} Q_0 \rangle \lambda_s ds . \quad (17)$$

158 In the second line we made use of the time-homogeneous nature of the equilibrium correlation func-
 159 tion. This is the celebrated Kubo formula [12] relating the non equilibrium linear response of the quantity
 160 Q to the equilibrium correlation function $\phi(s, \tau) = \langle Q_\tau \dot{Q}_s \rangle$. As Kubo showed it implies the fluctuation-
 161 dissipation relation [20], linking, for example, the mobility of a Brownian particle to its diffusion coeffi-
 162 cient [21], and the resistance of an electrical circuit to its thermal noise [22,23]

163 This classical derivation of Kubo's formula from the fluctuation theorem is a simplified version of the
 164 derivation given by Bochkov and Kuzovlev [10]. The corresponding quantum derivation was reported
 165 by Andrieux and Gaspard [14].

166 4. Implications for the arrow of time question


167 Jarzynski has analyzed in a transparent way how the fluctuation theorem for the inclusive work, W ,
 168 may be employed to make guesses about the direction of time's arrow [5]. Here we re-propose his
 169 argument and adapt it to the exclusive work, W_0 , fluctuation relation of Bochkov and Kuzovlev, Eq. (7).

170 Imagine we are shown a movie of an experiment in which a system starting at temperature $T =$
 171 $(k_B\beta)^{-1}$ is driven by a protocol, and we are asked to guess whether the movie is displayed in the same
 172 direction as it was filmed or in the backward direction. Imagine we can infer from the analysis of each
 173 single frame t the instantaneous values λ_t and Q_t taken by the parameter and its conjugate observable,
 174 respectively. With these we can calculate the work W_0 for the displayed process using Eq. (8). Imagine
 175 that we find, for the shown movie $\beta W_0 \gg 1$. If the film was shown in the “correct” direction it means
 176 that a process corresponding to $\beta W_0 \gg 1$ occurred. If the film was shown backward then it means that
 177 a process corresponding to $\beta W_0 \ll -1$ occurred (recall that W_0 is odd under time-reversal). The fluc-
 178 tuation theorem tells us that the former case occurs with an overwhelmingly higher probability relative
 179 to the probability of the latter case. Then we can be very much confident that the film was running in
 180 the correct direction. Likewise if we observe $\beta W_0 \ll -1$, then we can say with very much confidence
 181 the the film depicts the process in the opposite direction as it happened. Clearly when intermediate val-
 182 ues of βW_0 are observed we can still make well informed guesses about the direction of the movie, but
 183 with less confidence. The worst case arises when we observe $W_0 = 0$ in which case we cannot make
 184 any reliable guess. The question then arises of how to quantify the confidence of our guesses. In other
 185 words we have to quantify the likelihood $L_F[Q, \lambda]$ ($L_R[Q, \lambda]$) that the actual process occurred in the
 186 same (reversed) temporal order as it was shown, given the observed trajectory, Q and protocol λ .

187 Clearly $L_F[Q, \lambda]$ is proportional to the probability $p[Q, \lambda]$ that the observed process occurred. Like-
 188 wise, the likelihood $L_R[Q, \lambda] = 1 - L_F[Q, \lambda]$ is proportional to $p[\tilde{Q}, \tilde{\lambda}]$. Thus, normalizing over the two
 189 possibilities and using the fluctuation theorem (3), one finds

$$L_F[Q, \lambda] = \frac{p[Q, \lambda]}{p[Q, \lambda] + p[\tilde{Q}, \tilde{\lambda}]} = \frac{1}{e^{-\beta W_0} + 1} \quad (18)$$

Figure 3. Likelihood that a movie showing the nonautonomous evolution of a system is shown in the same temporal order as it was filmed, as a function of the observed work W_0 .



190 Figure 3 displays $L_F[Q, \lambda]$ as a function of W_0 . As it should be $L_F[Q, \lambda]$ is larger than 1/2 for
 191 positive W_0 , and vice versa, and is an increasing function of W_0 . If W_0 is large compared to β^{-1} , then
 192 $L_F[Q, \lambda] \simeq 1$, and we can be almost certain that the movie was shown in the forward direction. Vice
 193 versa, if $\beta W_0 \ll -1$, then we can say with almost certainty that the movie has been shown backward.
 194 The transition to certainty of guess occurs quite rapidly (in fact exponentially) around $|\beta W_0| \simeq 5$. Note

195 that that for an autonomous system $W_0 = 0$, implying $L_F[Q, \lambda] = L_R[Q, \lambda] = 1/2$, meaning that, as we
 196 have elaborated above, there is no way to discern the direction of time's arrow in an autonomous system
 197 at equilibrium.

198 Interestingly, since the fluctuation theorem (7) hold as a general law *regardless of the size of the*
 199 *system*, it appears that our ability to discern the direction of time's arrow does not depend on the system
 200 size. It is also worth mentioning the role played by thermal fluctuations in shaping our guesses. For a
 201 given observed value W_0 , the lower the temperature, the higher the confidence (and vice-versa).

202 5. Remarks

203 As we have mentioned in the introduction, traditionally the question of the emergence of the arrow of
 204 time from microscopic dynamics have been addressed within the framework of the Minus-First Law. In
 205 all existing approaches the arrow of time emerges from the introduction of some *extra ingredient* which
 206 in turn then dictates the time direction. Typically, one resorts to a coarse-graining procedure of the
 207 microscopic phase space to describe some state variables. For example, this is so in the theory of Gibbs
 208 and related approaches, see, e.g., in Ref. [24]. The time arrow is then generated via the observation that
 209 such coarse grained quantities no longer obey time-reversal symmetric Hamiltonian dynamics. More
 210 frequently, one resorts to additional assumptions which are of a probabilistic nature: Typical scenarios
 211 that come to mind are (i) the use of Boltzmann Stoßzahlansatz in the celebrated Boltzmann kinetic theory,
 212 (ii) the assumption of initial molecular chaos in more general kinetic theories that are in the spirit of
 213 Bogoliubov, or, likewise, with Fokker-Planck and master equation dynamics that no longer exhibit an
 214 explicit time-reversal invariant structure [24,25]. All such additional elements then induce the result of
 215 a *direction in time* with *future* not being equivalent with *past* any longer.

216 Having stressed the too often overlooked fact that the Second Law does not refer to the tradition-
 217 ally considered scenario of autonomously evolving systems, but rather to the case of nonautonomous
 218 dynamics, here we have focussed on the emergence of time's arrow in a driven system starting at equi-
 219 librium. Having based our derivation on the principle of nonautonomous microreversibility, Fig. 2, the
 220 question arises naturally regarding the origin of the time asymmetry in this case. It originates from
 221 the combination of the following two elements: i) The introduction of an explicit time dependence of
 222 the Hamiltonian, Eq. (4), ii) The particular shape of the initial equilibrium state, Eq. (5). The first
 223 breaks time homogeneity thus determining the emergence of an arrow of time, while the second deter-
 224 mines its direction. It is in particular the fact that the initial equilibrium is described by a probability
 225 density function which is a *decreasing* function of energy, that determines the \geq sign in Eq. (10). An
 226 increasing probability density function would result in the opposite sign [6,26,27]. In regard to breaking
 227 time homogeneity, it is worth commenting that the assumption of nonautonomous evolution has to be
 228 regarded itself as a convenient and often extremely good *approximation* in which the evolution λ of the
 229 external parameter influences the system dynamics without being influenced minimally by the system.⁶
 230 This indeed presupposes the intervention of a sort of Maxwell Demon (i.e., the experimentalist), who
 231 predisposes things in such a way that the wanted protocol actually occurs. This in turn evidences the

⁶ In principle one should treat the external parameter itself as a dynamical coordinate, and consider the autonomous evolution of the extended system.

232 phenomenological nature of the Second Law. It is not a law that dictates how things go by themselves,
233 but rather how they go in response to particular experimental investigations.

234 Acknowledgements

235 This work was supported by the cluster of excellence Nanosystems Initiative Munich (NIM) and the
236 Volkswagen Foundation (project I/83902).

237 References

- 238 1. Brown, H.R.; Uffink, J. The Origins of Time-Asymmetry in Thermodynamics: The Minus First
239 Law. *Stud. Hist. Phil. Mod. Phys.* **2001**, *32*, 525 – 538.
- 240 2. Campisi, M, On the mechanical foundations of thermodynamics: the generalized Helmholtz the-
241 orem *Stud. Hist. Phil. Mod. Phys.* **2005**, *36*, 275 – 290.
- 242 3. Campisi, M.; Kobe, D.H. Derivation of the Boltzmann principle. *Am. J. Phys.* **2010**, *78*, 608–
243 615.
- 244 4. Campisi, M.; Hänggi, P.; Talkner, P. Colloquium: Quantum fluctuation relations: Foundations
245 and applications. *Rev. Mod. Phys.* **2011**, *83*, 771–791.
- 246 5. Jarzynski, C. Equalities and Inequalities: Irreversibility and the Second Law of Thermodynamics
247 at the Nanoscale. *Annu. Rev. Condens. Matter. Phys.* **2011**, *2*, 329–351.
- 248 6. Allahverdyan, A.; Nieuwenhuizen, T. A mathematical theorem as the basis for the second law:
249 Thomson’s formulation applied to equilibrium. *Physica A* **2002**, *305*, 542 – 552.
- 250 7. Campisi, M.; Talkner, P.; Hänggi, P. Finite bath fluctuation theorem. *Phys. Rev. E* **2009**,
251 *80*, 031145.
- 252 8. Esposito, M.; Harbola, U.; Mukamel, S. Nonequilibrium fluctuations, fluctuation theorems, and
253 counting statistics in quantum systems. *Rev. Mod. Phys.* **2009**, *81*, 1665–1702.
- 254 9. Seifert, U. Stochastic thermodynamics: principles and perspectives. *Eur. Phys. J. B* **2008**,
255 *64*, 423–431.
- 256 10. Bochkov, G.N.; Kuzovlev, Y.E. General theory of thermal fluctuations in nonlinear systems. *Zh.*
257 *Eksp. Teor. Fiz.* **1977**, *72*, 238–247. [Sov. Phys. JETP **45**, 125–130 (1977)].
- 258 11. Messiah, A. *Quantum Mechanics*; North Holland: Amsterdam, 1962.
- 259 12. Kubo, R. Statistical-mechanical theory of irreversible processes. I. *J. Phys. Soc. Jpn.* **1957**,
260 *12*, 570–586.
- 261 13. Stratonovich, R.L. *Nonlinear nonequilibrium thermodynamics II : advanced theory*; Vol. 59,
262 *Springer series in synergetics*, Springer-Verlag: Berlin ; Heidelberg; New York, 1994.
- 263 14. Andrieux, D.; Gaspard, P. Quantum Work Relations and Response Theory. *Phys. Rev. Lett.*
264 **2008**, *100*, 230404.
- 265 15. Jarzynski, C. Comparison of far-from-equilibrium work relations. *C. R. Phys.* **2007**, *8*, 495–506.
- 266 16. Horowitz, J.; Jarzynski, C. Comparison of work fluctuation relations. *J. Stat. Mech.: Theory*
267 *Exp.* **2007**, p. P11002.
- 268 17. Campisi, M.; Talkner, P.; Hänggi, P. Quantum Bochkov-Kuzovlev work fluctuation theorems.
269 *Phil. Trans. R. Soc. A* **2011**, *369*, 291–306.

- 270 18. Jarzynski, C. Nonequilibrium equality for free energy differences. *Phys. Rev. Lett.* **1997**,
271 78, 2690–2693.
- 272 19. Crooks, G.E. Entropy production fluctuation theorem and the nonequilibrium work relation for
273 free energy differences. *Phys. Rev. E* **1999**, *60*, 2721–2726.
- 274 20. Callen, H.B.; Welton, T.A. Irreversibility and Generalized Noise. *Phys. Rev.* **1951**, *83*, 34–40.
- 275 21. Einstein, A. *Investigations on the Theory of Brownian Movement*; Methuen: London, 1926.
276 reprinted (Dover, New York), 1956.
- 277 22. Johnson, J.B. Thermal agitation of electricity in conductors. *Phys. Rev.* **1928**, *32*, 97–109.
- 278 23. Nyquist, H. Thermal agitation of electric charge in conductors. *Phys. Rev.* **1928**, *32*, 110–113.
- 279 24. Wu, T.-Y. On the Nature of Theories of Irreversible Processes. *Int. J. Theor. Phys.* **1969**, *2*, 325–
280 343.
- 281 25. Hänggi, P.; Thomas H. Stochastic Processes: Time-Evolution, Symmetries and Linear Response,
282 *Phys. Rep.* **1982**, *88*, 207–319.
- 283 26. Campisi, M. Statistical Mechanical Proof of the Second Law of Thermodynamics Based on Vol-
284 ume Entropy. *Stud. Hist. Phil. Mod. Phys.* **2008**, *39*, 181–194.
- 285 27. Campisi, M. Increase of Boltzmann entropy in a quantum forced harmonic oscillator. *Phys. Rev.*
286 *E* **2008**, *78*, 051123.

287 © November 8, 2011 by the authors; submitted to *Entropy* for possible open access publication under the
288 terms and conditions of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/3.0/>.