

# Information Causality, the Tsirelson Bound, and the ‘Being-Thus’ of Things\*

Michael E. Cuffaro<sup>a, b</sup>

<sup>a</sup>Rotman Institute of Philosophy, University of Western Ontario

<sup>b</sup>Munich Center for Mathematical Philosophy, Ludwig-Maximilians-Universität München

August 29, 2017

## Abstract

The principle of ‘information causality’ can be used to derive an upper bound—known as the ‘Tsirelson bound’—on the strength of quantum mechanical correlations, and has been conjectured to be a foundational principle of nature. In this paper, however, I argue that the principle has not to date been sufficiently motivated to play this role; the motivations that have so far been given are either unsatisfactorily vague or else amount to little more than an appeal to intuition. I then consider how one might begin to successfully motivate the principle. I argue that a compelling way of so doing is to understand it as a generalisation of Einstein’s principle of the mutually independent existence—the ‘being-thus’—of spatially distant things, (re-)interpreted as a special methodological principle. More specifically: I describe an argument, due to Demopoulos, to the effect that the quantum-mechanical no-signalling condition can be viewed as a generalisation, appropriate to an irreducibly statistical theory such as quantum mechanics, of the Einsteinian principle. And I then argue that a compelling way to motivate information causality is to in turn consider it as a further generalisation of the Einsteinian principle that is appropriate to a theory of communication. I nevertheless describe important obstacles that must yet be overcome if the project of establishing information causality as a foundational principle of nature is to succeed.

## 1 Introduction

Answering the question of precisely what distinguishes our experience with quantum as opposed to classical physical phenomena has historically been a central element of the overall project of interpreting quantum theory. For Schrödinger (1935), for instance, the sole distinguishing feature of quantum theory was none other than entanglement, while for Feynman

---

\*I am indebted to the late William Demopoulos for sharing a draft of the chapter “Quantum Reality” with me from his as of yet unpublished monograph *On Theories*. The present paper is motivated, illuminated, and informed both by that chapter and by my conversations with Bill during the years 2013–2017. Many thanks, also, to Jeffrey Bub, Markus Müller, and Ryan Samaroo for their very helpful comments on a previous draft of this paper and for discussion. I do not claim (though I hope for) the endorsement of my conclusions by any of the above. This project was supported financially by the Foundational Questions Institute (FQXi) and by the Rotman Institute of Philosophy.

the one and only quantum mystery was self-interference (Feynman et al., 1964, vol. 3, 1-1). The question continues to occupy many. However in much of the more recent literature it has taken on a different form. That is, it has become one of specifying a set of appropriately motivated constraints or ‘principles’ that serve to distinguish quantum from classical theory. Clifton, Bub, & Halvorson (2003), for instance, prove a theorem which they argue shows quantum mechanics to be essentially characterisable in terms of a small number of information-theoretic constraints. Spekkens (2007), meanwhile, shows that features often thought of as distinctively quantum can be manifested in a toy classical theory to which one adds a principled restriction on the maximal obtainable knowledge of a system.<sup>1</sup>

One feature that quantum and classical theory have in common is that the correlations manifested between the subsystems of a combined system satisfy the condition that the marginal probabilities associated with local experiments on a subsystem are independent of which particular experiments are performed on the other subsystems. It is a consequence of this condition that it is impossible to use either a classically correlated or entangled quantum system to signal faster than light. For this reason the condition is referred to as the ‘no-signalling’ condition or principle, even though the condition is not a relativistic constraint *per se*.

Quantum and classical theory do not exhaust the conceivable ways in which the world could be. The world could be such that neither quantum nor classical theory are capable of adequately describing the correlations between subsystems of combined systems. In particular the world could be such that correlations *stronger* than quantum correlations—those that exceed the so-called ‘Tsirelson bound’ (Tsirelson, 1980)—are possible within it. In a landmark paper, Popescu & Rohrlich (1994) asked the question of whether all such correlations must violate the no-signalling condition. The surprising answer to this question is no. As they showed, there do indeed exist conceivable correlations between the subsystems of combined systems that are stronger than quantum and yet non-signalling.

Popescu & Rohrlich’s result raises the question of whether some motivated principle or principles can be given which would pick out quantum theory—or at least some restricted subset of theories which includes quantum theory—from among the space of conceivable non-signalling physical theories in which correlations at or above the Tsirelson bound occur. This question has developed into an active research program. A particularly important result emerging from it is that of Pawłowski et al. (2009), who show that one can in fact derive the Tsirelson bound from a principle they call ‘information causality’, which they describe as a generalisation of no-signalling applicable to experimental setups in which the subsystems of a combined system (e.g. spatially separated labs) may be subluminally communicating classical information with one another. Pawłowski et al. conjecture that information causality may be a foundational principle of nature.

Below I will argue that, suitably interpreted (Bub, 2012), the principle can be regarded as a useful and illuminating answer to the question of what the Tsirelson bound expresses about correlations which exceed it. However I will argue that if one wishes to think of information causality as a fundamental principle of nature—in the sense that theories which violate the principle should thereby be regarded as unphysical or in some other sense impossible—then it requires more in the way of motivation than has hitherto been given.

What has typically been appealed to previously to motivate the principle is the intuition that a world in which information causality is not satisfied would be ‘too simple’ (Pawłowski et al., 2009, p. 1101), or ‘too good to be true’ (Bub 2012, p. 180, Bub 2016, p. 187); that

---

<sup>1</sup>For a discussion of both Clifton et al.’s and Spekkens’ results, and of the project in general, see Myrvold (2010); and see also Feline (2016).

it would allow one to “implausibly” access remote data (Pawłowski et al., 2009, *ibid.*), and that “things like this should not happen” (Pawłowski & Scarani, 2016, p. 429). I will argue below that these statements are unsatisfactorily vague. Nevertheless I will argue that they gesture at something that is importantly right; although they are right in, perhaps, a different sense than their authors envision.

More specifically, in contrast to Bub (2012), who in his otherwise illuminating analysis of information causality argues that it is misleadingly characterised as a generalisation of the no-signalling principle, I will argue that information causality can indeed be regarded as generalising no-signalling in a sense. To clarify this sense I will draw on the work of Demopoulos,<sup>2</sup> who convincingly shows that no-signalling can itself be thought of as a generalisation, appropriate for an irreducibly statistical theory such as quantum mechanics, of Einstein’s principle of the mutually independent existence of spatially distant things. Einstein regarded this principle as necessary for the very possibility of ‘physical thought’, and argued that it is violated by quantum mechanics (Howard, 1985, p. 187). However, suitably generalised and interpreted methodologically, Demopoulos convincingly argues that Einstein’s principle is satisfied both in Newtonian mechanics (despite its being an action-at-a-distance theory), and indeed (somewhat ironically) that it is satisfied in quantum mechanics, wherein it is expressed by none other than the no-signalling condition.

Coming back to information causality, I will then argue that it can likewise be thought of as a further generalisation of Einstein’s principle that is appropriate for a theory of communication. As I will clarify, in the context of the experimental setups to which the principle is applicable, a failure of information causality would imply an ambiguity in the way one distinguishes conceptually between a sender and a receiver of information. This ambiguity (arguably) makes communication theory as we know it in the context of such setups impossible, similarly to the way in which a failure of the principle of the mutually independent existence of spatially distant things (arguably) makes physical theory as we know it impossible.

Before beginning let me note that the general approach represented by the investigation into information causality is only one of a number of principle-theoretic approaches that one can take regarding the question of how to distinguish quantum from super-quantum theories. In the kind of approach exemplified by the investigation into information causality, one focuses on sets of static correlation tables associated with quantum and super-quantum theories, and in particular one disregards the dynamics of (super-)quantum systems. There is another family of principle-theoretic approaches to the question, however, wherein a richer framework is considered that does include dynamics.<sup>3</sup> Popescu & Rohrlich’s seminal (1994) investigation is an example of the former type of approach, though they themselves consider the latter, dynamical, approach to have the potential for deeper insight. For my part I do not consider any particular approach to be superior. Principle-theoretic approaches to the characterisation of quantum theory augment our understanding of the world by illuminating various aspects of it to us. Which particular aspect of the world is illuminated by an investigation will depend upon the particular question—and the framework which defines it—that is asked. I am highly skeptical of the idea that any one framework is sufficient by itself to illuminate all.

The rest of this paper will proceed as follows: I will introduce Popescu-Rohrlich (PR)

---

<sup>2</sup>I am referring to the chapter “Quantum Reality” of Demopoulos’s monograph *On Theories*, which is currently being prepared for posthumous publication.

<sup>3</sup>For further references, as well as an accessible description of one of these reconstructions of quantum theory, see Koberinski & Müller (forthcoming).

correlations in §2. In §3 I will introduce the ‘guessing game’ by which the principle of information causality is standardly operationally defined. The principle of information causality itself will be introduced in §4, wherein I will also describe how it can be used to derive the Tsirelson bound. I will argue in that section that information causality has not been sufficiently motivated to play the role of a foundational principle of nature, and in the remainder of the paper I will consider how one might begin to provide it with such a motivation. This analysis begins in §5 where I describe an argument, due to Demopoulos, to the effect that the no-signalling condition can be viewed as a generalisation, appropriate to an irreducibly statistical theory, of Einstein’s principle of mutually independent existence interpreted methodologically. Then in §6 I argue that a promising route toward successfully motivating information causality is to in turn consider it as a further generalisation of no-signalling that is appropriate to a theory of communication. I describe, however, some important obstacles that must yet be overcome if the project of establishing information causality as a foundational principle of nature is to succeed.

## 2 Popescu-Rohrlich correlations

Consider a correlated state  $\sigma$  of two two-level subsystems.<sup>4</sup> Let Alice and Bob each be given one of the subsystems, and instruct them to travel to distinct distant locations. Let  $p(A, B|a, b)$  be the probability that Alice and Bob obtain outcomes  $A$  and  $B$ , respectively, after measuring their local subsystems with the respective settings  $a$  and  $b$ . If  $A, B \in \{\pm 1\}$ , the expectation value of the outcome of their combined measurement is given by:

$$\langle a, b \rangle = \sum_{i, j \in \{1, -1\}} (i \cdot j) \cdot p(i, j|a, b),$$

where  $A = i$  and  $B = j$ . Less concisely, this is:

$$\begin{aligned} \langle a, b \rangle &= 1 \cdot p(1, 1|a, b) - 1 \cdot p(1, -1|a, b) - 1 \cdot p(-1, 1|a, b) + 1 \cdot p(-1, -1|a, b) \\ &= p(\text{same}|a, b) - p(\text{different}|a, b). \end{aligned}$$

Since  $p(\text{same}|a, b) + p(\text{different}|a, b) = 1$ , it follows that  $\langle a, b \rangle + 2 \cdot p(\text{different}|a, b) = 1$ , so that:

$$p(\text{different}|a, b) = \frac{1 - \langle a, b \rangle}{2}.$$

Similarly, we have that

$$p(\text{same}|a, b) = \frac{1 + \langle a, b \rangle}{2}.$$

Now imagine that  $\sigma$  is such that the probabilities for the results of experiments with settings  $a, b, a', b'$ , where  $a'$  and  $b'$  are different from  $a$  and  $b$  but arbitrary (Popescu & Rohrlich, 1994, p. 382), are:

$$\begin{aligned} p(1, 1|a, b) &= p(-1, -1|a, b) = 1/2, \\ p(1, 1|a, b') &= p(-1, -1|a, b') = 1/2, \\ p(1, 1|a', b) &= p(-1, -1|a', b) = 1/2, \\ p(1, -1|a', b') &= p(-1, 1|a', b') = 1/2. \end{aligned} \tag{2.1}$$

---

<sup>4</sup>Elements of the exposition in this and the next section have been adapted from Bub (2012, 2016) and Pawłowski et al. (2009).

In other words, if at least one of their settings is one of  $a$  or  $b$ , then Alice's and Bob's results are guaranteed to be the same. Otherwise they are guaranteed to be different. These correlations are called 'PR' correlations after Popescu & Rohrlich (1994).

Alice's marginal probability  $p(1_A|a, b)$  of obtaining the outcome 1 given that she measures  $a$  and Bob measures  $b$  is defined as:  $p(1_A, 1_B|a, b) + p(1_A, -1_B|a, b)$ . The no-signalling condition requires that her marginal probability of obtaining 1 is the same irrespective of whether Bob measures  $b$  or  $b'$ , i.e. that  $p(1_A|a, b) = p(1_A|a, b')$ , in which case we can write her marginal probability simply as  $p(1_A|a)$ . In general, no-signalling requires that

$$\begin{aligned} p(A|a, b) &= p(A|a, b'), & p(A|a', b) &= p(A|a', b'), \\ p(B|a, b) &= p(B|a', b), & p(B|a, b') &= p(B|a', b'). \end{aligned} \quad (2.2)$$

The reader can verify that the PR correlations (2.1) satisfy the no-signalling condition (2.2).

If we imagine trying to simulate the PR correlations (2.1) with some bipartite general non-signalling system  $\eta$ , then the probability of a successful simulation (assuming a uniform probability distribution over the possible joint measurements  $(a, b)$ ,  $(a, b')$ ,  $(a', b)$ , and  $(a', b')$ ) is given by:

$$\begin{aligned} & \frac{1}{4}(p(\text{same}|a, b) + p(\text{same}|a, b') + p(\text{same}|a', b) + p(\text{different}|a', b')) \\ &= \frac{1}{4} \left( \frac{1 + \langle a, b \rangle}{2} + \frac{1 + \langle a, b' \rangle}{2} + \frac{1 + \langle a', b \rangle}{2} + \frac{1 - \langle a', b' \rangle}{2} \right) \\ &= \frac{1}{2} \left( 1 + \frac{\langle a, b \rangle + \langle a, b' \rangle + \langle a', b \rangle - \langle a', b' \rangle}{4} \right). \end{aligned}$$

Notice that  $\langle a, b \rangle + \langle a, b' \rangle + \langle a', b \rangle - \langle a', b' \rangle$  is just the Clauser-Horne-Shimony-Holt (CHSH) correlation expression (Clauser et al., 1969). So the probability of a successful simulation of the PR correlations by  $\eta$  is:

$$p(\text{successful sim}) = \frac{1}{2} \left( 1 + \frac{\text{CHSH}}{4} \right), \quad (2.3)$$

with  $\text{CHSH} = 4$  if  $\eta$  is itself a PR-system.<sup>5</sup> As is well known, classically correlated systems are bounded by  $|\text{CHSH}| \leq 2$ . Thus the optimum probability of simulating PR correlations with a bipartite classical system is given by  $1/2(1 + 2/4) = 3/4$ . Quantum correlations are bounded by  $|\text{CHSH}| \leq 2\sqrt{2}$ .

### 3 Alice and Bob play a guessing game

At this point it will be convenient to change our notation. From now on I will refer to the measurement settings  $a$  and  $a'$  as 0 and 1, respectively, and likewise for  $b$  and  $b'$ . The outcomes 1 and -1 will also be respectively relabelled as 0 and 1. This will allow us to

---

<sup>5</sup>The reader may be familiar with the use of the term 'PR-box' to refer to systems whose subsystems are correlated as in (2.1). I find the term 'box' to be misleading since it conveys the idea of a spatially contiguous region occupied by a combined system. Bub's (2016) banana imagery is far less misleading in this sense. Below I will not use figurative language at all, but will (boringly) refer merely to such entities as 'PR-systems', 'PR-correlated systems', and so on.

describe PR correlations more abstractly using the exclusive-or (alternately: modulo two addition) operator as follows:

$$M_1 \oplus M_2 = m_1 \cdot m_2 \quad (3.1)$$

where capital letters refer to measurement outcomes and small letters to measurement settings. To illustrate, for a given 01-experiment (formerly  $(a, b')$ ) there are two possible outcomes: 00 and 11 (formerly:  $(1,1)$  and  $(-1,-1)$ ), and we have:  $0 \oplus 0 = 0 \cdot 1$  and  $1 \oplus 1 = 0 \cdot 1$ , respectively.

Now imagine the following game. At the start of each round of the game, Alice and Bob receive random and independently generated bit strings  $\mathbf{a} = a_{N-1}, a_{N-2}, \dots, a_0$  and  $\mathbf{b} = b_{n-1}, b_{n-2}, \dots, b_0$ , respectively, with  $N = 2^n$ . They win a round if Bob is able to guess the value of the  $\mathbf{b}^{\text{th}}$  bit in Alice's list. For example, suppose Alice receives the string  $a_7 a_6 a_5 a_4 a_3 a_2 a_1 a_0$ , and Bob receives the string 110. Then Bob must guess the value of  $a_6$ . They win the game if Bob is able to guess correctly over any sequence of rounds.

Besides this the rules of the game are as follows. Before the game starts, Alice and Bob are allowed to determine a mutual strategy and to prepare and share non-signalling physical resources such as classically correlated systems, or quantum systems in entangled states, or PR-systems, or other (bipartite) systems manifesting non-signalling correlations. They then go off to distinct distant locations, taking with them their portions of whatever systems were previously prepared. Once separated, Alice receives her bit string  $\mathbf{a}$  and Bob his bit string  $\mathbf{b}$ . She is then allowed to send Bob one additional classical bit  $c$ , upon receipt of which Bob must guess the value of Alice's  $\mathbf{b}^{\text{th}}$  bit.

Alice and Bob can be certain to win the game if they share a number of PR-systems. I will illustrate the case of  $N = 4$ , which requires three PR-systems (per round) labelled **I**, **II**, and **III**. Upon receiving the bit string  $\mathbf{a} = a_3 a_2 a_1 a_0$ , Alice measures  $a_0 \oplus a_1$  on her part of system **I** and gets the result  $A_I$ . She then measures  $a_2 \oplus a_3$  on her part of system **II** and gets the outcome  $A_{II}$ . She then measures  $(a_0 \oplus A_I) \oplus (a_2 \oplus A_{II})$  on her part of system **III** and gets the result  $A_{III}$ . She finally sends  $c = a_0 \oplus A_I \oplus A_{III}$  to Bob. Meanwhile, Bob, who has previously received  $\mathbf{b} = b_1 b_0$ , measures  $b_0$  on his parts of systems **I** and **II**, and gets back the results  $B_I$  and  $B_{II}$ . He also measures  $b_1$  on system **III** with the result  $B_{III}$ .

Bob's next step depends on the value of  $\mathbf{b}$ , i.e. on which of Alice's bits he has to guess. When  $\mathbf{b} = b_1 b_0 = 00$  (i.e. when Bob must guess the 0<sup>th</sup> bit) or  $\mathbf{b} = b_1 b_0 = 01$  (i.e. when Bob must guess the 1<sup>st</sup> bit) his guess should be:

$$c \oplus B_{III} \oplus B_I = a_0 \oplus A_I \oplus A_{III} \oplus B_{III} \oplus B_I. \quad (3.2)$$

For since  $A_{III} \oplus B_{III} = ((a_0 \oplus A_I) \oplus (a_2 \oplus A_{II})) \cdot b_1$ , we have:

$$\begin{aligned} & a_0 \oplus A_I \oplus A_{III} \oplus B_{III} \oplus B_I \\ &= a_0 \oplus A_I \oplus b_1(a_0 \oplus A_I) \oplus b_1(a_2 \oplus A_{II}) \oplus B_I \\ &= a_0 \oplus A_I \oplus B_I \\ &= a_0 \oplus b_0(a_0 \oplus a_1). \end{aligned} \quad (3.3)$$

If  $\mathbf{b} = 00$  then (3.3) correctly yields  $a_0$ . If  $\mathbf{b} = 01$  then (3.3) correctly yields  $a_1$ .

Suppose instead that  $\mathbf{b} = 10$  or  $\mathbf{b} = 11$ . In this case, Bob's guess should be

$$c \oplus B_{III} \oplus B_{II} = a_0 \oplus A_I \oplus A_{III} \oplus B_{III} \oplus B_{II}. \quad (3.4)$$

This is

$$\begin{aligned}
&= a_0 \oplus A_I \oplus b_1(a_0 \oplus A_I) \oplus b_1(a_2 \oplus A_{II}) \oplus B_{II} \\
&= (a_0 \oplus A_I) \oplus (a_0 \oplus A_I) \oplus (a_2 \oplus A_{II}) \oplus B_{II} \\
&= a_2 \oplus A_{II} \oplus B_{II} \\
&= a_2 \oplus b_0(a_2 \oplus a_3). \tag{3.5}
\end{aligned}$$

If  $\mathbf{b} = 11$  then (3.5) correctly yields  $a_3$ . If  $\mathbf{b} = 10$  then (3.5) correctly yields  $a_2$ .

In general, given  $N - 1$  PR-correlated systems per round,<sup>6</sup> and a single classical bit per round communicated by Alice to Bob, Alice and Bob can be certain to win the game for any value of  $N$ . In other words, given these resources and a single classical bit communicated to him by Alice, Bob can access the value of any single bit from her data set, however large that data set is. This result further generalises to the case where Alice is allowed to send not just one but  $m$  bits  $c_{m-1} \dots c_0$  to Bob in a given round, and Bob is required to guess an arbitrary set of  $m$  bits from Alice's data set. Note that if Alice is not allowed to send anything to Bob, i.e., when  $m = 0$ , then Bob will not be able to access the values of any of Alice's bits irrespective of how many PR-systems they share. This is a consequence of the fact that PR-correlations satisfy the no-signalling principle (2.2).

## 4 Information causality and the Tsirelson bound

As we saw in the last section, Alice and Bob can be certain to win the guessing game described there if they share a number of PR-correlated systems prior to going off to their respective locations. Note that if they do not use any correlated resources, they can still be sure to win the occasional round if Alice always sends Bob the value of whatever bit is at a previously agreed-upon fixed position  $a_k$  in her list. In this case, Bob will be guaranteed to guess correctly whenever  $b = k$  (but only then; otherwise he must rely on blind luck). If Alice and Bob share a sequence of classically correlated random bits, on the other hand, then Bob will be able to access the value of a single in general different  $a_i$  in Alice's list on each round.

Now consider the case where Alice and Bob share general no-signalling systems, i.e. bipartite systems such that the correlations between their subsystems satisfy the no-signalling condition. Recall that the probability that a non-signalling system simulates a PR-system on a given run depends on the value of CHSH in (2.3) that is associated with it. For convenience we will define  $E =_{df} \text{CHSH}/4$  so that (2.3) becomes:

$$p(\text{successful sim}) = \frac{1}{2}(1 + E). \tag{4.1}$$

When  $E = 1$  for a given non-signalling system, then it just is a PR-system, and the probability of a successful simulation is 1. When  $E < 1$ , then for given settings  $m_1, m_2$ , the values of the outputs  $M_1, M_2$ , will in general not satisfy the relation (3.1), i.e.  $M_1 \oplus M_2$  will not always equal  $m_1 \cdot m_2$ . For a given attempted simulation, let us say that  $M_2$  is 'correct' whenever (3.1) holds, and 'incorrect' otherwise.<sup>7</sup>

<sup>6</sup>These are to be arranged in an inverted pyramid so that the results of Alice's (respectively, Bob's) local measurements on the first  $2^{n-1}$  PR-systems are used to determine the local settings for her (his) next  $2^{n-2}$  measurements, and so on, for  $(n - i) \geq 0$ . Note that the cost in the number of PR-systems needed scales exponentially with respect to the length of  $\mathbf{b}$ . I will return to this point later.

<sup>7</sup>There is of course no reason why we should not say that  $M_1$  rather than  $M_2$  is incorrect, but for the analysis that follows it is convenient to take Bob's point of view.

Recall that in the  $N = 4$  game above, at the end of each round, Bob guesses either (i)  $c \oplus B_{III} \oplus B_I$ , or (ii)  $c \oplus B_{III} \oplus B_{II}$ , depending on the value of  $\mathbf{b}$ . We will consider only case (i), as the analysis is similar for (ii). If both  $B_I$  and  $B_{III}$  are ‘correct’, then for that particular round, the non-signalling systems will have yielded the same guess for Bob as PR-systems would have yielded:

$$(c \oplus B_{III} \oplus B_I)_{NS} = (c \oplus B_{III} \oplus B_I)_{PR}. \quad (4.2)$$

Note that if *both*  $B_I$  and  $B_{III}$  are *incorrect*, (4.2) will still hold, since in general  $x_1 \oplus x_2 = \overline{x_1} \oplus \overline{x_2}$ . So either way Bob will guess right. The probability of an unsuccessful simulation is

$$1 - \frac{1}{2}(1 + E) = \frac{1}{2}(1 - E).$$

Thus the probability that Bob makes the right guess on a given round in the  $N = 4$  game is:

$$\left(\frac{1}{2}(1 + E)\right)^2 + \left(\frac{1}{2}(1 - E)\right)^2 = \frac{1}{2}(1 + E^2).$$

In the general case, for  $N = 2^n$ , one can show (Pawłowski et al., 2009; Bub, 2012, 2016) that the probability that Bob correctly guesses Alice’s  $\mathbf{b}^{\text{th}}$  bit is

$$p_{\mathbf{b}} = \frac{1}{2}(1 + E^n). \quad (4.3)$$

The binary entropy  $h(p_{\mathbf{b}})$  associated with  $p_{\mathbf{b}}$  is given by

$$h(p_{\mathbf{b}}) = -p_{\mathbf{b}} \log_2 p_{\mathbf{b}} - (1 - p_{\mathbf{b}}) \log_2 (1 - p_{\mathbf{b}}).$$

In the case where Bob has no information about Alice’s  $\mathbf{b}^{\text{th}}$  bit,  $p_{\mathbf{b}} = 1/2$  and  $h(p_{\mathbf{b}}) = 1$ . If Alice then sends Bob  $m$  bits, then in general Bob’s information about that bit will increase by some non-zero amount. Pawłowski et al. (2009) propose the following constraint on this quantity, which they call the ‘information causality’ principle:

The information gain that Bob can reach about a previously unknown to him data set of Alice, by using all his local resources and  $m$  classical bits communicated by Alice, is at most  $m$  bits (2009, p. 1101).

For example, assuming that the  $N = 2^n$  bits in Alice’s bit string  $\mathbf{a}$  are unbiased and independently distributed, then if Alice sends Bob a single bit (i.e. when  $m = 1$ ), information causality asserts that Bob’s information about the  $\mathbf{b}^{\text{th}}$  bit in Alice’s string may increase by no more than  $1/2^n$ , i.e.,

$$h(p_{\mathbf{b}}) \geq 1 - \frac{1}{2^n}. \quad (4.4)$$

As Pawłowski et al. (2009) show,<sup>8</sup> the principle is satisfied within quantum mechanics. But within any theory which permits correlations with a value of  $E$  exceeding  $1/\sqrt{2}$  (i.e. any theory which allows correlations above the Tsirelson bound), one can find an  $n$  such that for a given  $m$  the principle is violated (for example, let  $E = .72$ ,  $m = 1$ , and  $n = 10$ ).

Given that any correlations above the Tsirelson bound will demonstrably violate the principle in this sense, it is tempting to view information causality as the answer to the

---

<sup>8</sup>See also Pawłowski & Scarani (2016).



question (i) of why nature does not allow correlations above this bound. And since the Tsirelson bound represents the maximum value of the CHSH expression for quantum correlations, one is further tempted to view information causality as the answer to the question (ii) of why only quantum correlations are allowable in nature. Indeed, Pawłowski et al. suggest that information causality “might be one of the foundational properties of nature” (2009, p. 1101).

There is a subtlety here, however. The set of (bipartite) quantum correlations forms a convex set which can be represented as a multi-dimensional region of points such that the points within this region that are furthest from the centre are at the Tsirelson bound (Bub, 2016, §5.1). Information causality disallows correlations beyond this bound, as we saw. It also disallows some correlations below the bound that are outside of the quantum convex set (for a discussion, see Pawłowski & Scarani, 2016). However there is numerical evidence that there exist correlations within the bound but outside of the quantum convex set that satisfy the information causality principle (Navascués et al., 2015). So it appears unlikely (though this was not known in 2009) that information causality can provide an answer to question (ii). It nevertheless remains promising as a principle with which to answer question (i) and can arguably still be thought of as a fundamental principle in that sense. Analogously, the fact that super-quantum no-signalling correlations are possible does not, in itself, undermine the status of no-signalling as a fundamental principle.

The information causality principle must be given some independent motivation if it is to play this explanatory role, however. Of course, the statement that the communication of  $m$  bits can yield no more than  $m$  bits of additional information to a receiver about a data set unknown to him is an intuitive one. But foundational principles of nature should require more for their motivation than such bare appeals to intuition. After all, quantum mechanics, which the principle aims to legitimate, arguably already violates many of our most basic intuitions. Pawłowski et al. (2009) unfortunately do not say very much to motivate information causality. But two ideas can be gleaned from statements made in their paper. The first is that in a world in which violations of information causality could occur, “certain tasks [would be] ‘too simple’” (p. 1101). The second is that in such a world there would be “implausible accessibility of remote data” (ibid.). The former idea has been expressed in this general context before. Van Dam (2013 [2005]), notably, shows that in a world in which PR-correlations exist and can be taken advantage of, only a trivial amount of communication is required to perform any distributed computational task. Van Dam argues (ibid., p. 12) that this is a reason to believe that such correlations cannot exist, for they violate the principle that “Nature does not allow a computational ‘free lunch’” (ibid., p. 9).<sup>9</sup> Bub (2012, pp. 180-181) echoes this thought by listing examples of distributed tasks (‘the dating game’ and ‘one-out-of-two’ oblivious transfer) which would become implausibly trivial if PR-correlated systems could be used.

Later in this paper I will argue that although such statements are unsatisfactorily vague, they nevertheless get at something that is importantly right; although they are right in, perhaps, a different sense than their authors envision. For now let me just say that even if one accepts van Dam’s argument that trivial communication complexity is implausible and should be ruled out—and that this should constitute a constraint on physical theory—not all correlations above the Tsirelson bound in fact result in trivial communication complexity. Brassard et al. (2006) have extended van Dam’s result by showing that (probabilistic) trivial communication complexity can be achieved for values of  $E > \sqrt{6}/3$ . But this still leaves a range of values for  $E$  open; physical correlations with associated values of  $E$  between the

---

<sup>9</sup>Cf. Aaronson (2005).

quantum mechanical maximum of  $1/\sqrt{2}$  and  $\sqrt{6}/3$  have not been shown to result in trivial communication complexity and cannot—at least not yet—be ruled out on those grounds. Thus the avoidance of trivial communication complexity cannot be used to motivate information causality in the way suggested by the statements of Pawłowski et al. (2009). In fairness, to say as they do that certain tasks would be ‘too simple’ in a world in which information causality is violated is not the same as saying that they would be trivial. The task remains, then, of expressing more precisely what is meant by ‘too simple’ in a way that is sufficient to motivate ruling out theories which violate the information causality principle in a less than maximal way (in particular with a value of  $E \leq \sqrt{6}/3$ ). We will return to this point later.

Regarding their second idea—that a world in which information causality is violated will lead to the “implausible accessibility of remote data” (p. 1101)—Pawłowski et al. again do not say enough,<sup>10</sup> although the idea is perhaps alluded to implicitly in another assertion they (too briefly) make, namely that information causality generalises the no-signalling principle (ibid., p. 1103). Fortunately the idea of implausible accessibility is expanded upon by Bub (2012), who motivates it in the following way:

the intuition is that if the correlations can be exploited to distribute one bit of communicated information among the  $N$  unknown bits in Alice’s data set, the amount of information distributed should be no more than  $\frac{1}{N}$  bits, because there can be no information about the bits in Alice’s data set in the previously established correlations themselves (p. 180).

Partly for this reason, Bub argues that the principle is misnamed. Drawing on the idea of implausible accessibility he argues that ‘information causality’ should rather be referred to as information *neutrality*: “The principle really has nothing to do with causality and is better understood as a *constraint on the ability of correlations to enhance the information content of communication in a distributed task*” (ibid., emphasis in original). Bub reformulates the principle as follows:

Correlations are informationally neutral: insofar as they can be exploited to allow Bob to distribute information communicated by Alice among the bits in an unknown data set held by Alice in such a way as to increase Bob’s ability to correctly guess an arbitrary bit in the data set, they cannot increase Bob’s information about the data set by more than the number of bits communicated by Alice to Bob (ibid.).

Stated in this way the principle sounds plausible and seems, intuitively, to be correct. However if the principle is to be of aid in ruling out classes of physical theory then it should be more than just intuitively plausible. If the goal of answering the question ‘Why the Tsirelson bound?’ is to give a convincing reason why correlations that are above the bound should be regarded as impossible, then if the fact that such correlations violate informational neutrality is to be one’s answer, one should give an independent motivation for why correlations must be informationally neutral. One might, for instance, motivate information neutrality by showing how it generalises or gives expression in some sense to a deeper

---

<sup>10</sup>Pawłowski & Scarani (2016, p. 429) do expand on the idea of implausible accessibility slightly: “we have transmitted only a single bit and the PR-boxes are supposed to be no-signalling so they cannot be used to transmit the other. Somehow the amount of information that the lab of Bob has is larger than the amount it received. Things like this should not happen.” I do not think this adds anything substantial to the idea expressed by Pawłowski et al. (2009) that such a situation is ‘implausible’.

underlying principle that is already well-motivated, or by pointing to ‘undesirable consequences’ of its failure. The consequence of a ‘free computational lunch’ given the existence of correlations above the bound, if it could be demonstrated, could (perhaps) constitute an example of the latter kind of motivation.

This said, there is a different way to think of the question ‘Why the Tsirelson bound?’ for which Bub’s explication of information causality in terms of informational neutrality is both a full answer and indeed an illuminating and useful one. In this sense the question represents a desire to understand what the Tsirelson bound expresses about correlations which violate it. Information neutrality answers this question by directing attention to a feature that no correlations above the bound can have. This feature, moreover, is one that we can easily grasp and explicitly connect operationally with our experience of correlated physical systems. On such a reading of the question, to answer ‘information neutrality’ is not of course to rule out that the world could contain non-informationally-neutral physical correlations. But on this view ruling out such a possibility is not the point, which is rather to provide a physically meaningful principle to help us to understand what our current physical theories, assuming they are to be believed, are telling us about the structure of the world.

In the remainder of this paper, however, I will continue to consider the information causality/neutrality principle as a possible answer in the first sense to the question ‘Why the Tsirelson bound?’. I will continue to consider, that is, whether there is some independent way of motivating the conclusion that correlations which violate the condition should be ruled out.

## 5 The ‘being-thus’ of spatially distant things

Our goal is to determine whether there is some sense in which we can motivate the idea that information causality must be satisfied by all physical theories which treat of correlated systems. I will now argue that some insight into this question can be gained if we consider the analogous question regarding no-signalling. As I mentioned earlier, the no-signalling condition (2.2) is not a relativistic constraint per se—in itself it is merely a restriction on the marginal probabilities associated with experiments on the subsystems of combined systems—but its violation entails the ability to instantaneously signal, which is in tension if not in outright violation of the constraints imposed by relativistic theory.<sup>11</sup> Indeed, the independently confirmed relativity theory can in this sense be thought of as an external motivation for thinking of the no-signalling principle as a constraint on the marginal probabilities allowable in any physical theory.

There is an arguably deeper way to motivate no-signalling, however, that can be drawn from the work of Einstein and which has been expanded upon by Demopoulos.<sup>12</sup> In the course of expressing his dissatisfaction with the ‘orthodox’ interpretation of quantum theory, Einstein described two foundational ideas—what Demopoulos calls *local realism* and *local action*. Realism in general, for Einstein, is a basic presupposition of any physical theory. It amounts to the claim that things in the world exist independently of our capability of knowing them; i.e.

the concepts of physics refer to a real external world, i.e., ideas are posited of things that claim a ‘real existence’ independent of the perceiving subject (bodies,

---

<sup>11</sup>For a discussion of signalling in the context of special and general relativity see Maudlin (2011, Ch. 4).

<sup>12</sup>This is done in Demopoulos’s monograph *On Theories*; see fn. 2.

fields, etc.), and these ideas are, on the other hand, brought into as secure a relationship as possible with sense impressions (Einstein 1948, as translated by Howard 1985, p. 187).

*Local realism*—alternately: the ‘mutually independent existence’ of spatially distant things—is the idea that things claim independent existence from one another insofar as at a given time they are located in different parts of space. Regarding this idea, Einstein writes:

Without such an assumption of the mutually independent existence (the ‘being thus’) of spatially distant things, an assumption which originates in everyday thought, physical thought in the sense familiar to us would not be possible (ibid.)

In the concrete context of a physical system made up of two correlated subsystems  $S_1$  and  $S_2$  (such as that described in the thought experiment of Einstein et al. 1935), local realism requires that

every statement regarding  $S_2$  which we are able to make on the basis of a complete measurement on  $S_1$  must also hold for the system  $S_2$  if, after all, no measurement whatsoever ensued on  $S_1$  (Einstein 1948, as translated by Howard 1985, p. 187).

In other words the value of a measurable theoretical parameter of  $S_2$  must not depend on whether a measurement is made on a system  $S_1$  that is located in some distant region of space. (And of course it must also not depend upon the *kind* of measurement performed on  $S_1$ ; cf. Howard 1985, p. 186.) Demopoulos notes that local realism as it is applied in such a context is a condition imposed on the measurable properties of the theory and hence it is a condition that is imposed at a theory’s ‘surface’ or operational level. This is an important point that I will return to later.

In the same *Dialectica* article Einstein also formulated a second principle:

For the relative independence of spatially distant things (A and B), this idea is characteristic: an external influence on A has no *immediate* effect on B; this is known as the ‘principle of local action’ ... The complete suspension of this basic principle would make impossible the idea of the existence of (quasi-) closed systems and, thereby, the establishment of empirically testable laws in the sense familiar to us (ibid., p. 188).

The thought expressed in the second part of this statement seems similar to Einstein’s earlier assertion that ‘physical thought’ would not be possible without the assumption of local realism. However Demopoulos convincingly argues that the principle of local realism, though it receives support from the principle of local action, is a conceptually more fundamental principle than the latter. For conceivably the principle of local realism—i.e. of ‘mutually independent existence’—could be assumed to hold, Demopoulos argues, even in the absence of local action. Indeed this is so in Newtonian mechanics. For example, Corollary VI to the laws of motion (Newton, 1999, p. 423) states that a system of bodies moving in any way whatsoever with respect to one another will continue to do so in the presence of equal accelerative forces acting on the system along parallel lines. This makes it

possible to treat the system of Jupiter and its moons, for example, as a quasi-closed system with respect to the sun. For owing to the sun's great distance, the forces exerted by it upon the Jovian system will be approximately equal and parallel. Thus despite its violation of local action, Demopoulos argues convincingly that Einstein would not (or anyway should not) have regarded a theory such as Newtonian mechanics as unphysical. It is still a basic *methodological* presupposition of Newtonian mechanics that spatially distant systems have their own individual 'being thus-ness', the description of which is made possible via the theory's characteristic methodological tool of successive approximation, in turn made possible by, for example, Corollary VI and the notion of quasi-closed system implied by it.<sup>13</sup>

Einstein's principle of local realism or mutually independent existence presupposes the framework of classical physics, which itself presupposes the framework of classical probability theory. Demopoulos argues, however, that the conceptual novelty of quantum theory consists in the fact that it is an 'irreducibly statistical theory', precisely in the sense that its probability assignments, unlike those described by classical probability theory, cannot in general be represented as weighted averages of two-valued measures over the Boolean algebra of all possible properties of a physical system (see also Pitowsky, 1989, 2006; Dickson, 2011). This raises the question of whether one can formulate a generalisation of the mutually independent existence condition that is appropriate for an irreducibly statistical theory such as quantum mechanics.<sup>14</sup>

Recall that Einstein's mutually independent existence condition, in the context of previously interacting but now spatially separated systems, is a condition that is imposed on the level of the measurable parameters of a theory and hence at its 'surface' or operational level. It requires, in particular, that the value of a measurable property of a system  $S_1$  in some region of physical space  $R_1$  is independent of what kind of measurement (or whether any measurement) is performed on some system  $S_2$  in a distant region of space  $R_2$ .

Demopoulos argues that in the context of an irreducibly statistical theory such as quantum mechanics, it is in fact the no-signalling condition which generalises the mutually independent existence condition. It does so in the sense that like mutually independent existence, no-signalling is a surface-level constraint on the local facts associated with a particular system, requiring that these facts be independent of the local surface-level facts associated with other spatially distant systems. Unlike the mutually independent existence condition, however, these local facts refer to the marginal probabilities associated with a system's measurable properties rather than with what one might regard as those properties themselves. Specifically, no-signalling asserts that the marginal probability associated with a measurement on a system  $S_1$  at a given location  $R_1$  is independent of what kind of measurement (or whether any measurement) is performed on some system  $S_2$  in a distant region of space  $R_2$ .<sup>15</sup> In this way no-signalling allows us to coherently treat systems in different regions of physical space as if they had mutually independent existences—i.e. as quasi-closed systems in the sense described above—and thus allows for the possibility of 'physical thought' in a methodological sense and for "the establishment of empirically testable laws in the sense familiar to us" (Einstein 1948, as translated by Howard 1985, p. 188). Demopoulos argues that quantum mechanics may in this way be thought of as a local

---

<sup>13</sup>For a detailed discussion of Newton's method of successive approximations and the methodological role therein played by Corollary VI, see Harper (2011). For a discussion of Corollary VI in the context of general relativity, see DiSalle (2006).

<sup>14</sup>I am not claiming here that Einstein himself would have been inclined to follow this line of reasoning.

<sup>15</sup>It is worth noting that the parameter independence condition (Shimony 1993) is just the no-signalling condition extended to include a hypothetical, possibly hidden, set of *underlying* parameters.

| $b_0$ | $a_2$ | $a_3$ | $a_2 \oplus a_3$ | $G$ |
|-------|-------|-------|------------------|-----|
| 0     | 0     | 0     | 0                | 0   |
| 0     | 0     | 1     | 1                | 0   |
| 0     | 1     | 0     | 1                | 1   |
| 0     | 1     | 1     | 0                | 1   |
| 1     | 0     | 0     | 0                | 0   |
| 1     | 0     | 1     | 1                | 1   |
| 1     | 1     | 0     | 1                | 0   |
| 1     | 1     | 1     | 0                | 1   |

**Figure 1:** A summary of the possible outcomes associated with Bob’s measurement  $G$  (his ‘guess’) in the guessing game of §3, based on Eq. (3.5). If all atomic variables are assumed to be equally likely to take on a value of 0 or 1, then  $G$  is probabilistically independent of Alice’s measurement setting  $a_2 \oplus a_3$ , but not of its components  $a_2$  and  $a_3$ , since, for example,  $p(G = 0|a_2 = 0) = 3/4 \neq p(G = 0|a_2 = 1)$ , and  $p(G = 0|a_3 = 0) = 3/4 \neq p(G = 0|a_3 = 1)$ .

theory of nonlocal correlations.

## 6 Mutually independent existence and communication

In the previous section we saw that no-signalling can be regarded as generalising a criterion for the possibility of ‘physical thought’ originally put forward by Einstein. Since quantum mechanics satisfies no-signalling, one may think of the theory, even under its orthodox interpretation, as in this sense legitimated methodologically by the principle. As we saw in §§2-4, however, other conceivable physical theories—some of which allow for stronger-than-quantum correlations—satisfy the condition as well. In light of this, ‘information causality’ (or ‘information neutrality’, in Bub’s terminology) was put forward by Pawłowski et al. (2009) as an additional foundational principle for more narrowly circumscribing the class of physically sensible theories. But in §4 I argued that the principle requires further motivation before it can legitimately be seen as playing this role. With our recent discussion of no-signalling in mind, let us now consider the proposal of Pawłowski et al. again.

*No-signalling* asserts that the marginal probabilities associated with Alice’s local measurements on a system  $S_A$  in a region  $R_A$  are independent of what kind of measurement (or whether any measurement) is performed by Bob locally on a system  $S_B$  in a distant region  $R_B$ . *Information causality* asserts that Bob can gain no more than  $m$  bits of information about Alice’s data set if she sends him only  $m$  bits. Pawłowski et al. (2009, p. 1101) remark that “The standard no-signalling condition is just information causality for  $m = 0$ ”. Bub (2012, p. 180) considers this remark to be misleading, but presumably all that Pawłowski et al. intend is that if Alice and Bob share *signalling* correlations, then Alice may provide Bob with information about her data set merely by measuring it, i.e. without actually sending him any bits. The information causality principle disallows this for any value of  $E$ , as does no-signalling.

On the other hand when (for instance)  $m = 1$ , then in the case where they have previously shared PR-correlated systems (i.e. systems such that  $E = 1$ ), one might argue that there arises a subtle sense in which the probabilities of Bob’s measurement outcomes can be influenced by Alice’s remote measurement settings. Consider the outcome of Bob’s combined measurement  $G =_{df} c \oplus B_{III} \oplus B_{II}$ , i.e. his ‘guess’ (3.4). From (3.5) it would appear

that Bob’s outcome is in part determined by the setting of Alice’s measurement on system **II**,  $a_2 \oplus a_3$ , since this appears explicitly in the equation. However in this case appearances are misleading, for the reader can verify that  $G$  is probabilistically independent of  $a_2 \oplus a_3$  (see figure 1).  $G$  is nevertheless probabilistically dependent on both of  $a_2$  and  $a_3$  considered individually. So one might say that although the outcome of  $G$  is not influenced by any of Alice’s measurement settings *per se*, it does seem to be influenced by the particular way in which those settings have been determined (despite the fact that neither  $a_2$  nor  $a_3$  are directly used by Alice to determine the value of the bit that she sends to Bob,  $c$ ). Put a different way, the constituents of Alice’s measurement setting on system **II** respectively determine the two possible outcomes of Bob’s guess whenever he performs the measurement  $G$  (for a given  $b_0$ ). Likewise in the case where Bob measures  $G' = c \oplus B_{III} \oplus B_I$  (i.e. his guess (3.2)); the two possible outcomes of  $G'$  are, respectively, determined by the constituents of Alice’s measurement settings on system **I**,  $a_0$  and  $a_1$  (for a given  $b_0$ ).

Note that since  $a_2$  and  $a_3$  (respectively:  $a_0$  and  $a_1$ ), besides being the constituents of Alice’s measurement settings on **II** (respectively: **I**), are also in fact the values of bits in Alice’s list **a**, the above considerations resonate with Bub’s remark (quoted above) that stronger-than-quantum correlations are such that they may themselves include information about Alice’s data set in the context of a game like that described in §3. These considerations further suggest a sense, *pace* Bub, in which it could be argued that the name ‘information causality’ is indeed apt. For the bit of information  $c$  that Alice sends to Bob can be thought of as the ‘enabler’ or ‘cause’, at least in a metaphorical sense, of Bob’s ability to use this aspect of the correlations to his advantage (cf. Pawłowski & Scarani, 2016, §3.4).<sup>16</sup>

Now we saw in the last section that Einstein’s principle of mutually independent existence can be thought of as a methodological principle such that it (arguably) must be presupposed if ‘physical thought’ is to be possible. We saw that this principle holds in the context of Newtonian mechanics, which one may think of as in that sense a local theory of nonlocal forces. We also saw that a generalisation of mutually independent existence appropriate for an irreducibly statistical theory—i.e. the no-signalling principle—holds in the context of quantum mechanics, and that it may thus be thought of analogously as a local theory of nonlocal correlations.

The context of our current investigation is one which involves considering communicating agents capable of building and manipulating physical systems—thought of now as resources—for their own particular purposes. Our context, that is, is the ‘practical’ one associated with quantum computation and information theory, recently described by Cuffaro (2017, forthcominga). As Cuffaro has argued, this context of investigation is in fact distinct from the more familiar ‘theoretical’ context that is associated with more traditional foundational investigations of quantum mechanics. A different way of putting this is that quantum computation and information theory, like the science of thermodynamics, is a ‘resource’ or ‘control’ theory (Myrvold, 2011; Wallace, 2014; Ladyman, forthcoming).

Without the methodological presupposition of mutually independent existence—according to which systems that occupy distinct regions of space are to be regarded as existing independently of one another—the idea of a quasi-closed system that can be subjected to empirical test, and in this sense ‘physical thought’, would (arguably) not be possible. Analogously, one might argue that in the context of a theory of communication—i.e. of the various resource costs associated with different communicational protocols and their interrelations—that it is necessary to presuppose that some significant operational distinction can be made between the parties involved in a communicational protocol. One might

---

<sup>16</sup>Perhaps, though, a better name would be the ‘no information causality’ principle.

argue, that is, that it is constitutive of the very idea of communication that it is an activity that takes place between what can be effectively regarded as two mutually independently existing entities for the purposes of quantifying the complexity of a particular protocol.<sup>17</sup>

Without the ability to make such an effective distinction between the sender and the receiver of information, it is not at all obvious how one should begin to quantify the amount of information that is required to be sent *from* Alice *to* Bob in the context of a particular protocol. Arguably this is the situation one is confronted with in the context of the game described above when it is played with super-quantum correlated systems. For the correlations, shared prior to the beginning of the game, upon being ‘activated’ by Alice’s classical message  $c$  to Bob, can be said to contribute something over and above  $c$  to the information Bob then gains about Alice’s data set. Indeed when Alice and Bob share *maximally* super-quantum systems (i.e. PR-systems), then after receiving  $c$  there is a sense in which Alice and her system can be said to be ‘a part’ of Bob and his system in the context of the game being played. Indeed from this point of view the fact that the complexity associated with any distributed computational task is trivial when PR-correlations are used seems natural. For after receiving  $c$  Bob has *immediate* access to the value of any single bit of Alice’s that he would like.

One might object that something similar could be said in the case where Alice and Bob share an entangled quantum system. Could not one say, that is, that Alice and Bob become likewise inseparable or ‘nonlocally joined’ (cf. Howard, 1989) upon sharing such a system in the context of the information causality game? And yet no one imagines the very possibility of the sciences of quantum information theory and quantum communication complexity to have been undermined as a result. This objection, however, appeals to a characterisation of the situation involving the sharing of an entangled quantum system that pertains to the metaphysical level of description. It therefore does not undermine the considerations of the previous paragraph. Consider the description of a classical bipartite communicational protocol. Both before and after communication has taken place, such a description may be regarded as decomposable into three parts: a sender, a receiver, and something communicated between them. In the case of a quantum protocol which utilises entangled states the situation regarding the possibility of this decomposition is more complex as a result of the well-known conceptual intricacies associated with entangled quantum states. However whether or not Alice and her system, and Bob and his system, are ‘in reality’ inseparably entangled with one another, it remains the case, both before (because of quantum mechanics’ satisfaction of the no-signalling condition) and after the communication of a classical message (because of quantum mechanics’ satisfaction of the information causality condition), that Alice, Bob, and the message  $c$  may be operationally distinguished from one another in the sense that Bob cannot take advantage of the underlying connection he has with Alice and her system via the correlations he shares with her to gain information about her data set over and above what has been provided to him via  $c$ . It is true that previously shared quantum correlations enable one to communicate with greater efficiency than is possible using only previously shared classical correlations. As (4.3) shows, Bob has a higher probability of guessing correctly in the information causality game if he and Alice have previously shared quantum as opposed to classical correlations.<sup>18</sup> And the question arises regarding the source of this increased communicational power. But whatever that source is, it is not

---

<sup>17</sup>Cf. Hushilevitz & Nisan (1997, p. x). Cf. also Maroney & Timpson’s (forthcoming) emphasis on the initialisation and readout stages of an information processing task.

<sup>18</sup>This is true in other contexts besides that of the information causality game. See, e.g., Buhrman et al. (2001); Brukner et al. (2002, 2004).



the case that it manifests itself in nonlocality or nonseparability at the *operational* level.<sup>19</sup>

But the game described by Pawłowski et al. (2009) involves the communication of *classical* bits from Alice to Bob. Might not this limitation in Bob's ability to take advantage of his underlying connection with Alice be overcome if we allow her to send him qubits rather than only classical bits? Indeed, it is well known that if Alice sends a qubit to Bob that is entangled with a qubit that is already in his possession, then Alice and Bob can implement the 'superdense coding' protocol (Nielsen & Chuang, 2000, §2.3); Alice's sending of a single qubit to Bob according to this protocol will allow him to learn two bits' worth of classical information.<sup>20</sup> Does this not undermine the claim that quantum correlations contribute nothing over and above whatever message is sent between Alice and Bob to the information gained by him?

It does not. On the one hand, before the transmission of the qubit(s) from Alice to Bob, no-signalling implies that Alice and Bob can be considered as operationally separable despite their sharing an entangled system, as we have seen above. On the other hand, in the superdense coding protocol, after Alice transmits her message to Bob, all of the correlated quantum system that was initially shared is now in Bob's possession. So after transmission there is no sense in which Bob can take advantage of correlations shared with Alice at that time. In a sense Alice's message to Bob 'just is' information regarding the correlations that exist between them at the time at which she sends it.<sup>21</sup>

As we have seen, when Alice and Bob share PR-correlated systems, they can win a round with certainty in the  $m = 1$  game for any  $N$  by exchanging a single classical bit. Earlier I also mentioned van Dam's (2013 [2005]) result to the effect that PR-correlated systems allow one to perform *any* distributed computational task with only a trivial amount of communication. These results are striking. However the reader may nevertheless feel somewhat suspicious of them for the following reason: the number of PR-correlated systems required to implement these protocols, as we have seen, is great. With respect to the length  $n$  of Bob's bit string  $\mathbf{b}$  (arguably the most appropriate measure of input size for the game), implementing the solution described above requires that they share  $2^n - 1$  PR-systems; i.e. the number of PR-systems required grows exponentially with the input size. Likewise for van Dam's protocol.<sup>22</sup> A reduction in *communication* complexity has therefore been achieved only at the expense of an increase in *computational* complexity. One might argue that it is in this sense misleading to consider the complexity of implementing the protocol with PR-correlated systems to be trivial—that they provide us with a 'free lunch'.

I will return to this point later. But for now let me say that, arguably, this is not a relevant consideration in this context. The theories of communication complexity and computational complexity are distinct sub-disciplines of computer science. The goal of communication complexity is to quantify the amount of communication necessary to implement

---

<sup>19</sup>Compare this with Buhrman et al. (2001), who writes that entanglement enables one to "*circumvent* (rather than simulate) communication" (p. 1831, emphasis in original), and also with Bub (2010)'s discussion of entanglement in the context of quantum computation, which he argues allows a quantum computer to compute a global property of a function by performing fewer, not more, computations than classical computers.

<sup>20</sup>In the context of a suitably generalised version of the information causality game, it turns out that a two-bit information gain per qubit constitutes an upper bound (Pitalúa-García, 2013).

<sup>21</sup>This conclusion is essentially that of Spekkens (2007, p. 032110-20). Fascinatingly, Spekkens also shows that the superdense coding protocol can be implemented in his toy classical theory.

<sup>22</sup>Specifically, van Dam's (2013 [2005]) protocol requires a number of systems that can grow exponentially with respect to the input size of an instance of the Inner Product problem, after which the solution can be efficiently converted into a solution to any other distributed computational problem.

various communicational protocols. For this purpose one abstracts away from any consideration of how complicated a computational system must be in other respects (Hushilevitz & Nisan, 1997). The question addressed in van Dam (2013 [2005]) and in Pawłowski et al. (2009); Pawłowski & Scarani (2016) concerns whether the availability of PR-correlated systems would make communicational, not computational, complexity theory superfluous. From this point of view any previously prepared PR-correlated systems are viewed as ‘free resources’ for the purposes of the analysis.

This said, one could imagine that the subsystems of PR-correlated systems employ some hidden means of communication with one another, and argue that this must be included in the complexity one ascribes to the protocol. To do so, however, would constitute a descent below the empirically verifiable level. Obviously this is not in itself objectionable. But it is hard to see what use this would be to a theory of communicational complexity, which after all, like computational complexity (Cuffaro, forthcomingb), aims to be a practical science whose goal is to guide us in making distinctions in practice between real problems related to data transmission that are of varying levels of difficulty. In this sense appealing to an underlying communication that takes place between the subsystems of PR-systems does not help with the conclusion that complexity theory, at least in an operational sense, becomes superfluous if PR-correlated systems are available. The objection described in the previous paragraph is nevertheless an important one that I will return to.

Above I have motivated the idea, of Pawłowski et al. (2009, p. 1101), that the kind of accessibility of remote data that is possible given the existence of super-quantum correlated systems is “implausible”. I have done so by describing, *pace* Bub (2012), the sense in which information causality can be taken to generalise no-signalling. In so doing I have gestured at a connection between the idea of implausible accessibility and the *prima facie* separate idea that a world in which super-quantum correlated systems exist would be ‘too good to be true’ in a communicational complexity-theoretic sense. My arguments have been mainly conceptual. I have argued, that is, that a kind of conceptual ambiguity at the operational level between the parties to a communicational protocol may result if stronger-than-quantum correlations are available to use. When the stronger-than-quantum correlations are strong enough (i.e. when  $E > \sqrt{6}/3$ ), this results in the trivial communicational complexity of any distributed computational task. But trivial communicational complexity does not result, or anyway has not yet been shown to result, for values of  $E$  above the Tsirelson bounded value of  $1/\sqrt{2}$  that are below  $\sqrt{6}/3$ . This is despite the fact that the conceptual ambiguity I have described is present for all such values of  $E$ .

Thus one may wonder whether ‘a little’ ambiguity may be tolerable for practical purposes—whether, that is, a theory which admits correlations which only ‘weakly’ violate the Tsirelson bound should be admitted within the space of possible physical theories from the point of view of the principle. The situation could be seen as analogous to the situation one is faced with in Newtonian Mechanics, in fact, for Corollary VI (which I described in §5) only guarantees that a given system can be treated as (quasi-) closed when it is acted upon by forces which are *exactly* equal in magnitude and *exactly* parallel. Clearly this is not the case for the Jovian system *vis-à-vis* the sun, for example. Corollary VI nevertheless functions as a methodological tool in that it allows us to maintain the idea of the mutually independent existence of spatially distant things as a methodological principle and treat the Jovian system, for the practical purpose of analysing its internal motions, as unaffected by the forces exerted upon it by the sun.

There is much work to be done before information causality can be considered as successful in ruling out—in the conceptual sense described in the previous paragraph—all

theories whose correlations violate the Tsirelson bound. Irrespective of whether this goal can be achieved, however, this does not (necessarily) undermine the status of information causality motivated as a methodological principle in something like the way that I have done in this paper. In particular, information causality would be especially compelling if one could draw a relation between the degree of violation of the principle and the degree of ‘superfluousness’ of the resulting theory of communication complexity with an eye to distinguishing ‘weak’ violations of the Tsirelson bound from more objectionable violations. Thus there is much work to do in any case.

I will close with the following more fundamental objection. Why should nature care whether beings such as us are able to engage in communication complexity theory? In fact there is no fundamental reason why nature should care. Analogously, there is no fundamental reason why nature should care whether beings such as us can do physics. But the goal of empirical science is not to derive the structure of the world or its constituent entities by way of a priori or ‘self-evident’ principles. It is rather to make sense of and explain our experience of and in the world. In fact we have a science which is called physics. And in fact we have a science which we refer to as communicational complexity theory. The principle of mutually independent existence, and analogously the principle of information causality, may be thought of as answers to the question: ‘how are such facts possible?’ (cf. Kant, 1998 [1781], pp. B20-B21).

That said, these may not be definitive answers. The necessity of presupposing Einstein’s mutual independence and local action principles for the purposes of theory testing has been questioned by Howard (1989). In a similar way, one might argue that it is misleading to think that the existence of correlated systems which ‘strongly’ violate the Tsirelson bound makes a science of communication complexity impossible. Rather, one might conclude instead that the idea of a science of communication complexity that is wholly independent of *computational* complexity-theoretic considerations is unachievable. This, one might argue, is the real lesson of the fact that an exponential number of PR-correlated systems is required to implement Alice’s and Bob’s solution to their guessing game. Yet even if this were all that we learned from information causality, it would still represent a significant advance in our understanding of the structure of our theoretical knowledge—an understanding of the physically motivated constraints under which two mathematical *theories* may be regarded as mutually independent.

## 7 Summary

Above I have argued that the principle of information causality has not yet been sufficiently motivated to play the role of a foundational principle of nature, and I have described a way in which one might begin to provide it with such a motivation. More specifically I described an argument, due to Demopoulos, to the effect that the no-signalling condition can be viewed as a generalisation, appropriate to an irreducibly statistical theory, of Einstein’s principle of mutually independent existence interpreted methodologically. I then argued that information causality can in turn be motivated as a further generalisation of no-signalling that is appropriate to a theory of communication. I closed by describing a number of important obstacles that are required to be overcome if the project of establishing information causality as a foundational principle is to succeed.

## References

- Aaronson, S. (2005). NP-complete problems and physical reality. *ACM SIGACT News*, 36, 30–52.
- Brassard, G., Buhrman, H., Linden, N., Méthot, A. A., Tapp, A., & Unger, F. (2006). Limit on nonlocality in any world in which communication complexity is not trivial. *Physical Review Letters*, 96, 250401–1–4.
- Brukner, Č., Žukowski, M., Pan, J.-W., & Zeilinger, A. (2004). Bell’s inequalities and quantum communication complexity. *Physical Review Letters*, 92, 127901.
- Brukner, Č., Žukowski, M., & Zeilinger, A. (2002). Quantum communication complexity protocol with two entangled qutrits. *Physical Review Letters*, 89, 197901.
- Bub, J. (2010). Quantum computation: Where does the speed-up come from? In A. Bokulich, & G. Jaeger (Eds.) *Philosophy of Quantum Information and Entanglement*, (pp. 231–246). Cambridge: Cambridge University Press.
- Bub, J. (2012). Why the Tsirelson bound? In Y. Ben-Menahem, & M. Hemmo (Eds.) *Probability in Physics*, (pp. 167–185). Berlin: Springer.
- Bub, J. (2016). *Bananaworld, Quantum Mechanics for Primates*. Oxford: Oxford University Press.
- Buhrman, H., Cleve, R., & van Dam, W. (2001). Quantum entanglement and communication complexity. *SIAM Journal on Computing*, 30, 1829–1841.
- Cirel’son, B. S. (1980). Quantum generalizations of Bell’s inequality. *Letters in Mathematical Physics*, 4, 93–100.
- Clauser, J. F., Horne, M. A., Shimony, A., & Holt, R. A. (1969). Proposed experiment to test local hidden-variable theories. *Physical Review Letters*, 23, 880–884.
- Clifton, R., Bub, J., & Halvorson, H. (2003). Characterizing quantum theory in terms of information-theoretic constraints. *Foundations of Physics*, 33, 1561–1591.
- Cuffaro, M. E. (2017). On the significance of the Gottesman-Knill theorem. *The British Journal for the Philosophy of Science*, 68, 91–121.
- Cuffaro, M. E. (forthcominga). Reconsidering no-go-theorems from a practical perspective. *The British Journal for the Philosophy of Science*.
- Cuffaro, M. E. (forthcomingb). Universality, invariance, and the foundations of computational complexity in the light of the quantum computer. In S. O. Hansson (Ed.) *Technology and Mathematics: Philosophical and Historical Investigations*. Berlin.
- Demopoulos, W. (In Preparation). *On Theories*.
- Dickson, M. (2011). Aspects of probability in quantum theory. In C. Beisbart, & S. Hartmann (Eds.) *Probabilities in Physics*, (pp. 171–200). Oxford: Oxford University Press.
- DiSalle, R. (2006). *Understanding Space-time*. Cambridge: Cambridge University Press.

- Einstein, A. (1948). Quanten-mechanik und wirklichkeit. *Dialectica*, 2, 320–324.
- Einstein, A., Podolsky, B., & Rosen, N. (1935). Can quantum-mechanical description of physical reality be considered complete? *Physical Review*, 47, 777–780.
- Felline, L. (2016). It’s a matter of principle: Scientific explanation in information-theoretic reconstructions of quantum theory. *Dialectica*, 70, 549–575.
- Feynman, R. P., Leighton, R. B., & Sands, M. (1964). *The Feynman Lectures on Physics*. Reading, Mass.: Addison-Wesley.
- Harper, W. L. (2011). *Isaac Newton’s Scientific Method*. Oxford: Oxford University Press.
- Howard, D. (1985). Einstein on locality and separability. *Studies in History and Philosophy of Science*, 16, 171–201.
- Howard, D. (1989). Holism, separability, and the metaphysical implications of the Bell experiments. In J. T. Cushing, & E. McMullin (Eds.) *Philosophical Consequences of Quantum Theory*, (pp. 224–253). Notre Dame: University of Notre Dame Press.
- Hushilevitz, E., & Nisan, N. (1997). *Communication Complexity*. Cambridge: Cambridge University Press.
- Kant, I. (1998 [1781]). *Kritik der Reinen Vernunft*. Hamburg: Felix Meiner.
- Koberinski, A., & Müller, M. P. (forthcoming). Quantum theory as a principle theory: Insights from an information theoretic reconstruction. In M. E. Cuffaro, & S. C. Fletcher (Eds.) *Physical Perspectives on Computation, Computational Perspectives on Physics*. Cambridge: Cambridge University Press.
- Ladyman, J. (forthcoming). Intension in the physics of computation: Lessons from the debate about Landauer’s principle. In M. E. Cuffaro, & S. C. Fletcher (Eds.) *Physical Perspectives on Computation, Computational Perspectives on Physics*. Cambridge: Cambridge University Press.
- Maroney, O. J. E., & Timpson, C. G. (forthcoming). How is there a physics of information? on characterising physical evolution as information processing. In M. E. Cuffaro, & S. C. Fletcher (Eds.) *Physical Perspectives on Computation, Computational Perspectives on Physics*. Cambridge: Cambridge University Press.
- Maudlin, T. (2011). *Quantum Non-Locality and Relativity*. Cambridge, MA: Wiley-Blackwell, third ed.
- Myrvold, W. (2011). Statistical mechanics and thermodynamics: A Maxwellian view. *Studies in History and Philosophy of Modern Physics*, 42, 237–243.
- Myrvold, W. C. (2010). From physics to information theory and back. In A. Bokulich, & G. Jaeger (Eds.) *Philosophy of Quantum Information and Entanglement*, (pp. 181–207). Cambridge: Cambridge University Press.
- Navascués, M., Guryanova, Y., Hoban, M. J., & Acín, A. (2015). Almost quantum correlations. *Nature Communications*, 6, 1–7.

- Newton, I. (1999). Mathematical principles of natural philosophy. Trans. I. B. Cohen, & A. Whitman. In I. B. Cohen (Ed.) *The Principia: A New Translation and Guide*, (pp. 371–946). Berkely and Los Angeles, California: University of California Press.
- Nielsen, M. A., & Chuang, I. L. (2000). *Quantum Computation and Quantum Information*. Cambridge: Cambridge University Press.
- Pawłowski, M., Paterek, T., Kaszlikowski, D., Scarani, V., Winter, A., & Żukowski, M. (2009). Information causality as a physical principle. *Nature*, *461*, 1101–1104.
- Pawłowski, M., & Scarani, V. (2016). Information causality. In G. Chiribella, & R. W. Spekkens (Eds.) *Quantum Theory: Informational Foundations and Foils*, (pp. 423–438). Dordrecht: Springer.
- Pitalúa-García, D. (2013). Quantum information causality. *Physical Review Letters*, *110*, 210402.
- Pitowsky, I. (1989). *Quantum Probability — Quantum Logic*. Hemsbach: Springer.
- Pitowsky, I. (2006). Quantum mechanics as a theory of probability. In W. Demopoulos, & I. Pitowsky (Eds.) *Physical Theory and its Interpretation*, (pp. 213–240). Dordrecht: Springer.
- Popescu, S., & Rohrlich, D. (1994). Quantum nonlocality as an axiom. *Foundations of Physics*, *24*, 379–385.
- Schrödinger, E. (1935). Discussion of probability relations between separated systems. *Mathematical Proceedings of the Cambridge Philosophical Society*, *31*, 555–563.
- Shimony, A. (1993). *Search for a Naturalistic Worldview, Volume 2: Natural Science and Metaphysics*. Cambridge: Cambridge University Press.
- Spekkens, R. W. (2007). Evidence for the epistemic view of quantum states: A toy theory. *Physical Review A*, *75*, 032110.
- van Dam, W. (2013 [2005]). Implausible consequences of superstrong nonlocality. *Natural Computing*, *12*, 9–12. arXiv:quant-ph/0501159v1.
- Wallace, D. (2014). Thermodynamics as control theory. *Entropy*, *16*, 699–725.