

# A Weyl-Type Theorem for Geometrized Newtonian Gravity

Erik Curiel

October 7, 2015

I state and prove, in the context of a space having only the metrical structure imposed by the geometrized version of Newtonian gravitational theory, a theorem analagous to that of Weyl's in a Lorentzian space. The theorem, loosely speaking, says that a projective structure and a suitably defined compatible conformal structure on such a space jointly suffice for fixing the metrical structure of a Newtonian spacetime model up to constant factors. It allows one to give a natural, physically compelling interpretation of the spatiotemporal geometry of a geometrized Newtonian gravity spacetime manifold, in close analogy with the way Weyl's Theorem allows one to do in general relativity.

## Contents

<b>1</b>	<b>Weyl's Theorem</b>	<b>1</b>
<b>2</b>	<b>Newtonian Conformal Structure</b>	<b>2</b>
<b>3</b>	<b>The Theorem</b>	<b>6</b>
<b>A</b>	<b>Appendix: Geometrized Newtonian Gravity</b>	<b>7</b>

## 1 Weyl's Theorem

Soon after Einstein first promulgated the theory of general relativity, [Weyl \(1918\)](#) formulated and proved a theorem that has since served as the foundation for one of the most influential and compelling ways to give a physical interpretation to the theory's mathematical machinery, Lorentzian geometry:<sup>1</sup>

---

<sup>1</sup>Researchers as varied in their backgrounds, aims and temperaments as [Eddington \(1923\)](#), [Møller \(1952\)](#), [Trautman \(1965\)](#), [Hawking and Ellis \(1973\)](#) and [Malament \(2012\)](#) have used it so, to mention only a small sample. This form of interpretation can be enlighteningly contrasted with one based, *e.g.*, on chronometry, as in [Einstein \(2001\)](#) or [Synge \(1960\)](#).

**Theorem 1.1** *Given a conformal structure on a differential manifold, and a projective structure agreeing with it on (images) of its null geodesics, there is a Lorentz metric, fixed up to a constant factor, having each as its associated structure of that kind.*

In essence, to know the conformal structure is to know which curves are timelike (the possible paths of massive bodies), which are null geodesics (the possible paths of light-rays *in vacuo*), and which are spacelike (the possible paths of no physical system); to know the projective structure allows one further to say which timelike curves are (images of) geodesics, *i.e.*, the possible paths of freely falling massive bodies. To know both, then, the theorem tells us, is to know the spacetime metric up to a fixed constant, *i.e.*, up to the choosing of a system of units of measurement, thus giving us the physical significance of the metric: it is determined by the behavior of light rays and freely falling bodies.<sup>2</sup> In this paper, I construct the necessary machinery for a formulation and proof of a natural analogue of this theorem in the context of geometrized Newtonian gravity. It provides in the same way the basis for a compelling physical interpretation of the spatiotemporal structures of that theory.

In §2, I characterize the analogue of a conformal structure in Newtonian spaces, *Newtonian conformal structure*, which comprises most of the heavy lifting. I conclude in §3 with a statement and proof of the theorem, and an explanation of the way it grounds a physical interpretation of the theory. In appendix A, I sketch the machinery of geometrized Newtonian gravity required for the paper’s definitions and proofs, following the treatment of Malament (2012, ch. 4, §2) (with minor emendations and simplifications).

## 2 Newtonian Conformal Structure

The following definition encapsulates almost the entirety of the formal structure of geometrized Newtonian gravity as a physical theory, the remainder being elaboration of and derivations from its elements.

**Definition 2.1** *A Newtonian spacetime model is an ordered quintuplet  $(\mathcal{M}, \rho, h^{ab}, t_{ab}, \nabla_a)$ , such that:*

1.  $\mathcal{M}$  is a four-dimensional, connected, paracompact, smooth, differential manifold
2.  $\rho$  is a smooth, non-negative, scalar function on  $\mathcal{M}$
3.  $h^{ab}$  is a smooth, symmetric tensor field on  $\mathcal{M}$  of signature  $(0, 1, 1, 1)$
4.  $t_{ab}$  is a smooth, symmetric tensor field on  $\mathcal{M}$  of signature  $(1, 0, 0, 0)$
5.  $t_{ab}$  and  $h^{ab}$  are compatible, *i.e.*,  $t_{an}h^{nb} = 0$

---

<sup>2</sup>See, *e.g.*, Ehlers, Pirani, and Schild (1972) for a thorough exposition of the mathematics behind the theorem and its intended physical significance, and Malament (2012, ch. 2, §1, pp. 120–121) for a lapidary account of such a physical interpretation, including a discussion of its virtues and possible problems.

6.  $\nabla_a$  is a smooth derivative operator on  $\mathcal{M}$ , compatible with  $t_{ab}$  and  $h^{ab}$  in the sense that  $\nabla_a t_{bc} = 0$  and  $\nabla_a h^{bc} = 0$

$\mathcal{M}$  represents spacetime, the “totality of all point-events”, and  $\nabla_a$  the physically relevant affine structure, *i.e.*, the one whose geodesics represent unaccelerated paths in the spacetime.  $\rho$  represents the mass-density distribution of matter.  $h^{ab}$  and  $t_{ab}$  represent, respectively, the closest we come to having spatial and temporal metric structures on  $\mathcal{M}$ . (See appendix A for an explanation of the sense in which these tensors represent such structure.) We refer to the ordered pair  $(h^{ab}, t_{ab})$  as a *Newtonian metrical structure*.

From hereon, we assume all Newtonian spacetime models to be spatially flat (*i.e.*,  $R^{abcd}$ , the spatialized Riemann tensor, vanishes; see appendix A for an explanation of the significance of this condition). It follows from the results and discussion of Malament (1986) that this represents no real loss of generality, as it is those models that best capture the idea of the possible spaces of classical Newtonian gravitational theory. We also assume in what follows that  $\mathcal{M}$  is simply connected and the spacetime model is temporally orientable (*i.e.*, that there exists a globally defined temporal function  $t$  such that  $t_{ab} = \nabla_a t \nabla_b t$ ). Again, this is no real loss of generality, for all arguments and conclusions would still go through without the assumption, at the cost of constant hedging about which results are local and which global, and hence much technical work of a nit-picky sort without any counter-balancing gain in physical insight.

From hereon, we will need to keep track of the difference between a curve considered, on the one hand, as a smooth, injective mapping from a real interval to  $\mathcal{M}$ , and, on the other, as the point-set image in  $\mathcal{M}$  of such a mapping. I will use ‘curve’ when I mean the former, and ‘image of a curve’ when I mean the latter.

Two affinities  $\nabla_a$  and  $\tilde{\nabla}_a$  are projectively equivalent if they agree on images of geodesics, which is to say, if they agree on geodesics up to arbitrary (smooth, monotonic) reparametrization:  $\xi^n \nabla_n \xi^a = 0$  if and only if  $\xi^n \tilde{\nabla}_n \xi^a = \lambda \xi^a$  for  $\lambda$  a smooth function on  $\xi$ . A projective structure, then, is a maximal collection of affine structures on a Newtonian spacetime all of which agree on images of all geodesics (“maximal” in the sense that we throw in every affine structure that meets the criterion). One can equally well define the projective structure as the complete family of images of geodesics on which the affine structures agree. A member of such a family of images of geodesics is a projective geodesic.

Now, to characterize the analogue of conformal structure in geometrized Newtonian gravity. In general relativity, one can define a conformal structure to be an assignment of a smoothly varying field of quadratic cones at every point of the spacetime manifold, the null-cones. Two Lorentz metrics  $g_{ab}$  and  $\tilde{g}_{ab}$  are conformally equivalent if they have the same null-cone structure. This holds if and only if the two metrics agree on their null geodesics, which holds if and only if  $g_{ab} = \Omega^2 \tilde{g}_{ab}$  for some smooth, non-zero scalar field  $\Omega$ . Because we have no such metric structure in geometrized Newtonian gravity, and correlatively no cone-structure (except the degenerate one, which will not do) or non-trivially null type of vector, one cannot apply such a characterization of conformal structure here.

The following observation provides the required clue for moving forward: two Lorentz metrics are conformally equivalent if and only if they agree on orthogonality-relations for all pairs of vectors. Indeed, two Lorentz metrics  $g_{ab}$  and  $\tilde{g}_{ab}$  agree on orthogonality-relations if and only if  $g_{ab} = \Omega^2 \tilde{g}_{ab}$  for some smooth, non-zero scalar field  $\Omega$ . Equivalently,  $g_{ab}$  and  $\tilde{g}_{ab}$  agree in assignments of ratios of lengths to any pair of tangent vectors. The conformal structure so characterized then allows one to distinguish among timelike, null and spacelike vectors, to distinguish null geodesics, and to reconstruct the null-cone structure. Null-vectors are those non-zero vectors orthogonal to themselves, which picks out the null-cones; and a null geodesic is one contained in a null-cone (in an appropriate sense—see [Ehlers, Pirani, and Schild 1972](#)). Timelike vectors are those pointing into the interior of the null cones, and spacelike vectors are all the rest.

This suggests that we attempt to characterize conformal structure in the context of geometrized Newtonian gravity by making use of orthogonality among (appropriate) pairs of vectors.

**Definition 2.2** *A Newtonian Conformal Structure  $\mathfrak{C}$  on a four-dimensional manifold  $\mathcal{M}$  consists of:*

1. *at each point  $p \in \mathcal{M}$ , a three-dimensional vector subspace  $\mathfrak{C}_p$  of  $T_p^*M$  (the tensor space of 1-forms over  $p$ ), smoothly varying from point to point*
2. *orthogonality relations fixed for all pairs of elements of  $\mathfrak{C}_p$*

This structure suffices for defining a family of symmetric tensor fields characterized by a given  $h^{ab}$  with signature  $(0, 1, 1, 1)$ , fixed up to multiplication by a positive scalar field  $\Omega^2$ : two 1-forms  $\alpha_a, \beta_a \in \mathfrak{C}_p$  are orthogonal if and only if  $h^{mn} \alpha_m \beta_n$ . We can now distinguish timelike from spacelike vectors: spacelike vectors are those that result from raising an index of a 1-form in a  $\mathfrak{C}_p$ ; timelike are those that cannot be so derived. If  $\xi^a = h^{an} \alpha_n$ , we say  $\alpha_a$  is a spacelike representative 1-form of  $x^a$ . The orthogonality relation extends in a natural way to pairs consisting of spacelike vectors, and pairs consisting of one spacelike and one timelike vector. We can now also determine a second family of symmetric tensor fields, characterized by a given  $t_{ab}$  with signature  $(1, 0, 0, 0)$ , fixed up to multiplication by a positive scalar field  $\chi^2$ . A vector  $\xi^a$  is timelike if and only if  $t_{mn} \xi^m \xi^n > 0$ . It follows that the timelike vectors at a point form a four-dimensional affine space modeled on the space of all tangent vectors at that point. Clearly, any two such representative tensor fields  $h^{ab}$  and  $t_{ab}$  are compatible, in the sense that  $t_{an} h^{nb} = 0$ . We say two symmetric tensor fields  $h^{ab}$  and  $\tilde{h}^{ab}$  of the appropriate signature are *conformally equivalent* if they live in the same family of tensor fields of a given Newtonian conformal structure, and similarly for  $t_{ab}$  and  $\tilde{t}_{ab}$ . A conformal derivative operator, then, is the family of derivative operators each of which is compatible with a pair of compatible spatial and temporal metrics in the conformal family, in the sense that each triplet of representatives satisfy  $\nabla_a h^{bc} = 0$  and  $\nabla_a t_{bc} = 0$ .

Because, given two spacelike vectors, there is always a third that, added to the first, makes it orthogonal to the second, the trigonometric functions allow one to define angles among them. In the same way, angles between spacelike and timelike vectors can be defined. Finally, one can define hyperbolic angles among pairs of timelike vectors by taking the difference between them,

fixing a spacelike vector, and using that as a “unit” to treat the difference vector as, in essence, a velocity difference. One can then use the hyperbolic trigonometric to define the angles. (This is essentially the same procedure one uses in general relativity to define hyperbolic angles between timelike vectors; and, in the same way, this structure does not allow one to define orthogonality relations among timelike vectors.) From this, one immediately derives ratios of lengths between all pairs of spacelike vectors, between all pairs consisting of a spacelike and a timelike vector, and between all pairs of timelike vectors.

It will give some insight into the character of a Newtonian conformal structure, as well as being useful in the proof of theorem 3.1, to perform here the calculation that shows that being twist-free for a vector field ( $\nabla^{[a}\xi^{b]} = 0$ ) is a conformally invariant notion, *i.e.*, a conformal derivative operator allows one to determine whether a given vector field is twist-free or not. This makes intuitive sense, as being twist-free essentially means that “nearby” vectors in the vector field have no “angular velocity” with respect to each other, but angular velocity is a conformal notion. More precisely, fix two representative derivative operators of a conformal derivative operator,  $\nabla^a$  and  $\tilde{\nabla}^a$  with difference vector  $C^a{}_{bc}$ , and the  $h^{ab}$  and  $\tilde{h}^{ab}$  in the conformal structure respectively compatible with them, where  $\tilde{h}^{ab} = \Omega^2 h^{ab}$ . First,

$$\begin{aligned} 0 &= \tilde{\nabla}_a \tilde{h}^{bc} \\ &= \tilde{\nabla}_a (\Omega^2 h^{bc}) \end{aligned}$$

so

$$\Omega^2 \tilde{\nabla}_a h^{bc} + h^{bc} \tilde{\nabla}_a \Omega^2 = 0$$

Using  $C^a{}_{bc}$  to re-express this, and noting that  $\nabla_a$  and  $\tilde{\nabla}_a$  agree in their action on  $\Omega^2$ , we get

$$\begin{aligned} 0 &= \Omega^2 \nabla_a h^{bc} + \Omega^2 C^b{}_{an} h^{nc} + \Omega^2 C^c{}_{an} h^{bn} + h^{bc} \nabla_a \Omega^2 \\ &= 2\Omega^2 h^{cn} C^b{}_{an} + h^{bc} \nabla_a \Omega^2 \end{aligned} \tag{2.1}$$

Thus  $2\Omega^2 h^{an} C^b{}_{nc} = -h^{ab} \nabla_c \Omega^2$  and so  $h^{n[a} C^{b]}{}_{nc} = 0$ . It follows immediately that

$$\tilde{\nabla}^{[a}\xi^{b]} = 0 \quad \text{if and only if} \quad \nabla^{[a}\xi^{b]} = 0$$

Now, for a Newtonian conformal structure to be a physically meaningful analogue of conformal structure in general relativity, it should allow one to distinguish a preferred family of images of curves, the conformal spacelike geodesics, *i.e.*, images of curves that can be reparametrized so as to be geodesics for each representative derivative operator of the conformal derivative operator associated with the conformal structure.<sup>3</sup> And it does. They are the spacelike curves that, in an appropriate sense, preserve orthogonality relations. Fix two spacelike vector-fields  $\xi^a$  and  $\eta^a$  everywhere orthogonal to each other. One can always do this. Pick a representative of the conformal

---

<sup>3</sup>Although a conformal structure in general relativity fixes the null geodesics as curves, not just images of curves, we cannot expect that here: that happens in the null case only because all vectors tangent to all curves formed by parametrizing the images have the same length (*viz.*, 0), and so the images remain true geodesics under arbitrary parametrizations. That will not be the case for the spacelike curves here, which will be true geodesics (not just re-parametrizable so as to be geodesics) only under a preferred family of parametrizations.

derivative operator,  $\nabla_a$ ; fix  $\xi^a$  and  $\eta^a$  at a point  $p$ ; then construct a curve (unique up to parametrization) that has  $\xi^a$  as its tangent vector at  $p$ , and that parallel-transport  $\eta^a$  along it with respect to  $\nabla_a$ . Then do the same with the roles of  $\xi^a$  and  $\eta^a$  reversed. Do the same along every point of the constructed curves. Pick a vector  $\theta^a$  orthogonal to both  $\xi^a$  and  $\eta^a$  at  $p$ , and parallel-transport the constructed curves along  $\theta^a$ . And so on. Because (recall) we have assumed that the Newtonian spacetime models we are working with are spatially flat ( $R^{abcd} = 0$ ), we are guaranteed that parallel-transport of all these vectors is path-independent, so the construction is consistent.

The integral curves of  $\xi^a$  and  $\eta^a$  are conformal spacelike geodesics. Pick a representative  $\nabla_a$  of the conformal derivative operator, with its associated  $h^{ab}$ . We will use  $h^{ab}$  to effectively lower indices in the following calculations, by using it to arbitrarily fix spacelike representative 1-forms of  $\xi^a$  and  $\eta^a$  respectively. (It is a simple calculation, following essentially the same lines as the calculation that being twist-free is a conformally invariant notion, to show that the following argument is independent of the initial choice of spacelike representative 1-forms; I leave it to the reader.) Now, it is clear that the two vector-fields Lie-derive each other. Thus  $\xi^n \nabla_n \eta^a = \eta^n \nabla_n \xi^a$ , and, by orthogonality,  $\nabla_a \xi^n \eta_m = 0$ . Now the result follows by playing these off each other in the standard way.

$$\begin{aligned} \eta_m \xi^n \nabla_n \xi^m &= \xi^m \xi^n \nabla_n \eta_m \\ &= \xi^n \nabla_n \xi^m \eta_m - \eta_m \xi^n \nabla_n \xi^m \end{aligned}$$

The first term in the last line is zero by orthogonality, and so we are left with  $\eta_m \xi^n \nabla_n \xi^m = -\eta_m \xi^n \nabla_n \xi^m$ , *i.e.*,  $\eta_m \xi^n \nabla_n \xi^m = 0$ . Since this is true for all vectors orthogonal to  $\xi^a$ , we conclude that  $\xi^n \nabla_n \xi^a = \lambda \xi^a$ , for some scalar field  $\lambda$ . Thus, the integral curves of  $\xi^a$  are geodesics up to reparametrization.

### 3 The Theorem

A projective structure and a Newtonian conformal structure are compatible with each other if the conformal spacelike geodesics determined by the Newtonian conformal structure are also projective geodesics. We can now state the main result of the paper.

**Theorem 3.1** *Two pairs of compatible spatial and temporal metrics  $(h^{ab}, t_{ab})$  and  $(\tilde{h}^{ab}, \tilde{t}_{ab})$  are both projectively and conformally equivalent with respect to compatible projective and conformal structures if and only if the two temporal metrics differ only by a constant factor, and the same holds for the two spatial metrics (though not necessarily by the same factor as for the temporal metrics).*

PROOF:

The “if” part of the theorem is immediate. Assume, then, that we have two pairs of conformally and projectively equivalent pairs of compatible spatial and temporal metrics,  $(h^{ab}, t_{ab})$  and  $(\tilde{h}^{ab}, \tilde{t}_{ab})$  with respect to compatible projective and conformal structures, where  $\tilde{h}^{ab} = \Omega^2 h^{ab}$  and  $\tilde{t}_{ab} = \xi^2 t_{ab}$ . Because their respective compatible derivative operators,  $\nabla_a$  and  $\tilde{\nabla}_a$ , agree on images of geodesics, their difference tensor  $C^a{}_{bc}$  is of the form  $\delta^a{}_{[b} \phi_{c]}$  for some 1-form  $\phi_a$ . In particular, because they are conformally related, from equation (2.1) we know that  $C^{abc} = -\frac{1}{2} h^{bc} \nabla^a \ln \Omega^2$ .

The analogous calculation shows that  $C_{abc} = \frac{1}{2}t_{bc}\nabla_a \ln \chi^2$ . The proof now proceeds in perfect exactly as in the Lorentzian case (Malament 2012, ch. 1, §9, p. 83), playing these expressions of each other, until one derives  $\nabla^a \Omega^2 = 0$  and  $\nabla_a \chi^2 = 0$ . ■

One can now use the theorem to render a physical interpretation to the Newtonian metric structure of a Newtonian spacetime model, in the following way.<sup>4</sup> We will use the following interpretive principles.

**C1** timelike curves represent the possible paths of massive bodies

**C2** segments of images of spacelike geodesics represent the spatial position and extent of (rigid) yard-sticks

**C3** spacelike vectors represent the possible accelerations of massive bodies

**P1** timelike geodesics represent the paths of freely falling massive bodies

**P2** the ratio of the lengths of two segments of a timelike geodesic represent the ratio of the temporal separations between the events at the ends of the two segments

Now, if we can determine when the accelerations of two given massive bodies are orthogonal to each other, then, by **C3**, we can determine the Newtonian conformal structure. This can be operationalized in any of a number of ways. Perhaps the simplest is to use the fact that the orthogonality relations among spacelike vectors are encoded in the ratios of their lengths, and then to use the fact that accelerations are linearly proportional to applied forces. Fix a standard yard-stick (by **C2**) and a standard clock (by **P2**). We can now measure the ratio of the magnitude of two forces by applying each in turn to the same body for a unit of time (based on our clock), and measuring the difference in resultant velocities (based on our yard-stick and our clock). The projective structure is fixed by determining the images of paths of freely falling bodies.

Thus, by the theorem, to know when two forces are orthogonal to each other, and to know the images of the paths of freely falling bodies, is to fix the entirety of the Newtonian metric structure of a Newtonian spacetime model.

## A Appendix: Geometrized Newtonian Gravity

A Newtonian spacetime model is as defined in definition 2.1.

$t_{ab}$  serves to define a notion of a vector's being spacelike or timelike.

**Definition A.1** *The temporal length of a vector  $\xi^a$  is  $(\xi^m \xi^n t_{mn})^{\frac{1}{2}}$ .  $\xi^a$  is timelike if its temporal length is non-zero, spacelike if the vector is non-zero and has a temporal length of zero.*

---

<sup>4</sup>I do not claim this is the unique or even just a canonical way of doing so, or the most elegant or concise or what have you, only that it is a perspicuous and natural way, with clear physical significance.

It is natural to think of the zero-vector as both timelike and spacelike, in so far as we know what it means for two events to occur at the same time as well as at the same place. The addition of any spacelike vector to a timelike one is always timelike. It follows that the family of timelike vectors forms a four-dimensional affine space modeled on the tangent space. (It cannot form a vector-space, as the sum of two timelike vectors may be spacelike.) It also follows that the family of spacelike vectors at a point forms a three-dimensional vector subspace of the tangent space at that point.

The signature of  $t_{ab}$  implies that at every point there exists a neighborhood and smooth 1-form  $t_a$  defined on that neighborhood such that  $t_{ab} = t_a t_b$  on that neighborhood. If there is a globally defined such 1-form, then the spacetime model is *temporally orientable* and  $t_a$  is a *temporal orientation* on it; a timelike vector  $\xi^a$  is future-directed if  $t_n \xi^n > 0$ . Item 6 in definition 2.1 implies that  $t_a$  is closed, and so, at least locally, there exists a smooth function  $t$  such that  $\nabla_a t = t_a$ , the (local) *time-function*. If  $\mathcal{M}$  is simply connected and the spacetime model temporally orientable,  $t$  is defined globally. In this case, its constant surfaces represent the Newtonian idea of “all of space at a given moment of time”, *i.e.*, a maximal collection of events all (absolutely) simultaneous with each other, a *simultaneity slice*, and so  $\mathcal{M}$  has the topology  $\mathbb{R} \times \Sigma$ , where  $\Sigma$  is a three-dimensional manifold diffeomorphic to a simultaneity slice. This  $t$  is unique up to the addition of a constant, which may be thought of as a change of temporal origin. Its scale, however, is fixed by the temporal length of vectors: to multiply  $t_{ab}$  by a constant factor changes nothing physically, but rather represents only a change in units of temporal measurement.

$t_{ab}$ , then, determines an absolute temporal structure—the temporal separation of any two events in the spacetime is fixed once and for all, independent of any other state of affairs in the world; the equivalence classes of spacetime points under the relation “having a temporal separation of zero with” are exactly the simultaneity slices. A *timelike curve* is one whose tangent vectors are everywhere timelike. Such curves represent the possible worldlines of ponderable bodies. The signature of  $t_{ab}$  has another consequence of note: it does not allow one to define orthogonality between two timelike vectors. This makes physical sense: two orthogonal, timelike vector-fields would define time-functions different in the sense that they would not share surfaces of constancy, *i.e.*, they would define incommensurable temporal structures.<sup>5</sup>

$h^{ab}$  defines the spatial metric structure in a more indirect way.

**Proposition A.2**

1. A vector  $\xi^a$  is spacelike at a point  $p$  if and only if there is a covector  $\alpha_a$  at  $p$  such that  $\xi^a = h^{an} \alpha_n$ .
2. For all covectors  $\alpha_a$  and  $\beta_a$  at a point, if  $h^{an} \alpha_n = h^{an} \beta_n$ , then  $h^{mn} \alpha_m \alpha_n = h^{mn} \beta_m \beta_n$ .

---

<sup>5</sup>One can see the physical sense this makes in another, more indirect way: in so far as one can consider all the spacelike vectors at a point in Newtonian spacetime to be the result, in the limit, of “flattening” the null-cones in a relativistic spacetime—“letting the upper bound of possible velocities go to infinity”—the timelike Newtonian vectors at that point encode essentially the same information as the timelike vectors in the interior of the original null-cones, and no two timelike vectors can be orthogonal to each other with respect to a fixed Lorentz metric. See Malament (1986) for a precise characterization and analysis of the process of flattening the null-cones in a relativistic spacetime.



In virtue of this proposition, the following is well formulated.

**Definition A.3** *The spatial length of a spacelike vector  $\xi^a$  is  $(h^{nm}\alpha_n\alpha_m)^{\frac{1}{2}}$ , where  $\alpha_a$  is any covector satisfying  $h^{an}\alpha_n = \xi^a$ .*

If a spacelike  $\eta^a$  equals  $h^{an}\alpha_n$ , then we say  $\alpha_a$  is a *representative spacelike 1-form* of  $\eta^a$ . In so far as  $h^{ab}$  determines lengths only for spacelike vectors, but not at all for timelike vectors (whereas  $t_{ab}$  determines lengths for all spacelike vectors: 0), it defines a spatial metric in only a Pickwickian sense, as it ought to according to Galileian relativity. We know what it means in Newtonian theory to assign a definite distance between two simultaneous events, by employing yard-sticks and the like, independent of any other state of affairs in the world. We do not know how to do so for non-simultaneous events.<sup>6</sup>

The following construction captures the content of this observation. Specify at a point  $p$  a constant timelike vector  $\xi^a$  of unit temporal length, and define  $\hat{h}_{ab}$  to be the unique symmetric tensor at  $p$  satisfying

$$\begin{aligned}\hat{h}_{an}h^{nb} &= \delta_a^b - t_{an}\xi^b\xi^n \\ \hat{h}_{an}\xi^n &= 0\end{aligned}$$

$\hat{h}_{ab}$  represents the covariant spatial metric determined by projection relative to  $\xi^a$ . Then the spatial length of any vector assigned indirectly by  $h^{ab}$  will agree with that assigned directly by  $\hat{h}_{ab}$ . The spatial separation of two non-simultaneous events, however, as determined by such a  $\hat{h}_{ab}$ , depends on the choice of  $\xi^a$ . Again, this is as it should be. If we knew how to assign a spatial length to timelike vectors, and so a fixed spatial separation between non-simultaneous events, then we could define a notion of absolute rest: “a particle is at absolute rest if the timelike curve representing its worldline has everywhere tangent vectors of zero spatial length”. This, however, we cannot do.

In contradistinction to  $t_{ab}$ ,  $h^{ab}$  allows one to define a relation of orthogonality among spacelike vectors: two spacelike vectors  $\eta^a$  and  $\theta^a$  are orthogonal if  $h^{mn}\alpha_m\beta_n = 0$ , where  $\alpha_m$  and  $\beta_n$  are any two representative spacelike 1-forms of  $\eta^a$  and  $\theta^a$  respectively. This makes physical sense as well: we know in Newtonian spacetime how to determine whether two yardsticks at a single moment of time are at a right angle to each other.  $h^{ab}$  also allows us to define a relation of orthogonality between a spacelike and a timelike vector: given a timelike vector  $\xi^a$ , a spacelike vector  $\eta^a$ , and any representative 1-form  $\alpha_a$  for  $\eta^a$ , then  $\xi^n\alpha_n = 0$  if and only if  $\xi^n\beta_n = 0$  for any other representative spacelike 1-form  $\beta_a$ . Though this may sound a little odd at first, it makes physical sense, too. A simple calculation shows that any timelike curve parametrized by arc-length (“proper time”) will satisfy  $\xi^n\nabla_n\xi^a = \eta^a$ , where  $\xi^a$  is the tangent to the curve and  $\eta^a$  is spacelike and orthogonal to  $\xi^a$ . This says that for any ponderable body, the worldline of which by definition instantiates a timelike curve, its acceleration is everywhere spacelike, and thus a viable candidate for entering Newton’s Second Law on the righthand side. If this were not the case, then one could have ponderable bodies accelerating in a timelike direction, with attendant timelike forces, and I see no possible way to make

---

<sup>6</sup>See [Stein \(1967\)](#) for a thorough discussion.

physical sense of such a conceit, except perhaps in a 1950’s B-movie “Great Scot! He’s beginning to mutater!”-sort of way. Thus, we can assign determinate magnitudes, fixed once and for all, to the acceleration of bodies, as Galileian relativity allows us to do, and Newtonian mechanics demands we do (Stein 1967). In sum,  $h^{ab}$  encodes the right amount of structure of the right kind, nothing more, nothing less. (One may think of this as physics practiced in accord with the Goldilocks Principle.)

In what follows, we assume that  $\mathcal{M}$  is simply connected and the spacetime model is temporally orientable; this is no real loss of generality, for all arguments and conclusions would still go through without the assumption, at the cost of constant hedging and much technical work of a nit-picky sort without any counter-balancing gain in physical insight. In particular, we do not need to worry about which results hold only locally and which globally.

Before moving on to the definition of the appropriate analogue of a conformal structure for a Newtonian metric structure, we rehearse a few of its properties and state the most important two theorems in geometrized Newtonian gravity, which will be of use later. When there is no chance for ambiguity, we will use raised indices to represent the action of  $h^{ab}$  on a tensorial object, e.g.,  $\nabla^a := h^{an}\nabla_n$ .

**Theorem A.4 (The Geometrization Lemma)** *Fix a Newtonian spacetime model  $(\mathcal{M}, \rho, h^{ab}, t_{ab}, \nabla_a)$ , such that  $\nabla_a$  is flat, i.e., its associated Riemann tensor  $R^a{}_{bcd} = 0$ , and a smooth scalar field  $\phi$  such that  $\nabla^n\nabla_n\phi = 4\pi\rho$  (the Poisson equation). Define another derivative operator  $\tilde{\nabla}_a = (\nabla_a, C^a{}_{bc})$ , where  $C^a{}_{bc} = -t_{cb}\nabla^a\phi$ . Then:*

1.  $(\mathcal{M}, \rho, h^{ab}, t_{ab}, \tilde{\nabla}_a)$  is a Newtonian spacetime model
2.  $\tilde{\nabla}_a$  is the unique derivative operator such that

$$\xi^n\tilde{\nabla}_n\xi^a = 0 \quad \text{if and only if} \quad \xi^n\nabla_n\xi^a = -\nabla^a\phi$$

3. the Riemann tensor  $\tilde{R}^a{}_{bcd}$  associated with  $\tilde{\nabla}_a$  satisfies

- a.  $\tilde{R}_{ab} = 4\pi\rho t_{ab}$
- b.  $\tilde{R}^a{}_b{}^c{}_d = \tilde{R}^c{}_d{}^a{}_b$
- c.  $\tilde{R}^{ab}{}_{cd} = 0$

A few remarks are in order to explain the physical significance of the theorem. In essence it says that, given a Newtonian spacetime model with a flat affine structure, we can always construct one in which the geodesics (possible paths of “freely falling bodies”) of a curved affine structure are the same paths as those representing bodies moving under the force of the ambient gravitational field associated with  $\rho$  in the original model. The constructed model “geometrizes” gravity: it incorporates the effects of gravity into the metric structure of the new model, in analogy with general relativity.

The effective converse of the Geometrization Lemma holds as well.

**Theorem A.5 (Trautman Recovery Theorem)** *Fix a Newtonian spacetime model  $(\mathcal{M}, \rho, h^{ab}, t_{ab}, \nabla_a)$  satisfying*

1.  $R_{ab} = 4\pi\rho t_{ab}$
2.  $R^a{}_b{}^c{}_d = R^c{}_d{}^a{}_b$
3.  $R^{ab}{}_{cd} = 0$

Then there is a derivative operator  $\tilde{\nabla}^a$  and a scalar field  $\tilde{\phi}$  such that

1.  $\tilde{\nabla}^n \tilde{\nabla}_n \tilde{\phi} = 4\pi\rho$  (the Poisson equation)
2.  $\tilde{\nabla}^a$  is compatible with  $t_{ab}$  and  $h^{ab}$
3.  $\tilde{\nabla}^a$  is flat
4. for all timelike curves with tangent vector  $\xi^a$

$$\xi^n \tilde{\nabla}_n \xi^a = 0 \quad \text{if and only if} \quad \xi^n \tilde{\nabla}_n \xi^a = -\tilde{\nabla}^a \tilde{\phi}$$

Moreover,  $(\tilde{\nabla}_a, \tilde{\phi})$  is not unique. Any other such pair  $(\hat{\nabla}_a, \hat{\phi})$  will satisfy the stated conditions if and only if

- a.  $\nabla^a \nabla^b (\tilde{\phi} - \hat{\phi}) = 0$
- b.  $\hat{\nabla}_a = (\tilde{\nabla}_a, C^a{}_{bc})$ , where  $C^a{}_{bc} = t_{bc} \nabla^a (\tilde{\phi} - \hat{\phi})$

Given a Newtonian spacetime model in which the gravitational effects of  $\rho$  are geometrized (incorporated into the curved affine structure), the theorem tells us that we can recover one in which the gravitational effects of  $\rho$  are rather represented by the acceleration of the paths of freely-falling bodies, as defined by a flat affine structure.

In disanalogy with general relativity, and crucially for our purposes, in this context  $R^{ab}{}_{cd} = 0$  is *not* equivalent to  $R^a{}_{bcd} = 0$ . The interpretation of the condition  $R^{ab}{}_{cd} = 0$  and its relative strength as compared to  $R^a{}_{bcd} = 0$  follow from a proposition whose proof is straightforward, albeit tedious (Malament 2012, §§4.2–4.3). To state it, we need a few more definitions. A Newtonian spacetime model is *spatially flat* if  $R^{abcd} = 0$ , *i.e.*, if the affine structure on the simultaneity slices derived by restricting to them the action of the global affine structure is flat. A unit, future-directed timelike vector-field  $\xi^a$  is *rigid* if  $\mathcal{L}_\xi h^{ab} = 0$ , *i.e.*, if  $\nabla^{(a} \xi^{b)} = 0$ . (One may think of a rigid vector field as the analogue of a Killing field in general relativity.)  $\xi^a$  is *twist-free* if  $\nabla^{[a} \xi^{b]} = 0$ , and is *acceleration-free* if  $\nabla_a \xi^b = 0$ .

**Proposition A.6** *Fix a spatially flat Newtonian spacetime model. Then:*

1.  $R^{ab}{}_{cd} = 0$  if and only if there exists a rigid and twist-free vector field
2.  $R^a{}_{bcd} = 0$  if and only if there exists a rigid, twist-free and acceleration-free vector field

Another illuminating characterization of  $R^{ab}{}_{cd} = 0$  follows from this proposition: the condition holds if and only if parallel-transport of spacelike vectors is path-independent.

## References

- Eddington, A. (1923). *Mathematical Theory of Relativity* (Second ed.). Cambridge: Cambridge University Press.
- Ehlers, J., F. Pirani, and A. Schild (1972). The geometry of free fall and light propagation. In L. O’Raifeartaigh (Ed.), *General Relativity: Papers in Honour of J. L. Synge*, Chapter 4, pp. 63–84. Oxford: Clarendon Press. Republished in *General Relativity and Gravitation*, 44(2012, 6):1587–1609, doi:10.1007/s10714-012-1353-4.
- Einstein, A. (2001). *Relativity: The Special and General Theory* (Fifteenth, enlarged ed.). London: Routledge. Translated by R. W. Lawson. The original German was published in 1916. The first English edition was published in 1920 by Methuen & Co., London. The Fifteenth edition was first published in 1954.
- Hawking, S. and G. Ellis (1973). *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press.
- Malament, D. (1986). Newtonian gravity, limits, and the geometry of space. In R. Colodny (Ed.), *From Quarks to Quasars: Philosophical Problems of Modern Physics*, pp. 181–201. Pittsburgh: Pittsburgh University Press.
- Malament, D. (2012). *Topics in the Foundations of General Relativity and Newtonian Gravitational Theory*. Chicago: University of Chicago Press. Uncorrected final proofs for the book are available for download at <http://strangebeautiful.com/other-texts/malament-founds-gr-ngt.pdf>.
- Møller, C. (1952). *The Theory of Relativity* (First ed.). Oxford: The Clarendon Press.
- Stein, H. (1967). Newtonian space-time. *Texas Quarterly* 10, 174–200.
- Synge, J. (1960). *Relativity: The General Theory*. Amsterdam: North-Holland Publishing Co.
- Trautman, A. (1965). Foundations and current problems of general relativity. In S. Deser. and K. Ford (Eds.), *Lectures on General Relativity*, Volume 1 of *Brandeis Summer Institute in Theoretical Physics*, pp. 1–248. Englewood Cliffs, NJ: Prentice-Hall, Inc.
- Weyl, H. (1918). *Raum-Zeit-Materie: Vorlesungen ber Allgeimene Relativitätstheorie*. Berlin: Springer-Verlag.