

Overview Harman ● Hempel ○ Carnap ○○ Goodman ○○○○ (RTE) ○○○ References Extras ○○○○○○

- Here is a “*reductio*” of classical deductive logic (this is quite naïve, but adding sophistication won’t help — see next slide):
  - (1) For all sets of statements  $X$  and all statements  $p$ , if  $X$  is inconsistent, then  $p$  is a logical consequence of  $X$ .
  - (2) If an agent  $S$ ’s belief set  $B$  entails  $p$  (and  $S$  knows  $B \models p$ ), then it would be reasonable for  $S$  to infer/believe  $p$ .
  - (3) Even if  $S$  knows their belief set  $B$  is inconsistent (and, hence, that  $B \models p$ , for any  $p$ ), there are still some  $p$ ’s such that it would *not* be reasonable for  $S$  to infer/believe  $p$ .
  - (4)  $\therefore$  Since (1)–(3) lead to absurdity, our initial assumption (1) must have been false — *reductio* of the “explosion” rule (1).
- Harman [8] would concede that (1)–(3) are inconsistent, and (as a result) that *something* is wrong with premises (1)–(3).
- But, he would reject the relevantists’ diagnosis that (1) must be rejected. I take it he’d say it’s (2) that is to blame here.
- 👉 (2) is a *bridge principle* [12] linking *entailment* and *inference*.
- (2) is correct *only* for *consistent*  $B$ ’s. [Even if  $B$  is consistent, the correct response *may* rather be to *reject* some  $B_i$ ’s in  $B$ .]

Branden Fitelson Goodman’s “Grue” Argument in Historical Perspective fitelson.org

Overview Harman ○● Hempel ○ Carnap ○○ Goodman ○○○○ (RTE) ○○○ References Extras ○○○○○○

- Note: the choice of *deductive* contexts in which  $S$ ’s belief set  $B$  is (known by  $S$  to be) *inconsistent* is intentional here.
- In such contexts, there is a *deep disconnect* between (known) *entailment* relations and (kosher) *inferential* relations.
- One might try a more sophisticated deductive bridge principle (2’) here. But, I conjecture a *dilemma*. Either:
  - (2’) will be *too weak* to yield a (classically) *valid* “*reductio*”.
  - or
  - (2’) will be *false*. [Our original BP (2) falls under this horn.]
- Let  $B$  be  $S$ ’s belief set, and let  $q$  be the conjunction of the elements  $B_i$  of  $B$ . Here are two more candidate BP’s:
  - (2’<sub>1</sub>) If  $S$  knows that  $B \models p$ , then  $S$  should *not* be such that *both*:  $S$  believes  $q$ , and  $S$  does not believe  $p$ .
  - (2’<sub>2</sub>) If  $S$  knows that  $B \models p$ , then  $S$  should *not* be such that *both*:  $S$  believes each of the  $B_i \in B$ , and  $S$  does not believe  $p$ .
- (2’<sub>2</sub>) is *false* (preface paradox) and too weak (it’s wide scope).
- (2’<sub>1</sub>) *may* be true, but it is also *too weak*. [It’s wide scope, and the agent can reasonably disbelieve *both*  $q$  and  $p$ .]

Branden Fitelson Goodman’s “Grue” Argument in Historical Perspective fitelson.org

Overview Harman ○○ Hempel ○ Carnap ○○ Goodman ○○○○ (RTE) ○○○ References Extras ○○○○○○

## “Potted History” Version of Goodman’s Argument

- Consider the following two inductive arguments:
 

$(\mathcal{A}_1)$ $(E_1)$ $a$ is a green emerald. $\therefore (H_1)$ All emeralds are green.		$(\mathcal{A}_2)$ $(E_2)$ $a$ is a grue emerald. $\therefore (H_2)$ All emeralds are grue.
---	--	---
- A “potted history” version of Goodman’s argument ([7]):
  - (1) Arguments  $(\mathcal{A}_1)$  and  $(\mathcal{A}_2)$  have the same logical form.
  - (2) Argument  $(\mathcal{A}_1)$  is “inductively valid” (i.e.,  $E_1$  confirms  $H_1$ ).
  - (3)  $(\mathcal{A}_2)$  is *not* “inductively valid” (i.e.,  $E_2$  does *not* confirm  $H_2$ ).
  - (4)  $\therefore$  “Inductive validity” is not *merely* a matter of logical form.
- My talk today aims mainly to undermine Goodman’s argument (in FF&F [7]) for premises (2) and (3).
  - Sidebar: I also think (1) is *question-begging*. I won’t be able to get to this today, but see my “Extras” slides for more.
- 👉 Goodman’s argument against *inductive* logic is analogous to the (unsound) argument above against classical *deductive* logic. This is what the rest of the talk will aim to establish.

Branden Fitelson Goodman’s “Grue” Argument in Historical Perspective fitelson.org

Overview Harman ○○ Hempel ● Carnap ○○ Goodman ○○○○ (RTE) ○○○ References Extras ○○○○○○

- I’ll begin by laying out the salient bits of the inductive logical (*viz.*, *confirmation*) theories of Hempel and Carnap.
- Hempelian confirmation theory uses *entailment* to explicate “inductive logical support” (confirmation), which is a logical relation between statements. [ $E$  confirms  $H$  iff  $E \models \text{dev}_E(H)$ ]
- Hempel’s theory has the following three key consequences:
  - (EQC) If  $E$  confirms  $H$  and  $E \models E'$ , then  $E'$  confirms  $H$ .
  - (NC) For all constants  $x$  and all (consistent) predicates  $\phi$  and  $\psi$ : ‘ $\phi x \ \& \ \psi x$ ’ confirms ‘ $(\forall y)(\phi y \supset \psi y)$ ’.
  - (M) For all  $x$ , for all (consistent)  $\phi$  and  $\psi$ , and all statements  $H$ : If ‘ $\phi x$ ’ confirms  $H$ , then ‘ $\phi x \ \& \ \psi x$ ’ confirms  $H$ .
- These three properties are the crucial ones needed to reconstruct Goodman’s “grue” argument against Hempel.
- Before giving a precise reconstruction of Goodman’s “grue” argument, we’ll look at the essentials of Carnapian IL/CT. [Goodman targeted both Hempel and Carnap in FF&F [7].]


Branden Fitelson Goodman’s “Grue” Argument in Historical Perspective fitelson.org

- Carnapian confirmation (*i.e.*, later Carnapian theory [13]) is based on *probabilistic relevance*, not deductive entailment:
  - $E$  confirms  $H$ , relative to  $K$  iff  $\Pr(H | E \& K) > \Pr(H | K)$ , for some “suitable” conditional probability function  $\Pr(\cdot | \cdot)$ .
    - Note how this is an *explicitly* 3-place relation. Hempel’s was only 2-place. This is because  $\Pr$  (unlike  $\models$ ) is *non-monotonic*.
    - Carnap thought “suitable  $\Pr$ ” meant “logical  $\Pr$ ” in a very strong/naive sense. But, Goodman’s argument (charitably reconstructed) will work against *any* probability function  $\Pr$ .
- 👉 Carnap’s theory implies *only 1* of our 3 Hempelian claims: (EQC). It does *not* imply either (NC) or (M) (see [3]/[13]).
  - This will allow Carnapian IL to avoid facing the full brunt of Goodman’s “grue” (but, it will still face a serious challenge).
- For Carnap, confirmation is a *logical* relation (akin to entailment). Like entailment, confirmation can be *applied*, but this requires *epistemic bridge principles* [akin to (2)].
- Carnap [1] discusses various bridge principles. The most well-known of these is the *requirement of total evidence*.

- **The Requirement of Total Evidence.** In the application of IL to a given knowledge situation, the total evidence available must be taken as a basis for determining the degree of confirmation.
- This *sounds* like a plausible principle. But, once it is made more precise, it will actually turn out to be subtly defective.
- More precisely, we have the following *bridge principle* connecting *confirmation* and *evidential support*:
  - (RTE)  $E$  evidentially supports  $H$  for  $S$  in  $C$  iff  $E$  confirms  $H$ , relative to  $K$ , where  $K$  is  $S$ ’s *total evidence* in  $C$ .
- The (RTE) has often been (implicitly) presupposed by Bayesian epistemologists (both subjective and objective).
- However, as we will soon see, the (RTE) is not a tenable bridge principle, and for reasons independent of “grue”.
- 👉 Moreover, Goodman’s “grue” argument will rely *more heavily* on (RTE) than the relevantists’ argument relies on (2). In this sense, Goodman’s argument will be *even worse*.
- Before reconstructing the argument, a brief “grue” primer.

- Let  $Gx \stackrel{\text{def}}{=} x$  is green,  $Ox \stackrel{\text{def}}{=} x$  is examined prior to  $t$ , and  $Ex \stackrel{\text{def}}{=} x$  is an emerald. Goodman introduces a predicate “grue”
 
$$Gx \stackrel{\text{def}}{=} x \text{ is grue} \stackrel{\text{def}}{=} Ox \equiv Gx.$$
- Consider the following two universal generalizations
  - ( $H_1$ ) All emeralds are green.  $[(\forall x)(Ex \supset Gx)]$
  - ( $H_2$ ) All emeralds are grue.  $[(\forall x)[Ex \supset (Ox \equiv Gx)]]$
- And, consider the following instantial evidential statement
  - ( $E$ )  $Ea \& Oa \& Ga$
- Hempel’s confirmation theory [(EQC) & (NC) & (M)] entails:
  - ( $\dagger$ )  $E$  confirms  $H_1$ , and  $E$  confirms  $H_2$ . [proof]
- As a result, his theory entails the following weaker claim
  - ( $\ddagger$ )  $E$  confirms  $H_1$  if and only if  $E$  confirms  $H_2$ .
- What about (later) Carnapian theory? Does it entail even ( $\ddagger$ )?
- 👉 Interestingly, NO! There are (later) Carnapian  $\Pr$ -models in which  $E$  confirms  $H_1$  but  $E$  *disconfirms*  $H_2$ .
- So, Hempel was an easier target for Goodman than (later) Carnap (to be fair, Goodman talks only about *early* Carnap).
- Now, we’re ready to reconstruct Goodman’s argument.

### A Proof of ( $\dagger$ ) From Hempel’s (NC), (M), and (EQC)



← back

Overview Harman Hempel Carnap Goodman (RTE) References Extras

- There is just one more ingredient in Goodman's argument:
  - The agent  $S$  who is assessing the evidential support that  $\mathcal{E}$  provides for  $H_1$  vs  $H_2$  in a Goodmanian "grue" context  $C_G$  has  $Oa$  as part of their total evidence in  $C_G$ . (e.g., [14].)
- Now, we can run the following Goodmanian *reductio*:
  - $E$  confirms  $H$ , relative to  $K$  iff  $\Pr(H | E \& K) > \Pr(H | K)$ .
  - $E$  evidentially supports  $H$  for  $S$  in  $C$  iff  $E$  confirms  $H$ , relative to  $K$ , where  $K$  is  $S$ 's total evidence in  $C$ .
  - The agent  $S$  who is assessing the evidential support  $\mathcal{E}$  provides for  $H_1$  vs  $H_2$  in a Goodmanian "grue" context  $C_G$  has  $Oa$  as part of their total evidence in  $C_G$  [i.e.,  $K \models Oa$ ].
  - If  $K \models Oa$ , then—c.p.— $\mathcal{E}$  confirms  $H_1$  relative to  $K$  iff  $\mathcal{E}$  confirms  $H_2$  relative to  $K$ , for **any**  $\Pr$  [i.e.,  $(\ddagger)$  holds,  $\forall \Pr$ 's].
  - Therefore,  $\mathcal{E}$  evidentially supports  $H_1$  for  $S$  in  $C_G$  if and only if  $\mathcal{E}$  evidentially supports  $H_2$  for  $S$  in  $C_G$ .
  - $\mathcal{E}$  evidentially supports  $H_1$  for  $S$  in  $C_G$ , but  $\mathcal{E}$  does *not* evidentially support  $H_2$  for  $S$  in  $C_G$ .
- $\therefore$  (i)–(vi) lead to an absurdity. Hence, our initial assumption (i) must have been false. Carnapian inductive logic refuted?

Branden Fitelson Goodman's "Grue" Argument in Historical Perspective fitelson.org

Overview Harman Hempel Carnap Goodman (RTE) References Extras

- Premise (vi) is based on Goodman's *epistemic intuition* that, in "grue" contexts,  $\mathcal{E}$  evidentially supports  $H_1$  but *not*  $H_2$ .
- Premise (v) follows logically from premises (i)–(iv).
- Premise (iv) is a theorem of probability calculus (**any**  $\Pr$ !).
  - The c.p. clause needed is  $\Pr(Ea | H_1 \& K) = \Pr(Ea | H_2 \& K)$ , which is assumed in all probabilistic renditions of "grue".
- Premise (iii) is an assumption about the agent's background knowledge  $K$  that's implicit in Goodman's set-up. See [14].
- Premise (ii) is (RTE). It's the *bridge principle*, akin to (2) in the relevantists' *reductio*. This is the premise I will focus on.
- Here are my two main points about Goodman's argument:
  - (ii) must be rejected by Bayesians for independent reasons.
  - Carnapian confirmation theory *doesn't even entail*  $(\ddagger)$ . [Hempel's theory does, just as deductive logic entails (1).]
- This suggests Goodman's argument is *even less a reductio* of (i) than the relevantists' argument is a *reductio* of (1).
- Moreover, a careful reading of *Fact, Fiction, and Forecast* reveals that this *was* Goodman's argumentative strategy.

Branden Fitelson Goodman's "Grue" Argument in Historical Perspective fitelson.org

Overview Harman Hempel Carnap Goodman (RTE) References Extras

## Three Salient Quotes from Goodman [7]

👉 The "new riddle" is *about inductive logic (not epistemology)*.

**Quote #1** (page 67): "Just as deductive logic is concerned primarily with a relation between statements — namely the consequence relation — that is independent of their truth or falsity, so inductive logic ... is concerned primarily with a comparable relation of confirmation between statements. Thus the problem is to define the relation that obtains between any statement  $S_1$  and another  $S_2$  if and only if  $S_1$  may properly be said to confirm  $S_2$  in any degree."

**Quote #2** (73): "Confirmation of a hypothesis by an instance depends ... upon features of the hypothesis other than its syntactical form".

👉 But, Goodman's *methodology* appeals to *epistemic intuitions*.

**Quote #3** (page 73): "... the fact that a given man now in this room is a third son *does not increase the credibility of* statements asserting that other men now in this room are third sons, *and so does not confirm* the hypothesis that all men now in this room are third sons."

Branden Fitelson Goodman's "Grue" Argument in Historical Perspective fitelson.org

Overview Harman Hempel Carnap Goodman (RTE) References Extras

- As Tim Willimson points out [16, ch. 9], Carnap's (RTE) must be rejected, because of the problem of old evidence [2].
- If  $S$ 's total evidence in  $C$  ( $K$ ) entails  $E$ , then, according to (RTE),  $E$  cannot evidentially support *any*  $H$  for  $S$  in  $C$ .
- As a result, there are  $C$ 's in which we can't use  $\Pr(\cdot | K)$  — for *any*  $\Pr$  — when assessing the *evidential import of*  $E$  in  $C$ .
- There are (basically) two kinds of strategies for revising (RTE). Carnap [1, p. 472] & Williamson [16, ch. 9] suggest:
 

(RTE<sub>+</sub>)  $E$  evidentially supports  $H$  for  $S$  in  $C$  iff  $S$  possesses  $E$  as evidence in  $C$  and  $\Pr_+(H | E \& K_+) > \Pr_+(H | K_+)$ . [ $K_+$  is "empty",  $\Pr_+$  is "inductive" [13]/"evidential" [16]/"logical" [1].]
- Note: Hempel explicitly *required* that confirmation be taken "*relative to*  $K_+$ " in all treatments of the paradoxes [9, 10]. (RTE<sub>+</sub>) is a charitable Carnapian reconstruction of Hempel.
- A more "standard" way to revise (RTE) is [(RTE')] to use  $\Pr_{S'}(\cdot | K')$ , where  $K \models K' \neq E$ , and  $\Pr_{S'}$  is the credence function of a "counterpart"  $S'$  of  $S$  with total evidence  $K'$ .

Branden Fitelson Goodman's "Grue" Argument in Historical Perspective fitelson.org

- Carnap never re-wrote the part of LFP [1] that discusses the (RTE), in light of a probabilistic *relevance* (“increase in firmness” [1]) notion of confirmation. This is too bad.
- If Carnap had discussed this (“old evidence”) issue, I suspect he would have used something like (RTE<sub>τ</sub>) as his bridge principle connecting confirmation and evidential support.
- Various other philosophers have proposed similar accounts of “support” as some probabilistic relation, taken relative to an “empty” (perhaps “*a priori*”) background &/∨ probability.
  - Richard Fumerton (who, unlike Williamson, is an epistemological *internalist*) proposes such a view in his [4].
  - Patrick Maher [13] applies such relations extensively in his recent (neo-Carnapian) work on confirmation theory.
  - Brian Weatherson [15] uses a similar, “Keynesian” [11] inductive-probability approach to evidential support.
- So, many Bayesians *already* reject (RTE). [Of course, “grue” gives Bayesians another important reason to reject (RTE). ]

- So far, I have left open (precisely) what I think Bayesian confirmation theorists *should* say (*logically & epistemologically*) in light of Goodman’s “grue” paradox.
- Clearly, BCTs will need to revise (RTE) in light of “grue”. But, the standard (RTE’) way of doing this to cope with “old evidence” isn’t powerful enough to avoid *both* problems.
- The more draconian (RTE<sub>τ</sub>) — suggested by the work of Carnap — avoids both problems, from a *logical* point of view (*if* “inductive”/“logical” probabilities *exist!*). But, what should would-be “Carnapians” say on the *epistemic* side?
- I’m not sure what the evidential relations *are* in “grue” contexts (but, see “Extras”). But, *that* doesn’t undermine my line on Goodman’s “grue” *argument* against *inductive logic*.
- Analogy: Harman doesn’t tell us (in general) how someone *should* respond to the discovery that their beliefs are inconsistent. But, *that* doesn’t undermine Harman’s points about relevantist “reductios” of classical deductive logic.

[1] R. Carnap, *Logical Foundations of Probability*, 2nd ed., Chicago Univ. Press, 1962.

[2] E. Eells, *Bayesian problems of old evidence*, in C. Wade Savage (ed.) *Scientific theories, Minnesota Studies in the Philosophy of Science* (Vol. X), 205-223, 1990.

[3] B. Fitelson, *The Paradox of Confirmation*, *Philosophy Compass* (online publication), Blackwell, 2006. URL: <http://fitelson.org/ravens.htm>.

[4] R. Fumerton, *Metaepistemology and Skepticism*, Rowman & Littlefield, 1995.

[5] C. Glymour, *Theory and Evidence*, Princeton University Press, 1980.

[6] I.J. Good, *The white shoe is a red herring*, *BJPS* 17 (1967), 322.

[7] N. Goodman, *Fact, Fiction, and Forecast*, Harvard University Press, 1955.

[8] G. Harman, *Change in View: Principles of Reasoning*, MIT Press, 1988.

[9] C. Hempel, *Studies in the logic of confirmation*, *Mind* 54 (1945), 1-26, 97-121.

[10] ———, *The white shoe: no red herring*, *BJPS* 18 (1967), 239-240.

[11] J. Keynes, *A Treatise on Probability*, Macmillan, 1921.

[12] J. MacFarlane, *In what sense (if any) is logic normative for thought?*, 2004.

[13] P. Maher, *Probability captures the logic of scientific confirmation*, *Contemporary Debates in the Philosophy of Science* (C. Hitchcock, ed.), Blackwell, 2004.

[14] E. Sober, *No model, no inference: A Bayesian primer on the grue problem*, in *grue! The New Riddle of Induction* (D. Stalker ed.), Open Court, Chicago, 1994.

[15] B. Weatherson, *The Bayesian and the Dogmatist*, manuscript, 2007. URL: <http://brian.weatherson.org/tbatd.pdf>.

[16] T. Williamson, *Knowledge and its Limits*, Oxford University Press, 2000.

## “Carnapian” Counterexamples to (NC) and (M)

- (K) Either: (H) there are 100 black ravens, no nonblack ravens, and 1 million other things, or (~H) there are 1,000 black ravens, 1 white raven, and 1 million other things.
- Let  $E \stackrel{\text{def}}{=} Ra \ \& \ Ba$  ( $a$  randomly sampled from universe). Then:
 
$$\Pr(E \mid H \ \& \ K) = \frac{100}{1000100} \ll \frac{1000}{1001001} = \Pr(E \mid \sim H \ \& \ K)$$
  - ∴ This  $K/\Pr$  constitute a counterexample to (NC), assuming a “Carnapian” theory of confirmation. This model can be emulated in the later Carnapian  $\lambda/\gamma$ -systems [13].
- 
- Let  $Bx \stackrel{\text{def}}{=} x$  is a black card,  $Ax \stackrel{\text{def}}{=} x$  is the ace of spades,  $Jx \stackrel{\text{def}}{=} x$  is the jack of clubs, and  $K \stackrel{\text{def}}{=} a$  card  $a$  is sampled at random from a standard deck (where  $\Pr$  is also standard):
    - $\Pr(Aa \mid Ba \ \& \ K) = \frac{1}{26} > \frac{1}{52} = \Pr(Aa \mid K)$ .
    - $\Pr(Aa \mid Ba \ \& \ Ja \ \& \ K) = 0 < \frac{1}{52} = \Pr(Aa \mid K)$ .

## A “Carnapian” Counterexample to (‡)

- (K) Either: ( $H_1$ ) there are 1000 green emeralds 900 of which have been examined before  $t$ , no non-green emeralds, and 1 million other things in the universe, or ( $H_2$ ) there are 100 green emeralds that have been examined before  $t$ , no green emeralds that have not been examined before  $t$ , 900 non-green emeralds that have not been examined before  $t$ , and 1 million other things.
- Imagine an urn containing true descriptions of each object in the universe (Pr  $\stackrel{\text{def}}{=}$  urn model). Let  $\mathcal{E} \stackrel{\text{def}}{=} “Ea \ \& \ Oa \ \& \ Ga”$  be drawn.  $\mathcal{E}$  confirms  $H_1$  but  $\mathcal{E}$  disconfirms  $H_2$ , relative to K:

$$\Pr(\mathcal{E} \mid H_1 \ \& \ K) = \frac{900}{1001000} > \frac{100}{1001000} = \Pr(\mathcal{E} \mid H_2 \ \& \ K)$$

- This  $K/\text{Pr}$  constitute a counterexample to (‡), assuming a “Carnapian” theory of confirmation. This probability model can be emulated in the later Carnapian  $\lambda/\gamma$ -systems [13].

## What Could “Carnapian” Inductive Logic Be? Part I

- Many logical empiricists dreamt that inductive logic (confirmation theory) could be formulated in such a way that it *supervenes* on deductive logic in a *very strong* sense.
  - Strong Supervenience (SS).** All confirmation relations involving sentences of a first-order language  $\mathcal{L}$  supervene on the *deductive-logical* (*viz.*, syntactical) structure *of*  $\mathcal{L}$ .
- Hempel clearly saw (SS) as a *desideratum* for confirmation theory. The early Carnap also seems to have (SS) in mind.
- I think it is fair to say that Carnap’s project — understood as requiring (SS) — was unsuccessful. [Note: I think this is true for reasons that are *independent* of Goodman’s “grue”.]
- The later Carnap seems to be aware of this. Most commentators interpret this shift as the later Carnap simply *giving up* on inductive logic (*qua logic*) altogether.
- I want to resist this “standard” reading of the history.

## What Could “Carnapian” Inductive Logic Be? Part II

- I propose a different reading of the later Carnap, which makes him much more coherent with the early Carnap.
- I propose *weakening* the supervenience requirement in such a way that it (a) ensures this coherence, and (b) maintains the “logicality” of confirmation relations in Carnap’s sense.
- Let  $\mathcal{L}$  be a formal language strong enough to express the fragment of probability theory Carnap needs for his later, more sophisticated confirmation-theoretic framework.
  - Weak Supervenience (WS).** All confirmation relations involving sentences of a first-order language  $\mathcal{L}$  supervene on the *deductive-logical* (*viz.*, syntactical) structure *of*  $\mathcal{L}$ .
- Happily,  $\mathcal{L}$  is pretty weak (Carnap’s  $\mathfrak{c}$ -theories are *decidable*). So, even by early (*logician*) Carnapian lights, satisfying (WS) is sufficient to ensure the “logical determinateness” of  $\mathfrak{c}$ .
- The specific (WS) approach I favor takes confirmation to be a 4-place relation: between  $E$ ,  $H$ ,  $K$ , and a Pr-model  $\mathcal{M}$ .

## What Could “Carnapian” Inductive Logic Be? Part III

- Consequences of moving to such a 4-place  $\mathfrak{c}$ -relation:
  - We need not try to “construct” “logical” probability functions from the syntax of  $\mathcal{L}$ . This is a dead-end anyhow.
  - Indeed, on this view, inductive logic has nothing to say about the *interpretation/origin* of Pr. That is *not* a *logical* question, but a question about the *application* of logic.
    - Analogy: Deductive logicians don’t owe us a “logical interpretation/construction” of the *valuation function*.
  - Moreover, this leads to a vast increase in the *generality* of inductive logic. Carnap was stuck with an impoverished set of “logical” probability functions (in his  $\lambda/\gamma$ -continuum).
    - On my approach, *any* probability function can be part of a confirmation relation (*via*  $\mathcal{M}$ ). Which functions are “appropriate” or “interesting” will depend on *applications*.
    - So, some confirmation relations will not be “interesting”, *etc.* But, this is (already) true of *entailments*, as Harman showed.
  - Questions: Now, what *is* the job of the inductive logician, and how (if at all) do they interact with *epistemologists*?

## What Could “Carnapian” Inductive Logic Be? Part IV

- The inductive logician must explain how it is that inductive logic can satisfy the following Carnapian *desiderata*.
  - The confirmation function  $c_{\mathcal{M}}(H, E | K)$  quantifies a *logical* (in a Carnapian sense) relation between  $E$ ,  $H$ , and  $K$ .
    - (D<sub>1</sub>) “Logical determinateness” of  $c$  is ensured by the move from (SS) to (WS) [from an  $\mathcal{L}$ -determinate to an  $\mathcal{L}$ -determinate  $c$ ].
    - (D<sub>2</sub>) Another aspect of “logicality” insisted upon by Carnap is that  $c_{\mathcal{M}}(H, E | K)$  should *generalize* the entailment relation.
      - At least:  $c_{\mathcal{M}}(H, E | K)$  should take a max (min) value when  $E \& K \models H$  ( $E \& K \models \sim H$ ) — for **all** (regular) Pr-models  $\mathcal{M}$ .
  - (D<sub>3</sub>) There must be *some* interesting “bridge principles” linking  $c$  and *some* relations of evidential support, in *some* contexts.
    - My basic “bridging” idea (rough): subject-context pairs  $\langle S, C \rangle$  will determine “epistemically appropriate” Pr-models  $\mathcal{M}$ .
    - (D<sub>2</sub>) implies that *if* there are any such bridge principles linking *entailment* and (say) *conclusive evidence*, these will be *inherited by*  $c$ . So, we also inherit Harman’s problem!

## “Potted History” Version of Goodman’s Argument (#2)

- Some say that “sensitivity to choice of language” is a central/essential theme/aspect of Goodman’s argument.
- But, this cannot be the case. It’s easy to see why.
  - 1 Goodman’s main target was *Hempel*.
  - 2 Hempel’s  $c$ -relation is defined in terms of  $\models$ .
  - 3  $\models$  is *not* (essentially) sensitive to choice of language.
  - 4 Or, if  $\models$  is sensitive to choice of language (and said sensitivity is *essential* to Goodman’s argument), then Goodman’s riddle is neither *new* nor peculiar to *induction*.
- Carnap’s *later* theories of  $c$  are sensitive to choice of language. But, (a) Goodman was not aware of those later theories, and (b) “grue” doesn’t reveal *that* problem anyway.
- In order to pinpoint the (pernicious) language-variance of Carnap’s later  $c$ -theories, more sophisticated constructions are required (*e.g.*, David-Miller-esque constructions).

## Is “Grue” an Observation Selection Effect? Part I

- **Canonical Example of an OSE:** I use a fishing net to capture samples of fish from various (randomly selected) parts of a lake. Let  $E$  be the claim that all of the sampled fish were over one foot in length. Let  $H$  be the hypothesis that all the fish in the lake are over one foot  $[(\forall x)((Fx \& Lx) \supset O_x)]$ .
- Intuitively, one might think  $E$  should evidentially support  $H$ . This may be so for an agent who knows *only* the above information ( $K$ ) about the observation process. That is, it seems plausible that  $\text{Pr}(E | H \& K) > \text{Pr}(E | \sim H \& K)$ , where  $\text{Pr}$  is taken to be “evidential” (or “epistemic”) probability.
- But, what if I *also* tell you that ( $D$ ) the net I used to sample the fish from the lake (which generated  $E$ ) has holes that are all over one foot in diameter? It seems that  $D$  *defeats* the support  $E$  provides for  $H$  (relative to  $K$ ), because  $D$  *ensures*  $O$ . Thus, intuitively,  $\text{Pr}(E | H \& D \& K) = \text{Pr}(E | \sim H \& D \& K)$ .

## Is “Grue” an Observation Selection Effect? Part II

- The “grue” hypothesis ( $H_2$ ) entails the following [& the green hypothesis ( $H_1$ ) entails a parallel claim  $H'_1$  about *grue* emeralds]: ( $H'_2$ ) All green emeralds have been (or will have been) examined prior to  $t$ .  $[(\forall x)((Ex \& Gx) \supset O_x)]$
- Now, consider the following two observation processes:
  - **Process 1.** For each green emerald in the universe, a slip of paper is created, on which is written a true description of that object as to whether it has property  $O$ . All the slips are placed in an urn, and one slip is sampled at random from the urn. By *this* process, we learn ( $\mathcal{E}$ ) that  $Ea \& Ga \& Oa$ .
  - **Process 2.** Suppose all the green emeralds in the universe are placed in an urn. We sample an emerald ( $a$ ) at random from this urn, and we examine it — *knowing antecedently* that the examination of  $a$  will take place prior to  $t$ , *i.e.*, that  $Oa$  is true. By *this* process, we learn ( $\mathcal{E}$ ) that  $Ea \& Ga \& Oa$ .
- Process 2 (which is Goodman-like) is uninformative wrt  $H'_1$  vs  $H'_2$ .