

# The Inexpressibility of Validity\*

Julien Murzi<sup>†</sup>

August 30, 2013

Tarski's Undefinability of Truth Theorem comes in two versions: that no consistent theory which interprets Robinson's Arithmetic (Q) can prove all instances of the T-Scheme

$$(T) \quad \text{Tr}(\ulcorner \phi \urcorner) \leftrightarrow \phi,$$

and hence *define* truth; and that no such theory, if sound, can even *express* truth.<sup>1</sup> In this note, I prove corresponding limitative results for *validity*. While Peano Arithmetic already has the resources to define a predicate expressing *logical validity*, as Jeff Ketland (2012) has recently pointed out, no theory which interprets Q closed under the standard structural rules can define nor express *validity*, on pain of triviality. The results put pressure on the widespread view that there is an *asymmetry* between truth and validity, viz. that while the former cannot be defined within the language, the latter can.<sup>2</sup> I argue that Vann McGee's and Hartry Field's arguments for the asymmetry view are problematic.

---

\*Many thanks to Jc Beall, Roy Cook, Hartry Field, Michael Glanzberg, Volker Halbach, Jeff Ketland, Hannes Leitgeb, Dave Ripley, Jack Woods, and audiences at the Reasoning Seminar, University of Kent, and at the Second Oxford-Buenos Aires Workshop on Truth, Inexpressibility and Paradoxes, Buenos Aires, for helpful comments and discussion. Special thanks to Luca Incurvati, Lionel Shapiro, and an anonymous referee for detailed comments on a previous draft that have led to significant improvements. I gratefully acknowledge the Alexander von Humboldt Foundation and the British Academy for generous financial support during the time this paper was written.

<sup>†</sup>University of Kent and Munich Center for Mathematical Philosophy, Ludwig-Maximilians Universität, München, j.murzi@gmail.com.

<sup>1</sup>Definitions of definability and expressibility will be provided in due course. For the time being, suffices to say that a theory defines truth iff it proves every instance of T, and that it expresses truth iff every such instance is true.

<sup>2</sup>See McGee (1991, pp. 44-5) and Field (2008). Field writes that while "[t]ruth can't possibly be given an extensionally correct definition within the language, ... validity presumably can" (Field, 2008, p. 127). What he means by that is that one can define the set of validities, without necessarily thereby defining the *concept* of validity. Kreisel's Squeezing Argument may then reassure us that standard model-theoretic definitions of first-order logical validity are extensionally adequate (Kreisel,

# 1 Logical validity and validity

We're all familiar with the standard Tarskian definition of logical validity:

(Tarski)  $\phi$  is logically valid iff  $\phi$  is true under all reinterpretations of its non-logical constituents.

Now let PA be an axiomatisation of Peano Arithmetic, with language  $\mathcal{L}_A$ . Idealising quite a bit, we may take PA to be a (simplified) model of our *maximal theory* at a certain time, i.e. a model of the closure of the set of sentences we (at least) implicitly accept at a certain time under the rules we (at least) implicitly accept at that time (Field, 1994, p. 401-5). We can then put our model to work, and ask: what can it say about logical validity?

Jeff Ketland (2012) has recently pointed out that PA already has the resources to derive the following principles about a predicate  $\text{Val}(x)$  intuitively reading ' $x$  is logically valid' (so that  $\text{Val}(n)$  is true iff  $n$  is the code of a valid  $\mathcal{L}_A$ -sentence):

(V-Intro) Given a *logical* derivation of  $\phi$ , infer  $\text{Val}(\ulcorner \phi \urcorner)$ ;<sup>3</sup>

(V-Out)  $\text{Val}(\ulcorner \phi \urcorner) \rightarrow \phi$ ;

(V-Imp)  $\text{Val}(\ulcorner \phi \rightarrow \psi \urcorner) \rightarrow (\text{Val}(\ulcorner \phi \urcorner) \rightarrow \text{Val}(\ulcorner \psi \urcorner))$ .<sup>4</sup>

The above principles can be easily generalised by means of a two-place predicate  $\text{Val}(x, y)$  expressing *argument-validity*, where  $x$  is to be replaced by (the code of) the conjunction of the premises of a given argument. Not only can PA, and hence our maximal theory, talk about numbers (and its own syntax): it can also talk, and prove intuitive principles, about *logical validity*.

To be sure, the extension of 'logically valid' will depend on one's view about what counts as logical. While arguably  $\wedge, \vee, \rightarrow, \neg, \exists, \forall$  and  $=$  are logical and terms such as 'water' aren't, the status of semantic predicates such as 'true' and 'valid' is less clear. According to Graham Priest (2007, p. 193), the T-Scheme (T) "ought ... to be considered part of logic", essentially on the grounds that its instances are analytic of 'true'. The view is controversial, however. As Roy Cook (2012, p. 235)

---

1967); see also Field (2008, Ch. 2.2). McGee's and Field's view express current logical orthodoxy, although the unorthodox view defended here, viz. that truth and validity are both paradox-prone semantic notions, is by no means new. For a representative sample of the literature, see e.g. Priest and Routley (1982), Read (1979), Read (2001) and, more recently, Shapiro (2011) and Beall and Murzi (2013).

<sup>3</sup>'Logical derivation' here means 'derivable in the logic of PA', viz. classical first-order logic.

<sup>4</sup>Here and throughout,  $\ulcorner \phi \urcorner$  indicates the numerical code of  $\phi$ , relative to some fixed coding scheme.

has recently observed, if “logical truth requires that uniform substitution instances of logical truths [be] logical truths”, some instances of (T) are not logically valid.

The issue, though, is largely terminological. Thus, McGee recommends that we use a broader notion of logical validity, while acknowledging that the notion doesn’t coincide with pure logical validity:

we must employ a richer notion of logical necessity, according to which there are certain sentences whose truth is so basic to our way of thinking and talking that they have the same epistemic status as logical validities, even though they are not actually logically valid. (McGee, 1991, p. 43)

In a similar spirit, Field suggests that we use “the term ‘valid’ ... in a very broad sense, one which counts ... a large amount of set theory and the basic principles of truth and satisfaction as valid” (Field, 2007, p. 99). I’ll distinguish, then, between *logical validity* and *validity (tout court)*. I mention two reasons why the distinction must be made.

First, it might be argued that there are clear examples of arguments that are valid, albeit not logically so. A first (admittedly controversial) example is given by the  $\omega$ -rule:

0 has property  $F$   
1 has property  $F$ ,  
2 has property  $F$ ,  
...  
Every number has property  $F$ .

Many would think that the rule is intuitively valid.<sup>5</sup> Yet, the rule is invalid in first-order logic.<sup>6,7</sup> Less controversially, other examples of valid but not logically valid arguments include analytic validities such as the following:

---

<sup>5</sup>Examples include Tarski (1983, p. 411), Etchemendy (1990) and McGee (2011, p. 33).

<sup>6</sup>First-order logic is *compact*: an argument is valid in first-order logic if and only if some *finite* sub-argument is valid.

<sup>7</sup>There are at least two ways of validating the  $\omega$ -rule. First, one could allow *infinitary quantifiers*, such as “there are infinitely many  $x$  such that ...”. Then,  $\{F(0), F(1), F(2), \dots\}$  would imply that there are infinitely many  $F$ s. The resulting system would be semantically incomplete, but it might be thought that this is not a problem. Relatedly, one could simply point out that the  $\omega$ -rule is valid in *second-order logic* (with standard semantics). Here I don’t have space to discuss either option, and simply observe that both are themselves controversial. For a criticism of infinitary quantifiers, see Hanson (1997, pp. 391-2).

$$\frac{x \text{ is a brother}}{x \text{ is male}} \quad \frac{\Phi(0) \quad \forall n(\Phi(n) \rightarrow \Phi(S(n)))}{\forall n(\Phi(n))},$$

where  $S(x)$  expresses the successor function. To be sure, one might point out that such rules would be logically valid if we held fixed the interpretation of ‘brother’, ‘male’, ‘successor’ and numerals. However, the dialectic here is a familiar one: intuitively invalid inferences such as ⟨Leslie was a US president  $\therefore$  Leslie was a man⟩, and intuitively invalid sentences such as ‘There are at least two numbers’, would thereby be declared logically valid (Etchemendy, 1990).<sup>8</sup> Whether rules such as

$$\frac{x \text{ is water}}{x \text{ is H}_2\text{O}}$$

are valid is perhaps less clear, but the examples, I hope, suffice to establish my point: logical validity is *one kind* of validity (see also Priest, 2006a).<sup>9</sup>

Second, as hinted in the above McGee quote, validity arguably has a role to play in our epistemic lives. Gil Harman (1986, p. 18) suggests that ordinary reasoning is partly governed by the following principles:

*Recognised Implication Principle.* One has a reason to believe  $\phi$  if one recognises that  $\phi$  is implied by one’s view.

*Recognised Inconsistency Principle.* One has a reason to avoid believing things one recognises to be inconsistent.

Similarly Field advocates the existence of a connection between validity and correct reasoning, this time framed in terms of degrees of belief. Where  $P(\phi)$  refers to one’s degrees of belief in  $\phi$ , Field’s principle reads:

(F) If it’s obvious that  $\phi_1, \dots, \phi_n$  together entail  $\psi$ , then one ought to impose the constraint that  $P(\psi)$  is to be at least  $P(\phi_1) + \dots + P(\phi_n) - (n - 1)$ , in any circumstance where  $\phi_1, \dots, \phi_n$  are in question. (Field, 2009, p. 259)

---

<sup>8</sup>Again, I don’t have space here to fully defend the claim that Tarski’s account of validity cannot handle analytic validities. This is a large issue related to the even larger issue whether, as forcefully argued by John Etchemendy (1990, 2008), Tarski’s account *undergenerates*. For more discussion on analytic validity and the issue of under generation, see e.g. Priest (1995, p. 288) and Etchemendy (2008, p. 278 and ff).

<sup>9</sup>For more on the slide between the foregoing examples and clear cases of logical validity, see e.g. Lycan (1989) and Sagi (2013).

Roughly: if  $\phi$  entails  $\psi$ , then one's degree of belief in  $\psi$  should be no lower than one's degree of belief in  $\phi$ .

Both Harman's and Field's principles are, I think, plausible. But neither principle is especially concerned with *logical validity*. Plainly, ordinary speakers typically don't distinguish between valid sentences and *logically* valid sentences (Harman, 1986, p. 17). Hence, if the principles apply at all, they apply to validity. As Harman puts it, "since there seems to be nothing special about *logical* implications and inconsistencies, ... there seems to be no significant way in which *logic* might be *specially* relevant to reasoning" (Harman, 2009, p. 334). Logical validity is not specially relevant to reasoning. But validity arguably is.

What, then, is validity? One standard view sees consequence as a kind of *modality*. For some understanding of 'possible', a set of premises entails a given conclusion iff it is impossible that all the premises are true and the conclusion false:

*Truth-preservation view.* An argument  $\langle \Gamma \therefore \beta \rangle$  is valid iff it is impossible that every member of  $\Gamma$  is true but  $\beta$  is false.

The view accommodates the examples of pp. 3-4, but has recently come under attack, chiefly on the grounds that the very claim that valid arguments preserve truth (i) entails Curry-driven triviality and (ii) is inconsistent with Gödel's Second Incompleteness Theorem (Priest, 2006a,b; Field, 2008, 2009; Beall, 2009).<sup>10</sup> However, the results to be presented below don't require that valid arguments be truth-preserving—they are indeed compatible with views that reject this assumption. One such view is endorsed by Field (2008, 2009). According to Field, validity is a primitive notion whose role is to constrain belief, and degrees of belief, via principles such as (F):

*Normative view.* validity normatively constrains our degrees of belief.

Now for some well-known, and less well-known, theorems.

## 2 Validity Curry

While V-Intro is adequate for a predicate expressing *logical validity*, it doesn't seem adequate for a predicate expressing *validity*. Suppose  $\phi$  has been derived by means

---

<sup>10</sup>For a critical discussion of Field's and Beall's arguments, see Shapiro (2011) and Murzi and Shapiro (forthcoming).

of rules that are themselves valid. Then, we would like to be able to say that  $\phi$  is *itself* valid. That is, the following rule should be in place:

(V-Intro\*) Given a valid derivation of  $\phi$ , infer  $\text{Val}(\ulcorner \phi \urcorner)$ .<sup>11</sup>

The rule is intuitively valid. On the *truth-preservation view*, it states that sentences proved only by means of necessarily truth-preserving rules are themselves necessary. On the *normative view*, it effectively says that, if we have proved  $\phi$  by means of valid rules, then, if we have thereby come to believe  $\phi$  to degree 1, then we should also believe  $\text{Val}(\ulcorner \phi \urcorner)$  to degree 1. Unlike V-Intro and V-Out, however, V-Intro\* and V-Out (with ‘Val( $x$ )’ now taken to express validity) spell trouble.

**Theorem 1** (Myhill 1960, Kaplan and Montague 1960). *Let  $T$  be a theory which (i) interprets  $\text{Q}$ , (ii) proves all instances of V-Out and (iii) is closed under V-Intro\*. Then,  $T$  is inconsistent.*

*Proof.* We reason in  $T$ . The Diagonal Lemma yields a sentence  $\kappa$  such that  $\kappa \leftrightarrow \neg \text{Val}(\ulcorner \kappa \urcorner)$ . We assume  $\text{Val}(\ulcorner \kappa \urcorner)$ , and derive  $\neg \text{Val}(\ulcorner \kappa \urcorner)$  by applications of V-Out and  $\leftrightarrow$ -E. Hence,  $\neg \text{Val}(\ulcorner \kappa \urcorner)$ . But, then,  $\kappa$  follows on no assumptions, and must therefore be valid. Contradiction.<sup>12</sup> □

In effect, the above reasoning—the Knower Paradox—is but a stronger Liar: it is the same paradox, except that the left-to-right direction of (T) is replaced by the weaker V-Intro\*.<sup>13</sup> Standard treatments of the Liar paradox still apply, however. On *paracomplete* treatments, one may not validly infer  $\neg \text{Val}(\ulcorner \kappa \urcorner)$  from a derivation of absurdity from  $\text{Val}(\ulcorner \kappa \urcorner)$ , i.e. the rule of *negation introduction* is rejected as (logically) invalid (see e.g. Kripke, 1975; Brady, 2006; Field, 2008; Horsten, 2009). On *paraconsistent* treatments, the above argument shows the existence of a true contradiction, viz.  $\text{Val}(\ulcorner \kappa \urcorner) \wedge \neg \text{Val}(\ulcorner \kappa \urcorner)$  (see e.g. Priest, 2006b; Beall, 2009).

<sup>11</sup>This is, in effect, a notational variant of the unrestricted Rule of Necessitation: given a derivation of  $\phi$ , infer  $\text{Tr}(\ulcorner \phi \urcorner)$ .

<sup>12</sup>The reasoning displayed in this proof is sometimes referred to as the Pseudo-Scotus Paradox. It might be objected that  $\text{Q}$ 's axioms and V-Intro\* are logically invalid. See e.g. Field (2008, p. 304 and p. 306), Ketland (2012) and Cook (2013). However, the assumption that they are valid is plausible on any interesting weakening of the notion of logical validity. For more discussion, see Murzi and Shapiro (forthcoming).

<sup>13</sup>To see this, note that both paradoxes have the same general form: from some  $\phi$  such that  $\vdash \phi \leftrightarrow \neg \Box(\ulcorner \phi \urcorner)$ , one proves  $\neg \Box(\ulcorner \phi \urcorner)$ , and hence  $\phi$ , on the assumption that  $\Box(\ulcorner \phi \urcorner)$ . Then,  $\Box(\ulcorner \phi \urcorner)$  follows on no assumptions via either the T-Scheme or Necessitation, depending on whether  $\Box$  is interpreted as, respectively, ‘true’ or ‘valid’.

Now suppose we'd like to attribute validity not only to sentences, but also to *arguments*. We'd then introduce in the language (at least) a two-place validity predicate  $\text{Val}(x, y)$  governed by the following rules:

(VP) Given a valid derivation of  $\psi$  from  $\phi$ , infer  $\text{Val}(\ulcorner \phi \urcorner, \ulcorner \psi \urcorner)$ .

(VD) From  $\phi$  and  $\text{Val}(\ulcorner \phi \urcorner, \ulcorner \psi \urcorner)$ , infer  $\psi$ .

More formally:

$$(VP) \frac{\phi \vdash \psi}{\vdash \text{Val}(\ulcorner \phi \urcorner, \ulcorner \psi \urcorner)} \quad (VD) \frac{\Gamma \vdash \text{Val}(\ulcorner \phi \urcorner, \ulcorner \psi \urcorner) \quad \Delta \vdash \phi}{\Gamma, \Delta \vdash \psi} .$$

VP is, again, intuitively valid. On the *truth-preservation view*, it states that, if  $\psi$  has been derived from  $\phi$  by means of truth-preserving rules, then the argument  $\langle \phi \therefore \psi \rangle$  is necessarily truth-preserving. On the *normative view*, it effectively says that, if  $\psi$  has been derived from  $\phi$  by means of valid rules, then the argument  $\langle \phi \therefore \psi \rangle$  is valid, and hence  $P(\phi) \leq P(\psi)$ . VD is also compelling on both views. On the truth-preservation view, it implies that, if we're in a position to assert that the argument  $\langle \phi \therefore \psi \rangle$  is truth-preserving, then we may infer  $\psi$  given  $\phi$ . On the normative view, it tells us that, if we're in a position to assert that the argument  $\langle \phi \therefore \psi \rangle$  is validity, then  $P(\phi) \leq P(\psi)$ .

Note that VP and VD are generalisations of, respectively, V-Intro\* and V-Out. To see this, it is sufficient to instantiate VP and VD using a constant  $\top$  expressing valid truth. Instantiating VP yields a notational variant of V-Intro\*, rewritten using our two place predicate  $\text{Val}(x, y)$  in place of a one-place validity predicate  $\text{Val}(x)$ :

$$(V\text{-Intro}^{**}) \frac{\top \vdash \psi}{\vdash \text{Val}(\ulcorner \top \urcorner, \ulcorner \psi \urcorner)} .$$

Likewise, instantiating VD thus

$$\frac{\Gamma \vdash \text{Val}(\ulcorner \top \urcorner, \ulcorner \psi \urcorner) \quad \top \vdash \top}{\Gamma, \top \vdash \psi}$$

yields a notational variant of a rule corresponding to the  $T$  axiom for a necessity operator, i.e. V-Out\*:

$$(V\text{-Out}^*) \text{Val}(\ulcorner \top \urcorner, \ulcorner \psi \urcorner), \top \vdash_{\top} \psi .$$

It is therefore no surprise that VP and VD *also* spell trouble. What is surprising is the minimal logical resources required to show this: one only needs to assume the validity of the standardly accepted *structural rules*, viz. Identity, Contraction and Cut:

$$(Id) \frac{}{\phi \vdash \phi} \quad (SContr) \frac{\Gamma, \phi, \phi \vdash \psi}{\Gamma, \phi \vdash \psi} \quad Cut \frac{\Gamma \vdash \phi \quad \Delta, \phi \vdash \psi}{\Gamma, \Delta \vdash \psi}.$$

The proof is a validity-involving version of Curry's Paradox: Validity Curry, or v-Curry, for short.<sup>14</sup>

**Theorem 2.** *Let  $T$  be any theory which interprets  $Q$  and is closed under VP, VD, Id and SContr. Then,  $T$  is trivial.*<sup>15</sup>

*Proof.* The Diagonal Lemma yields a sentence  $\pi$ , which intuitively says of itself, up to equivalence, that it validly entails that you will win the lottery:

$$\vdash_T \pi \leftrightarrow \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner).$$

Let  $\Sigma$  now be the following derivation of the further theorem  $\text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)$ :

$$\frac{\frac{\frac{\pi \vdash_T \pi \quad \vdash_T \pi \leftrightarrow \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)}{\pi \vdash_T \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)} \rightarrow\text{-E} \quad \pi \vdash_T \pi}{\frac{\pi, \pi \vdash_T \perp}{\pi \vdash_T \perp} \text{SContr}} \text{VD}}{\frac{\pi, \pi \vdash_T \perp}{\pi \vdash_T \perp} \text{SContr}} \text{VP}} \text{VP}$$

Using  $\Sigma$ , we can then 'prove' that you will win the lottery:

$$\frac{\frac{\frac{\Sigma}{\vdash_T \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)} \quad \frac{\frac{\vdash_T \pi \leftrightarrow \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)}{\vdash_T \pi} \quad \frac{\frac{\Sigma}{\vdash_T \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)}}{\vdash_T \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)}}{\vdash_T \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)}} \rightarrow\text{-E}}{\vdash_T \perp} \text{VD}}{\vdash_T \perp} \text{VD}} \text{VD}$$

□

In short: validity, just like truth, is plagued by paradox. More precisely, since VD and VD are generalisations of the validity rules that yield the Knower Paradox, the v-Curry Paradox is a generalisation of the Knower Paradox, which, recall, is but a stronger Liar.

★★ *Parenthetical note.* Theorem 2 can be strengthened. While the theorem assumes that  $Q$  be valid in the target sense, this assumption is not strictly needed, as Cook (2013) has recently pointed out.

<sup>14</sup>To the best of my knowledge, the first known occurrence of v-Curry paradox is in the 16th-century author Jean de Celaya. See Read (2001, fn. 11-12) and references therein. Albert of Saxony discusses a contrapositive version of the paradox in his *Insolubles* (Read, 2010, p. 211). More recent versions surface in Whittle (2004, fn. 3), Clark (2007, pp. 234-5) and Shapiro (2011, fn. 29). For a first comprehensive discussion of the v-Curry Paradox, see Beall and Murzi (2013).

<sup>15</sup>It is worth noting that Cut is effectively built in our formulation of VD. Hence, should the latter rule be formulated differently, Cut may need to be mentioned alongside Id and SContr in the statement of Theorem 2 below.



**Theorem 3** (Cook). *If VP and VD are valid, Q is trivial.*

*Proof.* Let  $Q^*$  be the conjunction of  $Q$ 's axioms. Using the Diagonal Lemma, we can derive

$$Q^* \vdash \kappa \leftrightarrow \text{Val}(\ulcorner \kappa \wedge Q^* \rightarrow \perp \urcorner)$$

We then 'prove'  $\text{Val}(\ulcorner \kappa \wedge Q^* \rightarrow \perp \urcorner)$  and, from this, conclude  $Q^* \vdash \perp$  □

Beyond Id, SContr and Cut,<sup>16</sup> all is needed for deriving the Validity Curry is that VP and VD be valid. It immediately follows from Theorem 2 and 3 that theories of semantic paradox which retain the standard structural rules—including the standard paracomplete and paraconsistent ones—still validate the v-Curry Paradox. Such theories can in general express *sentence* validity, but they're unable to express *argument* validity. For more discussion, see Zardini (2011), Beall and Murzi (2013) and Murzi (2012). *End note.* \*\*

### 3 Validity and expressibility

The results of §2 suggest natural generalisations of Tarski's undefinability and inexpressibility of truth theorems. I'll closely follow, *mutatis mutandis*, Peter Smith's presentation of Tarski's original results (Smith, 2007, pp. 180-2). As above, we'll take some (possibly not recursively axiomatisable) theory  $T$  which interprets PA to represent our maximal theory at a certain time, and hence the consequence relation of our language at that time. That is, ignoring for simplicity's sake the relativisation to time, we'll effectively assume that  $T$ 's consequence relation is *the* consequence relation of our maximal theory, so that  $T$  is, so to speak, validity *sound and complete*. In short:  $\phi \vdash_T \psi$  iff  $\phi$  entails  $\psi$ .

We can now show that  $T$  can neither define, nor express, validity. To this end, we make use of the two following (standard) definitions:

**Definition 4** (Definability).  $\Phi(x, y)$  *defines* a binary relation  $R$  in  $T$  iff:  $mRn$  iff  $\vdash_T \Phi(\bar{m}, \bar{n})$ , where  $\bar{m}$  and  $\bar{n}$  are names of, respectively,  $m$  and  $n$ .<sup>17</sup>

**Definition 5** (Expressibility).  $\Phi(x, y)$  *expresses* a binary relation  $R$  iff:  $mRn$  iff  $\Phi(\bar{m}, \bar{n})$  is true.

<sup>16</sup>This last rule is in effect built into VD. See *infra* fn. 15.

<sup>17</sup>Definability, as defined in Definition 4, is sometimes also referred to as *weak representability*.

Now let  $Valid$  be a numerical relation such that  $Valid(m, n)$  is true iff  $m$  and  $n$  are the Gödel numbers of formulae  $\phi, \psi$  of  $\mathcal{L}_A$  such that  $\phi$  entails  $\psi$ . Suppose an open wff  $Val(x, y)$  belonging to an arithmetical language  $\mathcal{L}'_A$  including  $\mathcal{L}_A$  expresses this numerical relation  $Valid$ . Then, for any  $\mathcal{L}_A$  sentences  $\phi, \psi$ ,  $\phi$  entails  $\psi$  iff  $Valid(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$  iff  $Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$  is true. That is, for any  $\mathcal{L}_A$  sentences  $\phi, \psi$ ,  $\phi$  entails  $\psi$  iff  $Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$  is true. In turn, this motivates the following definitions.

**Definition 6** (Formal validity predicate). An open  $\mathcal{L}'_A$ -wff  $Val(x, y)$  is a *formal validity-predicate* for  $\mathcal{L}_A$  iff  $Val(x, y)$  expresses validity, i.e. iff for every  $\mathcal{L}_A$ -sentence  $\phi, \psi$ ,  $\phi$  entails  $\psi$  iff  $Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$  is true.

**Definition 7** (Validity theory). A theory  $T$  (with language  $\mathcal{L}'_A$  which includes  $\mathcal{L}_A$ ) is a *validity theory* for  $\mathcal{L}_A$  iff, for some  $\mathcal{L}'_A$ -wff  $Val(x, y)$ ,  $\phi$  entails  $\psi$  iff  $\vdash_T Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$ . A validity theory for  $\mathcal{L}_A$  is also a *validity definition* for  $\mathcal{L}_A$ .

Note that, given our idealising assumption that  $T$  is validity sound and complete, it follows from Definitions 5 and 6 that  $Val(x, y)$  is a formal validity predicate iff  $Val(x, y)$  expresses  $T$ 's consequence relation. More formally,  $Val(x, y)$  is a formal validity predicate iff:  $Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$  is true iff  $\phi \vdash_T \psi$ . Furthermore, on the foregoing assumption,  $T$  is a validity theory iff the following holds:  $\phi \vdash_T \psi$  iff  $\vdash_T Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)$ . That is, if  $T$  is validity sound and complete,  $T$  is a validity theory iff it is closed under the naïve validity rules:

$$(VP) \frac{\phi \vdash_T \psi}{\vdash_T Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner)} \quad (VD) \frac{\Gamma \vdash_T Val(\ulcorner\phi\urcorner, \ulcorner\psi\urcorner) \quad \Delta \vdash_T \phi}{\Gamma, \Delta \vdash_T \psi}.$$

We're now ready to introduce our main results.

Can  $T$  define its own consequence relation, i.e. can  $T$  contain its own validity definition? This question is answered in the negative by the following theorem.

**Theorem 8** (Indefinability of  $\vdash_T$  in  $T$ ). *Let  $T$  be any non-trivial theory which interprets  $Q$  with language  $\mathcal{L}_V$  including a fresh predicate  $Val(x, y)$ . Then,  $T$  cannot define its own consequence relation.*

*Proof.* Since  $T$  is at least as strong as  $Q$ , the Diagonal Lemma applies, and gives us a sentence  $\pi$ —a v-Curry sentence!—such that

$$\vdash_T \pi \leftrightarrow Val(\ulcorner\pi\urcorner, \ulcorner\perp\urcorner).$$

However, since  $T$  defines its own consequence relation,  $T$  validates VP and VD. Hence, we can run the v-Curry reasoning in  $T$ , and conclude  $\vdash_T \perp$ , which contradicts our assumption that  $T$  isn't trivial.  $\square$

It immediately follows that, if  $T$  is validity sound and complete, then  $T$  cannot define *validity*.

**Corollary 9** (Indefinability of validity). *Let  $T$  be as above. Then, if  $T$  is validity sound and complete,  $T$  cannot define validity.*

*Proof.* This follows at once from Theorem 8 and the observation that, if  $T$  is validity sound and complete,  $T$  defines validity iff it validates VP and VD.  $\square$

Can at least  $T$  express validity? This question must also receive a negative answer, provided Q is sound. The proof makes use in the metalanguage of disquotational truth-predicate, i.e. we assume that, for all  $\phi \in \mathcal{L}_V$ ,  $\phi$  is true iff  $\phi$ , where ‘true’  $\notin \mathcal{L}_V$ .<sup>18</sup> It also assumes that our meta-theory validates the standard structural rules, and that the metalanguage entailment connective we’ve been using so far satisfies (a version of) *conditional proof* and *modus ponens*, i.e. that, for all  $\phi \in \mathcal{L}_V$ , if there is a derivation of  $\psi$  from  $\phi$  in the metatheory, then  $\phi$  entails  $\psi$ , and if  $\phi$  and  $\phi$  entails  $\psi$ , then  $\psi$ , where ‘entails’  $\notin \mathcal{L}_V$ .

**Theorem 10** (Inexpressibility of validity). *Let  $T$  be any non-trivial theory which interprets Q, with language  $\mathcal{L}_V$ . Then, if Q is sound,  $T$  cannot contain a predicate expressing validity.*

*Proof.* Let  $T$  be a theory that interprets Q, and suppose there is a  $\mathcal{L}_V$  predicate  $\text{Val}(x, y)$  expressing validity. We then have:

1.  $\vdash_T \pi \leftrightarrow \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)$ .

However, (the subtheory of  $T$  which interprets) Q is *ex hypothesi* sound. So

2.  $\pi \leftrightarrow \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)$ .

Since  $\text{Val}(x, y)$  expresses validity, the following holds:

3.  $\pi$  entails  $\perp$  iff  $\text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner)$  is true.

We can now reason the standard Curry way. We assume  $\pi$ , and derive that  $\pi$  entails  $\perp$  from 2. and 3. Assuming again  $\pi$ , we derive  $\perp$  and, discharging both occurrences of  $\pi$ , we deduce that  $\pi$  entails  $\perp$ . From 2. and 3., we conclude  $\perp$ .  $\square$

---

<sup>18</sup>While I’m assuming the metalanguage to be classical, and hence take the truth-predicate to be typed, the proof would also go through using a type-free truth-predicate in a non-classical metalanguage.

Notice that the multiple discharge of  $\pi$  is effectively equivalent to assuming that SContr holds (see e.g. Negri and von Plato, 2001, Ch. 8).

We're now confronted with a familiar dilemma: either we invalidate one of the naïve validity principles at work in the foregoing proofs, or, if  $T$  is to either define or express validity,  $T$ 's logic must be (radically) weakened: one among Id, SContr and Cut must go.

## 4 Incompleteness and classical validity

It might be objected that the foregoing results are inconsistent with both Gödel's Second Incompleteness Theorem and Löb's Theorem and that, for this reasons, our definitions of a formal validity predicate and of a validity theory must be incorrect. The Second Incompleteness Theorem states that no consistent recursively axiomatised theory  $T$  which interprets Q and is strong enough to prove the Hilbert-Bernays conditions for a predicate  $Prov_T$  expressing provability-in- $T$  can prove its own consistency.<sup>19</sup> Löb's Theorem is a formalised version of Curry's Paradox, which, as it turns out, entails the Second Incompleteness Theorem. Where  $T$  is as above, Löb's Theorem states that, if  $\vdash_T Prov_T(\ulcorner \phi \urcorner) \rightarrow \phi$ , then  $\vdash_T \phi$ .

In a nutshell, one might use Gödel's and Löb's results to argue against VD thus. Consider the following version of the Second Incompleteness Theorem.

**Theorem 11.** *Let  $T$  be any recursively axiomatised theory that interprets Q, and let  $Prov_T(x)$  be a provability predicate for  $T$ . Suppose  $T$  asserts that theorems of  $T$  are valid and that validity is factive:*

1.  $Prov_T(\ulcorner \phi \urcorner) \rightarrow Val(\ulcorner \phi \urcorner)$ ;
2.  $Val(\ulcorner \phi \urcorner) \rightarrow \phi$ .

*It follows by Löb's Theorem that  $T$  is trivial.*

*Proof.* Suppose  $T$  derives both 1. and 2. Then, since  $Prov_T(\ulcorner \phi \urcorner) \rightarrow \phi$  follows from 1. and 2. by the transitivity of  $\rightarrow$ ,  $T$  also derives  $Prov_T(\ulcorner \phi \urcorner) \rightarrow \phi$ . By Löb's Theorem,  $T$  derives  $\phi$ . □

That this is in effect the Second Incompleteness Theorem can be seen by contraposing the proof while setting  $\phi := 0 \neq 1$ . We can now derive the following corollary.

---

<sup>19</sup>The derivability-conditions are the predicate-analogues of the Rule of Necessitation (see *infra*, fn. 11) and of the K and 4 axioms of modal logic.

**Corollary 12.** *Let  $T$  be as above and suppose  $T$  is closed under VD. Then,  $T$  is trivial.*

*Proof.* One need only notice that VD entails V-Out (see §2). Then, the result immediately follows from Theorem 10. □

Have we *shown* that VD must be rejected? I think not, but let's first examine the main assumption on which the arguments depends.

The argument assumes that our theory of validity  $T$  is *recursively axiomatisable*. McGee defends the assumption thus:

[i]f ... we identify the [validities] as those statements from whose denials one can derive a contradiction, and if we take derivability to be provability is some explicitly describable system of rules, it will follow that the set of necessary truths is a recursively enumerable set. (McGee, 1991, pp. 44-5)

In short: if  $\text{Val}(\ulcorner \phi \urcorner)$  is true iff  $\neg\phi \vdash_T \perp$ , for some recursively axiomatisable  $T$ , then the set of validities is recursively enumerable. It then follows from Corollary 12 that VD must fail. Moreover, as McGee observes, "if the set [of validities] is recursively enumerable ... [then we can] explicitly define it" (McGee, 1991, p. 45). That is, there is, in McGee's view, a fundamental *asymmetry* between truth and validity: unlike truth, validity *is* definable in the language, and "[w]e must confront the paradox [the Knower Paradox] directly; there is no escaping in the metalanguage" (McGee, 1991, p. 45). Yet, it is doubtful that the claim that validity is recursively axiomatisable can be made plausible in the present context. For one thing, the claim is not available to someone who accepts the validity of arguments with infinitely many premises, such as the example of §2.<sup>20</sup> For another, McGee's argument for the recursive axiomatisability of validity fails to convince. The argument assumes that, if  $\phi$  is valid, then we can *derive* a contradiction from  $\neg\phi$ . But since to derive  $\perp$  from  $\neg\phi$  is to (classically) *prove*  $\phi$ , this is tantamount to assuming a form of *verificationism*, to the effect that validities must have proofs of a certain kind. This is implausible, however.

To begin with, Goldbach's Conjecture could be true but (absolutely) unprovable. It would then follow that, since one wouldn't be able to *derive*  $\perp$  from the negation of Goldbach's Conjecture, Goldbach's Conjecture, unlike, say, Fermat's Theorem, or some other provable arithmetical theorem, would be invalid in the target sense. This

---

<sup>20</sup>Incidentally, McGee concedes that instances of the *omega*-rule are at least informally valid (McGee, 2011, p. 33).

seems counterintuitive, though. Since both Goldbach’s Conjecture and Fermat’s Theorem are arithmetical sentences, it would seem that either they are *both* valid, or neither is.<sup>21</sup> Second, even if all mathematical truths are absolutely provable, the set of absolutely provable sentences would still not be guaranteed to be recursively enumerable. For example, perhaps every mathematical truth is absolutely provable by some possible finite being or other, even if no possible finite being can give absolute proofs of all mathematical truths.<sup>22</sup>

In any event, the claim that validity is recursively axiomatisable is itself problematic. The claim implies that  $\text{Val}(x)$  can be interpreted as  $\text{Prov}_T(x)$ , which in turn implies, because of Löb’s Theorem, that our theory of validity cannot assert that valid sentences are true, on pain of triviality. Yet we would seem to *know* that validity is a mode of truth. As William Reinhardt puts it in the case of knowledge, “we know that what is known is true, but the reason for this is that if it were not so we would not call it known” (Reinhardt, 1986, pp. 468-9). The point, I take it, carries over to validity.

McGee (1991, pp. 45-9) confronts this second issue directly. While discussing what he calls *logical necessity*, i.e. what we are simply calling *validity*, he concedes that the principles which generate the Knower Paradox are “intuitively obvious”. In particular, he acknowledges that it is natural to think, of someone who denies the necessity of all instances of the schema ‘If  $\phi$  is necessary, then  $\phi$ ’ (i.e. V-Out), that he “would not know what he was talking about[:] even though he uses the word ‘necessary’, he could not be talking about necessity; he must be using the word in a deviant way” (p. 48). Still, McGee’s recommendation is that we substitute V-Out with the weaker, but consistent

$$(L) \text{Val}(\ulcorner \text{Val}(\ulcorner \phi \urcorner) \rightarrow \phi \urcorner) \rightarrow \text{Val}(\ulcorner \phi \urcorner)),$$

which, he claims, “our dogmatic insistence upon V-Out has obscured to view” (p. 49). Validity can be seen to be non-paradoxical, McGee suggests, provided we give up some “intuitively obvious” principles about validity, such as the truism that valid sentences are true.

This reasoning also fails to convince. In effect, McGee is here merging together two different issues: the question whether validity is paradoxical, and the problem how the validity paradoxes should be solved. While it may be that V-Out has

---

<sup>21</sup> Admittedly, this objection is more plausible when validity is equated with necessary preservation of truth. Thanks to Jack Woods for helping me appreciate this point.

<sup>22</sup> Thanks to an anonymous referee for suggesting this point.

obscured (L) to view, the point remains that V-Out *is* intuitively obvious: more obvious, I submit, than the assumption that validity is recursively axiomatisable. We may eventually revert to a recursively axiomatisable theory of validity, as McGee suggests, and invalidate V-Out. But this would be a *reaction* to the validity paradoxes: the recursive axiomatisability of the set of validities, and the subsequent invalidation of V-Out, would not be one of our initial assumptions.

It might be insisted that validity cannot be paradoxical on the grounds that “the notion of validity is to be ... defined in set-theory” (Field, 2008, p. 298) and that, since set-theory is consistent, so must be *validity*. This argument clearly would not work, however. For if it were legitimate to assume that validity is model-theoretically definable in order to show that there are no paradoxes of validity, then it would also be legitimate to assume that *truth* is model-theoretically definable in order to show that there are no paradoxes of truth. That truth-in- $\mathcal{L}$  and validity $_{\mathcal{L}}$  can be defined in set-theory by no means imply that there are no paradoxes of *truth* and *validity*.

Still, it might be thought that there are independent reasons to assume that validity is *classical*, which again would imply that validity cannot be a paradoxical notion. To see this, notice that, if validity is classical, then the following disjunction would hold:

$$(D) \pi \wedge \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner) \vee \neg \pi \wedge \neg \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner).$$

It is now easy to check that both disjuncts are incompatible with VD. The second entails

$$(D^*) \neg \text{Val}(\ulcorner \text{Val}(\ulcorner \pi \urcorner, \ulcorner \perp \urcorner) \wedge \pi \urcorner, \ulcorner \perp \urcorner),$$

which effectively says that an instance of VD is invalid. As for the first, given VD, it immediately entails  $\perp$ . Field (2008, p. 307) mentions two reasons why we should think that validity is classical:

- (a) that assuming excluded middle for validity claims leads to simpler reasoning about validity;

and

- (b) that it would seem to be somewhat detrimental to the role of logic as regulator of reasoning if we were unable to say that any given piece of reasoning is either valid or not valid.

He concludes that “these considerations together seem to ... make the assumption that validity claims obey excluded middle a reasonable working hypothesis” (Field, 2008, p. 307). Both (a) and (b), though, validate parallel arguments against the claim that *truth* is paradoxical. *Ad (i)*, if logic is a guide for correct reasoning and if, as Field supposes, only a simple logic can serve such a purpose, then reasoning in classical or intuitionistic logic is arguably simpler than reasoning in any weakening of these logics. But it would be too quick to conclude that one direction of the T-Scheme should fail on the grounds that assuming excluded middle for truth claims leads to simpler reasoning about truth. Certainly this argument would not be available to Field, who has long been advocating a non-classical, paracomplete logic of paradox (Field, 2003, 2007, 2008). *Ad (ii)*, a parallel argument would conclude that it would be detrimental to the role of truth as a regulator of assertion if we were unable to say that any given sentence is either true or not true. Either way, it would then follow that *truth* must be a classical notion and that, for this reason, at least one direction of the T-Scheme must fail. But this would hardly be a reason for thinking that the Liar reasoning isn’t paradoxical.

## 5 Concluding remarks

I’ve argued that we have reasons—largely epistemic ones—for countenancing a broader notion of validity, alongside logical validity. Such a notion is no more, and *no less*, paradoxical than truth. As John Myhill puts it, commenting on the notion of absolute provability (a kind of validity):

the situation is completely analogous to that Epimenes paradox which arises when we try to formalize the notion of truth, and does not show the notion of [absolute] provability to be any more paradoxical than the notion of truth. (Myhill, 1960, p. 470)

The paradoxes of validity are *generalisations* of the truth-theoretic paradoxes. The Knower Paradox is but a stronger Liar, and v-Curry is, in turn, a generalisation of the Knower, one that employs stronger, though equally compelling, validity principles, as well as weaker logical resources. Consequently, validity gives rise to stronger indefinability and inexpressibility results: the standardly assumed structure of the validity relation suffices to make validity indefinable and, if Q is sound, inexpressible.



## References

- Beall, J.: 2009, *Spandrels of Truth*, Oxford University Press, Oxford.
- Beall, J. and Murzi, J.: 2013, Two flavors of Curry's paradox, *The Journal of Philosophy* **CX**(3), 143–65.
- Brady, R.: 2006, *Universal Logic*, CSLI Publications, Stanford.
- Clark, M.: 2007, *Paradoxes from A to Z*, second edn, Routledge, London and New York.
- Cook, R.: 2012, The T-schema is not a logical truth, *Analysis* **72**(2), 231–239.
- Cook, R.: 2013, There is no paradox of logical validity. Forthcoming in *Logica Universalis*.
- Corcoran, J. (ed.): 1983, *Logic, Semantics, Metamathematics*, Second edn, Hackett Publishing Company, Indianapolis.
- Etchemendy, J.: 1990, *The Concept of Logical Consequence*, Harvard University Press, Harvard (Mass.).
- Etchemendy, J.: 2008, Reflections on consequence, in D. Patterson (ed.), *New Essays on Tarski and Philosophy*, Oxford University Press, pp. 263–99.
- Field, H.: 1994, Are our logical and mathematical concepts highly indeterminate?, in P. A. French, T. E. Uehling, and H. K. Wettstein (eds), *Midwest Studies in Philosophy*, Vol. 19, pp. 391–429.
- Field, H.: 2003, A revenge-immune solution to the semantic paradoxes, *Journal of Philosophical Logic* **32**, 139–77.
- Field, H.: 2007, Solving the paradoxes, escaping revenge, in J. Beall (ed.), *Revenge of the Liar*, Oxford University Press, pp. 53–144.
- Field, H.: 2008, *Saving Truth from Paradox*, Oxford University Press, Oxford.
- Field, H.: 2009, What is the normative role of logic?, *Proceedings of the Aristotelian Society* **83**, 251–68.
- Hanson, W. H.: 1997, The concept of logical consequence, *The Philosophical Review* **106**(3), 365–409.

- Harman, G.: 1986, *Change in View. Principles of Reasoning*, MIT Press, Cambridge (Mass.).
- Harman, G.: 2009, Field on the normative role of logic, *Proceedings of the Aristotelian Society* **109**(1), 333–335.
- Horsten, L.: 2009, Levity, *Mind* **118**(471), 555–581.
- Kaplan, D. and Montague, R.: 1960, A paradox regained, *Notre Dame Journal of Formal Logic* **1**, 79–90.
- Ketland, J.: 2012, Validity as a primitive. Forthcoming in *Analysis*.
- Kreisel, G.: 1967, Informal rigour and completeness proofs, in I. Lakatos (ed.), *Problems in the Philosophy of Mathematics*, North-Holland, Amsterdam, pp. 138–171.
- Kripke, S.: 1975, Outline of a theory of truth, *Journal of Philosophy* **72**, 690–716.
- Lycan, W.: 1989, Logical constants and the glory of truth-conditional semantics, *Notre Dame Journal of Formal Logic* **30**(390-400).
- McGee, V.: 1991, *Truth, Vagueness, and Paradox*, Hackett Publishing Company, Indianapolis.
- McGee, V.: 2011, Logical consequence, in L. Horsten and R. Pettigrew (eds), *The Continuum Companion to Philosophical Logic*, Continuum, London and New York.
- Murzi, J.: 2012, On Heck's new liar, *Thought* **1**(4), 258–69.
- Murzi, J. and Shapiro, L.: forthcoming, Validity and truth-preservation, in H. G. T. Achourioti F. Fujimoto and J. Martinez-Fernandez (eds), *Unifying the Philosophy of Truth*, Springer.
- Myhill, J.: 1960, Some remarks on the notion of proof, *Journal of Philosophy* **57**(14), 461–71.
- Negri, S. and von Plato, J.: 2001, *Structural Proof Theory*, Cambridge University Press, Cambridge.
- Priest, G.: 1995, Etchemendy on logical consequence, *Canadian Journal of Philosophy* **25**(2), 283–92.

- Priest, G.: 2006a, *Doubt Truth to be a Liar*, Oxford University Press, Oxford.
- Priest, G.: 2006b, *In Contradiction*, Oxford University Press, Oxford. Expanded edition (first published 1987 Kluwer-Dordrecht).
- Priest, G.: 2007, Parconsistency and dialetheism, in D. M. Gabbay and J. Woods (eds), *Handbook of the History of Logic Vol 8: The Nonmonotonic Turn in Logic*, North Holland, Amsterdam, pp. 129–204.
- Priest, G. and Routley, R.: 1982, Lessons from Pseudo-Scotus, *Philosophical Studies* **42**, 189–99.
- Read, S.: 1979, Self-reference and validity, *Synthese* **42**, 265–74.
- Read, S.: 2001, Self-reference and validity revisited, in M. Yrjonsuuri (ed.), *Medieval Formal Logic*, Kluwer Academic Publishers, pp. 183–96.
- Read, S.: 2010, Field’s paradox and its medieval solution, *History and Philosophy of Logic* **31**(2), 161–176.
- Reinhardt, W. N.: 1986, Some remarks on extending and interpreting theories, with a partial predicate for truth, *Journal of Philosophical Logic* **15**, 219–51.
- Sagi, G.: 2013, Formality in logic: From logical terms to semantic constraints. Forthcoming in *Logique et Analyse*.
- Shapiro, L.: 2011, Deflating logical consequence, *The Philosophical Quarterly* **61**, 320–42.
- Smith, P.: 2007, *An Introduction to Gödel’s Theorems*, Cambridge University Press, Cambridge.
- Tarski, A.: 1983, *On the Concept of Logical Consequence*, Hackett Publishing Company, pp. 409–20. In Corcoran (1983).
- Whittle, B.: 2004, Dialetheism, logical consequence and hierarchy, *Analysis* **64**(4), 318–26.
- Zardini, E.: 2011, Truth without contra(di)ction, *Review of Symbolic Logic* **4**, 498–535.