

Bounding Nonsplitting Enumeration Degrees

Thomas F. Kent
Andrea Sorbi

Università degli Studi di Siena
Italia

July 18, 2007



Goal: Introduce a form of Σ_2^0 -permitting for the enumeration degrees.

Till now, density was the only known property that held in all ideals of Σ_2^0 -enumeration degrees.

A is enumeration reducible to B ($A \leq_e B$) if we can enumerate A given any enumeration of B .

Definition

$A \leq_e B$ iff there is c.e. set Φ such that

$$A = \{x : \exists \langle x, P \rangle \in \Phi (P \text{ finite and } P \subseteq B)\} = \Phi^B$$



Basic Facts

We can embed the Turing degrees into the enumeration degrees via the embedding $\iota : \text{deg}_T(A) \mapsto \text{deg}_e(A \oplus \bar{A})$.

- The image of the Turing degrees under ι is known as the “total degrees”.

$$\mathbf{0}_e = \{W : W \text{ is c.e.}\}.$$

$$\mathbf{0}'_e = \text{deg}_e(\bar{K}).$$


Theorem (Cooper, 1984)

A is Σ_2^0 iff $A \leq_e \bar{K}$.

Theorem (Cooper, 1984)

The Σ_2^0 -enumeration degrees are dense.

The Global and Local Picture



Nonsplitting Degrees

Definition

A degree \mathbf{a} is nonsplitting if $\mathbf{a} > \mathbf{0}_e$ and for every $\mathbf{x}, \mathbf{y} < \mathbf{a}$,
 $\mathbf{x} \vee \mathbf{y} < \mathbf{a}$.

Theorem (Ahmad 1989 (c.f. Ahmad, Lachlan 1998))

There exists a nonsplitting Σ_2^0 -enumeration degree.

The requirements:

- Nontrivial

\mathcal{N}_Φ : $A \neq \Phi$, and

- Nonsplitting

$\mathcal{S}_{\Psi, \Omega_0, \Omega_1}$: $A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow \exists \Gamma_0, \Gamma_1 [A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A}]$.



Bounding Nonsplitting Degrees

Theorem (Kent, Sorbi 2007)

Every nontrivial Σ_2^0 -enumeration degree bounds a nonsplitting degree.

The requirements:

- $A \leq_e B$

$$\mathcal{R} \quad : \quad A = \Theta^B$$

- Nontrivial

$$\mathcal{N}_\Phi \quad : \quad A = \Phi \Rightarrow \exists \Delta (B = \Delta), \text{ and}$$

- Nonsplitting

$$\mathcal{S}_{\Psi, \Omega_0, \Omega_1} \quad : \quad A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow \exists \Gamma_0, \Gamma_1 \left[A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \right] \\ \text{or } \exists \Lambda [B = \Lambda].$$



Some Corollaries

Corollary

The nonsplitting degrees are downwards dense in the Δ_2^0 -enumeration degrees.

Corollary

There is a properly Σ_2^0 nonsplitting enumeration degree.

Corollary


The c.e. Turing degrees are not elementarily equivalent to any ideal of the Σ_2^0 -enumeration degrees.

Question

Are the nonsplitting degrees dense in the Σ_2^0 or Δ_2^0 enumeration degrees?




N-Requirement - Standard



1. Pick x and set $x \in A$.
2. If ever $x \in \Phi$, set $x \notin A$.




N-Requirement - Bounded



1. Assume $x_0, \dots, x_{n-1} \in \Phi \cap A$.
2. Pick x_n .
3. While 1. holds, enumerate $\langle x_n, B \upharpoonright x_n \rangle \in \Theta$.
4. If ever $x_n \in \Phi$, stop defining x_n axioms, and enumerate $D_n = \bigcap \{D : \langle x_n, D \rangle \in \Theta\}$ into Δ .




N-Requirement - Bounded



- If infinitely many x_i are defined and each $x_i \in A$, then since $D_0 \subseteq D_1 \subseteq \dots \subseteq B$, we can conclude $\Delta = B$.
- If $x_i \notin A$ then $x_j \notin A$ for all $j > i$.
- (Conditional Dumping) While $x_i \in A$, for all $y \in S(s_i)$, enumerate $\langle y, B \upharpoonright y \rangle$ into Θ .




S-Requirement - Standard



1. Pick x and set $x \in A$.
2. Wait for $x \in \Psi^{\Omega_0^A \oplus \Omega_1^A}$ via $\langle x, F_0 \oplus F_1 \rangle \in \Psi$.
3. Enumerate $\langle x, F_i \rangle$ into Γ_i , x into $S(\gamma_0)$ and return to Step 1.
 - Hopefully $x \in A$ iff $F_0 \subseteq \Omega_0^A$.
 - If true for co-finitely many x , then $A = \Gamma_0^{\Omega_0^A}$.
 - Strategies below γ_0 can only use x which have this property.




S-Requirement - Standard



4. If ever see $x \notin A$ and $F_0 \subseteq \Omega_0^A$ (hence $x \in \Gamma_0^{\Omega_0^A} - A$), dump $S(\gamma_0) - \{x\}$ into A , enumerate x into $S(\gamma_1)$.
- For this x , $F_0 \subseteq \Omega_0^{A - \{x\}}$, killing Γ_0 .
 - Hopefully $x \in A$ iff $F_1 \subseteq \Omega_1^A$.
 - If true for infinitely many x , then $A = \Gamma_1^{\Omega_1^A}$.
 - Strategies below γ_1 can only use x which have this property.



S-Requirement - Standard




5. If ever see $x \notin A$ and $F_1 \subseteq \Omega_1^A$ (hence $x \in \Gamma_1^{\Omega_1^A} - A$), dump $S(\gamma_1) \cup S(\gamma_0) - x$ into A , and set $x \notin A$.
- For this x , $F_1 \subseteq \Omega_1^{A - \{x\}}$, killing Γ_1 .
 - Not a problem since now $x \in \Psi^{\Omega_0^{A - \{x\}} \oplus \Omega_1^{A - \{x\}}}$.



S-Requirement - Bounded (v. 1.0)

$$S_{\Psi, \Omega_0, \Omega_1} : A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \text{ or } B = \Lambda$$




- As with the N -strategy, expand the outcome s to s_0, s_1, \dots
- If we choose x_0, x_1, \dots as possible diagonalization witness, and for all $i, x_i \in \Psi^{\Omega_0^A \oplus \Omega_1^A} \cap A$, then $B = \Lambda$.
- (Conditional Dumping) While $x_i \in A$, for all $y \in S(s_i)$, enumerate $\langle y, B \upharpoonright y \rangle$ into Θ .



S-Requirement - Potential Problem

$$S_{\Psi, \Omega_0, \Omega_1} : A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \text{ or } B = \Lambda$$




- Γ_0 assumes all elements of $S(\gamma_1)$ have settled down.
- Possibly there is $x \in S(\gamma_1)$ and $y \in S(\gamma_0)$ such that
 - while $x \in A$, $y \in A$ iff $y \in \Gamma_0^{\Omega_0^A}$, but
 - while $x \notin A$, $y \notin \Gamma_0^{\Omega_0^A}$.
 - $\lim_s A(x)$ does not exist, i.e. A is Σ_2^0 .



S-Requirement - Solution

$$S_{\Psi, \Omega_0, \Omega_1} : A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \text{ or } B = \Lambda$$



- Assume $S(\gamma_1) = \{x\}$.
- Construct two enumeration operators: $\Gamma_{0,0}$ and $\Gamma_{0,1}$.
- $\Gamma_{0,0}$ assumes $x \notin A$ and $\Gamma_{0,1}$ assumes $x \in A$.
- Accounts for Σ_2^0 nature of A .
- In general, if $|S(\gamma_1)| = n$, then we construct 2^n enumeration operators.



Quasi-Lexicographical Ordering

Definition

Define the quasi-lexicographical ordering $<_b$ on $2^{<\omega}$ by $\sigma <_b \tau$ if


1. $|\sigma| < |\tau|$, or
2. $|\sigma| = |\tau|$ and there is a $k < |\sigma|$ such that $\sigma(k) = 0$, $\tau(k) = 1$, and for all $i < k$, $\sigma(i) = \tau(i)$.

0 $<_b$ 1 $<_b$
00 $<_b$ 01 $<_b$ 10 $<_b$ 11 $<_b$
000 $<_b$ 001 $<_b$ 010 $<_b$ 011 $<_b$ 100 $<_b$ 101 $<_b$...



S-Requirement - Bounded


$$S_{\Psi, \Omega_0, \Omega_1} : A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \text{ or } B = \Lambda$$



- The strategy proceeds basically the same as before.
- We have added infinitely many outcomes to the tree to account for all possible states of elements to the left of the current outcome.
- When an element is moved left through the streams, it must make sure that the assumptions of the new stream are consistent with the assumptions of the previous stream.

S-Requirement - Bounded

$$S_{\Psi, \Omega_0, \Omega_1} : A = \Psi^{\Omega_0^A \oplus \Omega_1^A} \Rightarrow A = \Gamma_0^{\Omega_0^A} \text{ or } A = \Gamma_1^{\Omega_1^A} \text{ or } B = \Lambda$$



- Other technical concerns not covered here include:
 - Other uses of conditional dumping.
 - Local approximations to B .
 - etc.



The End!

Kent, Sorbi “Bounding Nonsplitting Enumeration Degrees,”
to appear in The Journal of Symbolic Logic.

